



MATHS

BOOKS - CENGAGE MATHS (ENGLISH)

STRAIGHT LINES

Others

1. If the lines joining the origin and the point of intersection of curves $ax^2 + 2hxy + by^2 + 2gx = 0$ and $a_1x^2 + 2h_1xy + b_1y^2 + 2g_1x = 0$ are mutually perpendicular, then prove that $g(a_1 + b_1) = g_1(a + b)$.

Watch Video Solution

2. Prove that the angle between the lines joining the origin to the points of intersection of the straight line y = 3x + 2 with the curve

$$x^2+2xy+3y^2+4x+8y-11=0 ext{ is } an^{-1}iggl(rac{2\sqrt{2}}{3}iggr)$$

Watch Video Solution

3. Prove that the straight lines joining the origin to the point of intersection of the straight line hx + ky = 2hk and the curve $(x - k)^2 + (y - h)^2 = c^2$ are perpendicular to each other if $h^2 + k^2 = c^2$.

Watch Video Solution

4. If $x^2 - 2pxy - y^2 = 0$ and $x^2 - 2qxy - y^2 = 0$ bisect angles between

each other, then find the condition.

Watch Video Solution

5. Find the value of a for which the lines represented by $ax^2 + 5xy + 2y^2 = 0$ are mutually perpendicular.

6. Find the acute angle between the pair of lines represented by $(x\coslpha-y\sinlpha)^2=ig(x^2+y^2ig)\sin^2lpha.$

Watch Video Solution

7. If the angle between the two lines represented by $2x^2+5xy+3y^2+6x+7y+4=0$ is $an^{-1}(m),$ then find the value of m.

Watch Video Solution

8. If the pair of straight lines $ax^2 + 2hxy + by^2 = 0$ is rotated about the origin through 90^0 , then find the equations in the new position.



9. The orthocentre of the triangle formed by the lines xy=0 and x+y=1 is (a) $\left(rac{1}{2},rac{1}{2}
ight)$ (b) $\left(rac{1}{3},rac{1}{3}
ight)$ (c) (0,0) (d) $\left(rac{1}{4},rac{1}{4}
ight)$

Watch Video Solution

10. The lines joining the origin to the point of intersection of $3x^2 + mxy - 4x + 1 = 0$ and 2x + y - 1 = 0 are at right angles. Then

which of the following is a possible value of m?

Watch Video Solution

11. If the slope of one line is double the slope of another line and the

combined equation of the pair of lines is $\left(\frac{x^2}{a}\right) + \left(\frac{2xy}{h}\right) + \left(\frac{y^2}{b}\right) = 0$, then find the ratio ab : h^2 .



13. The value k for which $4x^2 + 8xy + ky^2 = 9$ is the equation of a pair

of straight lines is_____

Watch Video Solution

14. The two lines represented by $3ax^2 + 5xy + (a^2 - 2)y^2 = 0$ are perpendicular to each other for (a) two values of a (b) a (c) for one value of a (d) for no values of a



15. If two lines represented by $x^4 + x^3y + cx^2y^2 - xy^3 + y^4 = 0$ bisect the angle between the other two, then the value of c is (a) 0 (b) -1 (c) 1 (d) -6

Watch Video Solution

16. The straight lines represented by $x^2 + mxy - 2y^2 + 3y - 1 = 0$ meet at (a) $\left(-\frac{1}{3}, \frac{2}{3}\right)$ (b) $\left(-\frac{1}{3}, -\frac{2}{3}\right)$ (c) $\left(\frac{1}{3}, \frac{2}{3}\right)$ (d) none of these

Watch Video Solution

17. The straight lines represented by the equation $135x^2 - 136xy + 33y^2 = 0$ are equally inclined to the line (a) x - 2y = 7 (b) x+2y=7 (c) x - 2y = 4 (d) 3x + 2y = 4

18. If one of the lines of $my^2+ig(1-m^2ig)xy-mx^2=0$ is a bisector of the angle between the lines xy=0 , then m is (a)1 (b) 2 (c) $-rac{1}{2}$ (d) -1

Watch Video Solution

19. Statement 1 : If -2h = a + b, then one line of the pair of lines $ax^2 + 2hxy + by^2 = 0$ bisects the angle between the coordinate axes in the positive quadrant. Statement 2 : If ax + y(2h + a) = 0 is a factor of $ax^2 + 2hxy + by^2 = 0$, then b + 2h + a = 0

Watch Video Solution

20. Show that all chords of the curve $3x^2 - y^2 - 2x + 4y = 0$, which subtend a right angle at the origin, pass through a fixed point. Find the coordinates of the point.



23. x + y = 7 and $ax^2 + 2hxy + ay^2 = 0$, $(a \neq 0)$, are three real distinct lines forming a triangle. Then the triangle is (a) isosceles (b) scalene equilateral (d) right angled

Watch Video Solution

24. If the slope of one of the lines represented by $ax^2 + 2hxy + by^2 = 0$ is the square of the other, then $\frac{a+b}{h} + \frac{8h^2}{ab} =$ (a) 4 (b) 6 (c) 8 (d) none

of these



25. Find the area of the triangle formed by the line x+y=3 and the angle bisectors of the pair of lines $x^2-y^2+4y-4=0$

Watch Video Solution

26. The sides of a triangle have the combined equation $x^2 - 3y^2 - 2xy + 8y - 4 = 0$. The third side, which is variable, always passes through the point (-5, -1). Find the range of values of the slope of the third line such that the origin is an interior point of the triangle.

27. Let PQR be a right-angled isosceles triangle, right angled at P(2, 1). If the equation of the line QR is 2x + y = 3, then the equation representing the pair of lines PQ and PR is (a) $3x^2 - 3y^2 + 8xy + 20x + 10y + 25 = 0$ (b) $3x^2 - 3y^2 + 8xy - 20x - 10y + 25 = 0$ (c) $3x^2 - 3y^2 + 8xy + 10x + 15y + 20 = 0$ (d) $3x^2 - 3y^2 - 8xy - 15y - 20 = 0$

Watch Video Solution

28. The combined equation of three sides of a triangle is $(x^2 - y^2)(2x + 3y - 6) = 0$. If (-2, a) is an interior point and (b, 1) is an exterior point of the triangle, then `2

Watch Video Solution

29. Find the equation of the bisectors of the angles between the lines joining the origin to the point of intersection of the straight line

x-y=2 with the curve $5x^2+11xy-8y^2+8x-4y+12=0$

Watch Video Solution

30. If θ is the angle between the lines given by the equation $6x^2 + 5xy - 4y^2 + 7x + 13y - 3 = 0$, then find the equation of the line passing through the point of intersection of these lines and making an angle θ with the positive x-axis.

> Watch Video Solution

31. The distance of a point (x_1, y_1) from two straight lines which pass through the origin of coordinates is p. Find the combined equation of these straight lines.

32. Prove that the product of the perpendiculars from (α, β) to the pair

of lines
$$ax^2+2hxy+by^2=0$$
 is $\displaystyle rac{alpha^2+2hlphaeta+beta^2}{\sqrt{\left(a-b
ight)^2+4h^2}}$

Watch Video Solution

33. Find the area enclosed by the graph of $x^2y^2 - 9x^2 - 25y^2 + 225 = 0$.

Watch Video Solution

34. Show that the pairs of straight lines $2x^2 + 6xy + y^2 = 0$ and $4x^2 + 18xy + y^2 = 0$ have the same set of angular bisector.

Watch Video Solution

35. Show that the equation of the pair of lines bisecting the angles between the pair of bisectors of the angles between the pair of lines $ax^2 + 2hxy + by^2 = 0$ is $(a - b)(x^2 - y^2) + 4hxy = 0$.

36. Find the angle between the straight lines joining the origin to the point of intersection of $x^2 + 2xy + 3y^2 + 4x + 8y - 11 = 0$ and 3x - y = -2

Watch Video Solution

37. Through a point A on the x-axis, a straight line is drawn parallel to the y-axis so as to meet the pair of straight lines $ax^2 + 2hxy + by^2 = 0$ at B and C. If AB = BC, then $h^2 = 4ab$ (b) $8h^2 = 9ab$ $9h^2 = 8ab$ (d) $4h^2 = ab$

Watch Video Solution

38. Find the lines whose combined equation is $6x^2 + 5xy - 4y^2 + 7x + 13y - 3 = 0$

39. Does equation $x^2 + 2y^2 - 2\sqrt{3}x - 4y + 5 = 0$ satisfies the condition $abc + 2gh - af^2 - bg^2 - ch^2 = 0$? Does it represent a pair of straight lines?

Watch Video Solution

40. Find the value of λ if $2x^2 + 7xy + 3y^2 + 8x + 14y + \lambda = 0$

represents a pair of straight lines

41. Find the distance between the pair of parallel lines $x^2 + 4xy + 4y^2 + 3x + 6y - 4 = 0$

42. If the pair of lines $ax^2+2hxy+by^2+2gx+2fy+c=0$ intersect on the y-axis , then prove that $2fgh=bg^2+ch^2.$

43. Find the lines whose combined equation is $6x^2 + 5xy - 4y^2 + 7x + 13y - 3 = 0$ using the concept of parallel lines through the origin.

Watch Video Solution

44. If the lines $px^2 - qxy - y^2 = 0$ makes the angles lpha and eta with X-axis

, then the value of an(lpha+eta) is

45. Find the joint equation of the pair of lines which pass through the origin and are perpendicular to the lines represented the equation $y^2 + 3xy - 6x + 5y - 14 = 0$



46. If the sum of the slopes of the lines given by $x^2 - 2cxy - 7y^2 = 0$ is

four times their product, then the value of c is_____

Watch Video Solution

47. The distance between the two lines represented by the equation

$$9x^2-24xy+16y^2-12x+16y-12=0$$
 is (a) $rac{8}{5}$ (b) $rac{6}{5}$ (c) $rac{11}{5}$ (d)

none of these

48. If the gradient one of the lines $ax^2 + 2hxy + by^2 = 0$ is twice that of

the other, then h^2 =____`

Watch Video Solution

49. If one of the lines of $my^2 + (1 - m^2)xy - mx^2 = 0$ is a bisector of the angle between the lines xy = 0 , then m is 3 (b) 2 (c) $-rac{1}{2}$ (d) -1

Watch Video Solution

50. Two pairs of straight lines have the equations $y^2 + xy - 12x^2 = 0$ and $ax^2 + 2hxy + by^2 = 0$. One line will be common among them if. a + 8h - 16b = 0 (b) a - 8h + 16b = 0 a - 6h + 9b = 0 (d) a + 6h + 9b = 0

51. If the equation of the pair of straight lines passing through the point (1, 1), one making an angle θ with the positive direction of the x-axis and the other making the same angle with the positive direction of the y-axis, is $x^2 - (a+2)xy + y^2 + a(x+y-1) = 0, a \neq 2$, then the value of $\sin 2\theta$ is

(a)
$$a-2$$
 (b) $a+2$ (c) $\displaystyle rac{2}{a+2}$ (d) $\displaystyle rac{2}{a}$

Watch Video Solution

52. If one of the lines given by the equation $2x^2 + pxy + 3y^2 = 0$ coincide with one of those given by $2x^2 + qxy - 3y^2 = 0$ and the other lines represented by them are perpendicular, then p = 5 (b) p = -5 q = -1 (d) q = 1

Watch Video Solution

53. If $x^2+2hxy+y^2=0$ represents the equation of the straight lines through the origin which make an angle lpha with the straight line

$$y + x = 0$$
 (a) $sec2lpha = h \cos lpha$ (b) $= \sqrt{rac{(1+h)}{(2h)}}$ (c) $2\sin lpha = \sqrt{rac{(1+h)}{h}}$ (d) $\cot lpha = \sqrt{rac{(1+h)}{(h-1)}}$

Watch Video Solution

54. The equation to a pair of opposite sides of a parallelogram are $x^2 - 5x + 6 = 0$ and $y^2 - 6y + 5 = 0$. The equations to its diagonals are x + 4y = 13, y = 4x - 7 (b) 4x + y = 13, 4y = x - 74x + y = 13, y = 4x - 7 (d) y - 4x = 13, y + 4x - 7

Watch Video Solution

55. The equation $a^2x^2 + 2h(a+b)xy + b^2y^2 = 0$ and $ax^2 + 2hxy + by^2 = 0$ represent two pairs of perpendicular straight lines two pairs of parallel straight lines two pairs of straight lines which are equally inclined to each other none of these

56. The equation $x^3 + x^2y - xy^2 = y^3$ represents (a)three real straight lines (b)lines in which two of them are perpendicular to each other (c)lines in which two of them are coincident (d)none of these

Watch Video Solution

57. The image of the pair of lines represented by $ax^2 + 2hxy + by^2 = 0$ by the line mirror y = 0 is $ax^2 - 2hxy - by^2 = 0$ $bx^2 - 2hxy + ay^2 = 0$ $bx^2 + 2hxy + ay^2 = 0$ $ax^2 - 2hxy + by^2 = 0$

Watch Video Solution

58. The combined equation of the lines l_1andl_2 is $2x^2 + 6xy + y^2 = 0$ and that of the lines m_1andm_2 is $4x^2 + 18xy + y^2 = 0$. If the angle between l_1 and m_2 is α then the angle between l_2andm_1 will be

59. If the equation $ax^2 - 6xy + y^2 = 0$ represents a pair of lines whose slopes are m and m^2 , then the value(s) of a is/are

Watch Video Solution

60. The equation of a line which is parallel to the line common to the pair of lines given by $6x^2 - xy - 12y^2 = 0$ and $15x^2 + 14xy - 8y^2 = 0$ and at a distance of 7 units from it is 3x - 4y = -35 5x - 2y = 73x + 4y = 35 2x - 3y = 7

Watch Video Solution

61. If the sum of the slopes of the lines given by $x^2 - 2cxy - 7y^2 = 0$ is

four times their product, then the value of c is_____

62. Area of the triangle formed by the line x + y = 3 and the angle bisectors of the pairs of straight lines $x^2 - y^2 + 2y = 1$ is 2squnits (b) 4squnits 6squnits (d) 8squnits

Watch Video Solution

63. The equation $x^2y^2 - 9y^2 - 6x^2y + 54y = 0$ represents (a) a pair of straight lines and a circle (b) a pair of straight lines and a parabola (c) a set of four straight lines forming a square (d) none of these

Watch Video Solution

64. The straight lines represented by $(y - mx)^2 = a^2(1 + m^2)$ and $(y - nx)^2 = a^2(1 + n^2)$ from a (a) rectangle (b) rhombus (c) trapezium (d) none of these

65. If the pairs of lines $x^2 + 2xy + ay^2 = 0$ and $ax^2 + 2xy + y^2 = 0$ have exactly one line in common, then the joint equation of the other two lines is given by (1) $3x^2 + 8xy - 3y^2 = 0$ (2) $3x^2 + 10xy + 3y^2 = 0$ (3) $y^2 + 2xy - 3x^2 = 0$ (4) $x^2 + 2xy - 3y^2 = 0$

Watch Video Solution

66. The condition that one of the straight lines given by the equation $ax^2 + 2hxy + by^2 = 0$ may coincide with one of those given by the equation $a'x^2 + 2h'xy + b'y^2 = 0$ is $(ab' - a'b)^2 = 4(ha' - h'a)(bh' - b'h)$ $(ab' - a'b)^2 = (ha' - h'a)(bh' - b'h)$ $(ha' - h'a)^2 = 4(ab' - a'b)(bh' - b'h)$ $(bh' - b'h)^2 = 4(ab' - a'b)(ha' - h'a)$

67. If the represented by the equation $3y^2 - x^2 + 2\sqrt{3}x - 3 = 0$ are rotated about the point $(\sqrt{3}, 0)$ through an angle of 15^0 , on in clockwise direction and the other in anticlockwise direction, so that they become perpendicular, then the equation of the pair of lines in the new position is (1) $y^2 - x^2 + 2\sqrt{3}x + 3 = 0$ (2) $y^2 - x^2 + 2\sqrt{3}x - 3 = 0$ (3) $y^2 - x^2 - 2\sqrt{3}x + 3 = 0$ (4) $y^2 - x^2 + 3 = 0$

Watch Video Solution

68. A point moves so that the distance between the foot of perpendiculars from it on the lines $ax^2 + 2hxy + by^2 = 0$ is a constant 2d. Show that the equation to its locus is $(x^2 + y^2)(h^2 - ab) = d^2 \{(a - b)^2 + 4h^2\}$.

Watch Video Solution

69. The angle between the pair of lines whose equation is $4x^2 + 10xy + my^2 + 5x + 10y = 0$ is

(a)
$$an^{-1}igg(rac{3}{8}igg)$$
 (b) $an^{-1}igg(rac{3}{4}igg)$ (c) $an^{-1}igg\{2rac{\sqrt{25-4m}}{m+4}igg\}, m\in R$ (d)

none of these



70. Find the point of intersection of the pair of straight lines represented

by the equation $6x^2 + 5xy - 21y^2 + 13x + 38y - 5 = 0$.





Watch Video Solution

72. If the pair of lines $\sqrt{3}x^2 - 4xy + \sqrt{3}y^2 = 0$ is rotated about the origin by $\frac{\pi}{6}$ in the anticlockwise sense, then find the equation of the pair in the new position.

73. If the equation $2x^2 + kxy + 2y^2 = 0$ represents a pair of real and distinct lines, then find the values of k.

Watch Video Solution

74. If the equation $x^2+(\lambda+\mu)xy+\lambda uy^2+x+\mu y=0$ represents two parallel straight lines, then prove that $\lambda=\mu.$

Watch Video Solution

75. If one of the lines of the pair $ax^2 + 2hxy + by^2 = 0$ bisects the angle between the positive direction of the axes. Then find the relation for a, band h.

76. Prove that the equation $2x^2 + 5xy + 3y^2 + 6x + 7y + 4 = 0$ represents a pair of straight lines. Find the coordinates of their point of intersection and also the angle between them.

77. A line L passing through the point (2, 1) intersects the curve $4x^2 + y^2 - x + 4y - 2 = 0$ at the point AandB. If the lines joining the origin and the points A, B are such that the coordinate axes are the bisectors between them, then find the equation of line L.



79. If one of the lines denoted by the line pair $ax^2 + 2hxy + by^2 = 0$ bisects the angle between the coordinate axes, then prove that $(a+b)^2 = 4h^2$

Watch Video Solution

80. If the middle points of the sides BC, CA, and AB of triangle ABC are (1, 3), (5, 7), and (-5, 7), respectively, then find the equation of the side AB.

Watch Video Solution

81. Find the equations of the lines which pass through the origin and are

inclined at an angle $\tan^{-1} m$ to the line y = mx + c.

82. If (-2,6) is the image of the point (4,2) with respect to line L=0, then L

is:



83. If the lines x+(a-1)y+1=0 and $2x+a^2y-1=0$ are perpendicular, then find the value of a.

Watch Video Solution

84. Find the equation of the right bisector of the line segment joining the

points (3, 4) and (-1, 2).



85. If the coordinates of the points A, B, C and D be (a, b), (a', b'), (-a, b) and (a', -b') respectively, then the equation





89. Find the equation of the straight line passing through the intersection of the lines x - 2y = 1 and x + 3y = 2 and parallel to 3x + 4y = 0.



concurrent.

Watch Video Solution

91. If the point $P(a, a^2)$ lies completely inside the triangle formed by the lines x = 0, y = 0, and x + y = 2, then find the exhaustive range of values of a is (A) (0, 1) (B) $(1, \sqrt{2})$ (C) $(\sqrt{2} - 1, 1)$ (D) $(\sqrt{2} - 1, 2)$

92. If the point (a, a) is placed in between the lines |x + y| = 4, then find the values of a_{\cdot}



94. If the point $P(a^2, a)$ lies in the region corresponding to the acute angle between the lines 2y = x and 4y = x, then find the values of a.

Watch Video Solution

95. Find the range of values of the ordinate of a point moving on the line

x=1 , which always remain in the interior of the triangle formed by the

lines y = x, the x-axis and x + y = 4.



96. The point (8, -9) with respect to the lines 2x + 3y - 4 = 0 and 6x + 9y + 8 = 0 lies on (a) the same side of the lines (b) the different sides of the line (c)one of the line (d) none of these



97. If the point $(a^2, a+1)$ lies in the angle between the lines 3x-y+1=0 and x+2y-5=0 containing the origin, then find the value of a.

Watch Video Solution

98. Find the range of (lpha,2+lpha) and $\left(rac{3lpha}{2},a^2
ight)$ lie on the opposite sides

of the line 2x + 3y = 6.

99. Which pair of points lies on the same side of 3x - 8y - 7 = 0? a) (0, -1)and(0, 0) b)(4, -3) and (0, 1) c) (-3, -4)and(1, 2) d) (-1, -1)and(3, 7)

Watch Video Solution

100. The line $\frac{x}{a} + \frac{y}{b} = 1$ moves in such a way that $\frac{1}{a^2} + \frac{1}{b^2} = \frac{1}{c^2}$, where c is a constant. The locus of the foot of the perpendicular from the origin on the given line is $x^2 + y^2 = c^2$.

Watch Video Solution

101. Consider the lines given by $L_1: x + 3y - 5 = 0$ $L_2: 3x - ky - 1 = 0$ $L_3: 5x + 2y - 12 = 0$ Column I|Column II L_1, L_2, L_3 are concurrent if|p. k = -9 One of L_1, L_2, L_3 is parallel to at least one of the other two if|q. $k=-rac{6}{5}$ L_1,L_2,L_3 form a triangle if|r. $k=rac{5}{6}$ L_1,L_2,L_3 do not

form a triangle if s. k = 5

Watch Video Solution

102. A variable straight line is drawn through the point of intersection of the straight lines $\frac{x}{a} + \frac{y}{b} = 1$ and $\frac{x}{b} + \frac{y}{a} = 1$ and meets the coordinate axes at A and B. Show that the locus of the midpoint of ABis the curve 2xy(a + b) = ab(x + y)

Watch Video Solution

103. The line 3x + 2y = 24 meets the y-axis at A and the x-axis at B. The perpendicular bisector of AB meets the line through (0, -1) parallel to the x-axis at C. If the area of triangle ABC is A, then the value of $\frac{A}{13}$ is

104. Find equation of the line passing through the point (2, 2) and cutting

off intercepts on the axes whose sum is 9.



105. The area of the parallelogram formed by the lines

$$y = mx, y = xm + 1, y = nx, andy = nx + 1$$
 equals. $\frac{|m + n|}{(m - n)^2}$ (b)
 $\frac{2}{|m + n|} \frac{1}{(|m + n|)}$ (d) $\frac{1}{(|m - n|)}$
Watch Video Solution

106. A ray of light is sent along the line 2x - 3y = 5. After refracting across the line x + y = 1 it enters the opposite side after torning by 15^0 away from the line x + y = 1. Find the equation of the line along which the refracted ray travels.
107. Let $P = (-1, 0), Q = (0, 0) and R = (3, 3\sqrt{3})$ be three points. The equation of the bisector of the angle PQR (1) $\sqrt{3}x + y = 0$ (2) $x + \frac{\sqrt{3}}{2}y = 0$ (3) $\frac{\sqrt{3}}{2}x + y = 0$ (4) $x + \sqrt{3}y = 0$

Watch Video Solution

108. A ray of light is sent along the line x - 2y - 3 = 0 upon reaching the line 3x - 2y - 5 = 0, the ray is reflected from it. Find the equation of the line containing the reflected ray.

Watch Video Solution

109. Line *L* has intercepts *a* and *b* on the coordinate axes. When the axes are rotated through a given angle keeping the origin fixed, the same line *L* has intercepts *p* and *q*. Then (a) $a^2 + b^2 = p^2 + q^2$ (b) $\frac{1}{a^2} + \frac{1}{b^2} = \frac{1}{p^2} + \frac{1}{q^2}$ (c) $a^2 + p^2 = b^2 + q^2$ (d) $\frac{1}{a^2} + \frac{1}{p^2} = \frac{1}{b^2} + \frac{1}{q^2}$

110. If the sum of the distances of a point from two perpendicular lines in a plane is 1, then its locus is(A) a square(b) a circle (C) a straight line(d) two intersecting lines

111. A line 4x + y = 1 passes through the point A(2,-7) and meets line BC at B whose equation is 3x - 4y + 1 = 0, the equation of line AC such that AB = AC is (a) 52x +89y +519=0(b) 52x +89y-519=0 c) 82x +52y+519=0 (d) 89x +52y -519=0

Watch Video Solution

112. A straight canal is $4\frac{1}{2}$ miles from a place and the shortest route from this place to the canal is exactly north-east. A village is 3miles north and four miles east from the place. Does it lie by the nearest edge of the canal?

113. Let PS be the median of the triangle with vertices P(2, 2), Q(6, -1)andR(7, 3) Then equation of the line passing through (1, -1) and parallel to PS is 2x - 9y - 7 = 02x - 9y - 11 = 0 2x + 9y - 11 = 0 2x + 9y + 7 = 0

Watch Video Solution

114. Find the equation of the line perpendicular distance from the origin is 5 units and the angle made by the perpendicular with the positive x-axis is $30 \circ$.

115. The number of integral values of m for which the x-coordinate of the point of intersection of the lines 3x + 4y = 9 and y = mx + 1 is also an integer is (a) -2 (b) 0 (c) 4 (d) 1



116. Reduce the line 2x - 3y + 5 = 0 in slope-intercept, intercept, and normal forms.

Watch Video Solution

117. The line 5x + 4y = 0 passes through the point of intersection of

straight lines x+2y-10 = 0, 2x + y =-5

Watch Video Solution

118. If the intercept of a line between the coordinate axes is divided by the

point (-5, 4) in the ratio 1:2, then find the equation of the line.

119. The lines 2x + 3y + 19 = 0 and 9x + 6y - 17 = 0 , cut the

coordinate axes at concyclic points.



120. The straight lines 3x + y - 4 = 0, x + 3y - 4 = 0 and x + y = 0 form a triangle which is :

Watch Video Solution

121. A line through the variable point A(k+1,2k) meets the lines

7x + y - 16 = 0, 5x - y - 8 = 0, x - 5y + 8 = 0 at B, C, D, T

respectively. Prove that AC, AB, AD are in HP.

122. Two particles start from point (2, -1), one moving two units along the line x + y = 1 and the other 5 units along the line x - 2y = 4, If the particle move towards increasing y, then their new positions are:



123. If P=(1,0); Q=(-1.0)& R=(2,0) are three given points, then

the locus of the points S satisfying the relation, $SQ^2+SR^2=2SP^2$ is -

Watch Video Solution

124. Distance of point (1,3) from the line 2x - 3y + 9 = 0 along

x - y + 1 = 0

125. A rectangle ABCD has its side AB parallel to line y = x, and vertices A, BandD lie on y = 1, x = 2, and x = -2, respectively. The locus of vertex C is x = 5 (b) x - y = 5 y = 5 (d) x + y = 5

126. Two adjacent vertices of a square are (1, 2) and (-2, 6) Find the other vertices.

Watch Video Solution

127. The equation of a line through the point (1, 2) whose distance from

the point (3,1) has the greatest value is (a)y=2x (b) y=x+1 (c)

$$x+2y=5$$
 (d) $y=3x-1$

128. Find the equation of the line through the point A(2,3) and making an angle of 45 \circ with the x axis Also determine the length of intercept on it between A and the line x+y+1=0

Watch Video Solution

129. The line $\frac{x}{a} + \frac{y}{b} = 1$ meets the x-axis at A, the y-axis at B, and the line y = x at C such, that the area of DeltaAOC is twice the area of DeltaBOC. Then the coordinates of C are $\left(\frac{b}{3}, \frac{b}{3}\right)$ (b) $\left(\frac{2a}{3}, \frac{2a}{3}\right)$ (c) $\left(\frac{2b}{3}, \frac{2b}{3}\right)$ (d) none of these

Watch Video Solution

130. The line joining two points A(2,0) and B(3,1) is rotated about A in anticlockwise direction through an angle of 15° . find the equation of line in the new position. If b goes to c in the new position what will be the coordinates of C.



131. The area of the triangle formed by the lines y = ax, x + y - a = 0

and the y-axis is (a)
$$rac{1}{2|1+a|}$$
 (b) $rac{1}{|1+a|}$ (c) $rac{1}{2}\Big|rac{a}{1+a}\Big|$ (d) $rac{a^2}{2|1+a|}$

Watch Video Solution

132. Find the equation of the lines through the point (3, 2) which make an

angle of 45 with the line x - 2y = 3 .

Watch Video Solution

133. Consider the points A(0, 1)andB(2, 0), andP be a point on the line 4x + 3y + 9 = 0. The coordinates of P such that |PA - PB| is maximum are $\left(-\frac{24}{5}, \frac{17}{5}\right)$ (b) $\left(-\frac{84}{5}, \frac{13}{5}\right)\left(\frac{31}{7}, \frac{31}{7}\right)$ (d) (0, 0)

134. A straight line is drawn through the point P(2,3)and is inclined at an angle of 30° with the x axis. Then the coordinates of two points on it at a distance 4 from P on either side of P will be..

135. A line of fixed length 2 units moves so that its ends are on the positive x-axis and that part of the line x + y = 0 which lies in the second quadrant. Then the locus of the midpoint of the line has equation.

(a)
$$x^2 + 5y^2 + 4xy - 1 = 0$$
 (b) $x^2 + 5y^2 + 4xy + 1 = 0$ (c)

$$x^2+5y^2-4xy-1=0$$
 (d) $4x^2+5y^2+4xy+1=0$

Watch Video Solution

136. The perpendicular from the origin to a line meets it at the point (2, 9), find the equation of the line.



137. The line $\frac{x}{3} + \frac{y}{4} = 1$ meets the y-axis and x-axis at A and B respectively. A square *ABCD* is constructed on the line segment *AB* away from the origin. The coordinates of the vertex of the square farthest from the origin are

A. (a) (7,3)

B. (b) (4,7)

C. (c) (6,4)

D. (d) (3,8)

Answer: null

Watch Video Solution

138. Find the direction in which a straight line must be drawn through the point (1, 2) so that its point of intersection with the line x + y = 4 may be at a distance of 3 units from this point.

139. The centroid of an equilateral triangle is (0, 0). If two vertices of the triangle lie on $x + y = 2\sqrt{2}$, then one of them will have its coordinates. (a) $(\sqrt{2} + \sqrt{6}, \sqrt{2} - \sqrt{6})$ (b) $(\sqrt{2} + \sqrt{3}, \sqrt{2} - \sqrt{3})$ (c) $(\sqrt{2} + \sqrt{5}, \sqrt{2} - \sqrt{5})$ (d) none of these

Watch Video Solution

140. Two fixed points A and B are taken on the coordinates axes such that OA=a and OB =b. Two variable points A' and B' are taken on the same axes such that OA'+OB' = OA+OB. Find the locus of the point of intersection of AB' and A'B.



141. Find the equations of the lines, which cut-off intercepts on the axes

whose sum and product are 1 and -6 , respectively.

142. Find the equation of the straight line which passes through the origin and makes angle 60^0 with the line $x + \sqrt{3}y + \sqrt{3} = 0$.

143. The equation of a straight line passing through the point (2, 3) and inclined at an angle of $\tan^{-1}\left(\frac{1}{2}\right)$ with the line y + 2x = 5 is: (a)y = 3 (b) x = 2 (c)3x + 4y - 18 = 0 (d) 4x + 3y - 17 = 0

Watch Video Solution

144. If we reduce 3x + 3y + 7 = 0 to the form $x \cos \alpha + y \sin \alpha = p$,

then find the value of p.

145. The equation of line on which the perpendicular from the origin make 30^0 angle with the x-axis and which form a triangle of area $\frac{50}{\sqrt{3}}$ with the axes is a. $\sqrt{3}x + y - 10 = 0$ b. $\sqrt{3}x + y + 10 = 0$ c. $x + \sqrt{3}y - 10 = 0$ d. $x - \sqrt{3}y - 10 = 0$

Watch Video Solution

146. In the given figure, PQR is an equilateral triangle and OSPT is square. If $OT = 2\sqrt{2}$ units, find the equation of lines OT, OS, SP, QR, PR, and PQ.

Watch Video Solution

147. Line *L* has intercepts *aandb* on the coordinate axes. When the axes are rotated through a given angle keeping the origin fixed, the same line *L* has intercepts *pandq*. Then (a) $a^2 + b^2 = p^2 + q^2$ (b) $\frac{1}{a^2} + \frac{1}{b^2} = \frac{1}{p^2} + \frac{1}{q^2}$ (c) $a^2 + p^2 = b^2 + q^2$ (d) $\frac{1}{a^2} + \frac{1}{p^2} = \frac{1}{b^2} + \frac{1}{q^2}$

148. A line intersects the straight lines 5x-y-4=0 and 3x-4y-4=0 at A and B, respectively. If a point P(1,5) on the line AB is such that AP : PB = 2:1(internally), find point A.

Watch Video Solution

149. A line is a drawn from P(4, 3) to meet the lines L_1 and l_2 given by 3x + 4y + 5 = 0 and 3x + 4y + 15 = 0 at points A and B respectively. From A, a line perpendicular to L is drawn meeting the line L_2 at A_1 Similarly, from point B_1 Thus a parallelogram $\forall_1 BB_1$ is formed. Then the equation of L so that the area of the parallelogram $\forall_1 BB_1$ is the least is (a) x - 7y + 17 = 0 (b) 7x + y + 31 = 0 (c) x - 7y - 17 = 0 (d) x + 7y - 31 = 0

150. A straight line through the point A (3,4) is such that its intercept between the axis is bisected at A then its equation is : A. x + y = 7 B. 3x - 4y + 7 = 0 C. 4x + 3y = 24 D. 3x + 4y = 24

Watch Video Solution

151. Two straight lines u = 0 and v = 0 pass through the origin and the angle between them is $\tan^{-1}\left(\frac{7}{9}\right)$. If the ratio of the slope of v = 0 and u = 0 is $\frac{9}{2}$, then their equations are (a) y + 3x = 0 and 3y + 2x = 0 (b) 2y + 3x = 0 and 3y + x = 0 (c) 2y = 3x and 3y = x (d) y = 3x and 3y = 2x

Watch Video Solution

152. A straight line through the point (2, 2) intersects the lines $\sqrt{3}x + y = 0$ and $\sqrt{3}x - y = 0$ at the point A and B, respectively. Then find the equation of the line AB so that triangle OAB is equilateral.

153. Let $u\equiv ax+by+ab3=0, v\equiv bx-ay+ba3=0, a,b\in R$, be two straight lines. The equations of the bisectors of the angle formed by $k_1u-k_2v=0$ and $k_1u+k_2v=0$, for nonzero and real k_1 and k_2 are u=0 (b) $k_2u+k_1v=0$ $k_2u-k_1v=0$ (d) v=0

Watch Video Solution

154. If the foot of the perpendicular from the origin to a straight line is at

 $(3, \ -4)$, then find the equation of the line.

Watch Video Solution

155. Two sides of a triangle are parallel to the coordinate axes. If the slopes of the medians through the acute angles of the triangle are 2 and m, then m is (a) $\frac{1}{2}$ (b) 2 (c) 4 (d) 8

156. The diagonals AC and BD of a rhombus intersect at (5, 6). If A = (3, 2), then find the equation of diagonal BD.

157. A line which makes an acute angle θ with the positive direction of the x-axis is drawn through the point P(3, 4) to meet the line x = 6 at R and y = 8 at S. Then, (a) $PR = 3 \sec \theta$ (b) $PS = 4 \csc ec\theta$ (c) $PR + PS = \frac{2(3 \sin \theta + 4 \cos \theta)}{\sin 2\theta}$ (d) $\frac{9}{(PR)^2} + \frac{16}{(PS)^2} = 1$

Watch Video Solution

158. Find the values of non-negative real number h_1 , h_2 , h_3 , k_1 , k_2 , k_3 such that the algebraic sum of the perpendiculars drawn from the points $(2, k_1), (3, k_2), \cdot 7, k_3), (h_1, 4), (h_2, 5), (h_3, -3)$ on a variable line passing through (2, 1) is zero.

159. The sides of a triangle ABC lie on the lines 3x + 4y = 0, 4x + 3y = 0and x = 3. Let (h, k) be the centre of the circle inscribed in riangle ABC. The value of (h + k) equals

Watch Video Solution

160. If a and b are two arbitrary constants, then prove that the straight line (a - 2b)x + (a + 3b)y + 3a + 4b = 0 will pass through a fixed point. Find that point.

Watch Video Solution

161. Statement 1: The incenter of a triangle formed by the lines

$$x\cos\left(\frac{\pi}{9}\right) + y\sin\left(\frac{\pi}{9}\right) = \pi, x\cos\left(\frac{8\pi}{9}\right) + y\sin\left(\frac{8\pi}{9}\right) = \pi$$
 and

 $x\cos\left(rac{13\pi}{9}
ight)+y\sin\left(rac{13\pi}{9}
ight)=\pi$ is (0,0) Statement 2: Any point

equidistant from the given three non-concurrent straight lines in the plane is the incenter of the triangle formed by these lines.



163. The lines x + y - 1 = 0, $(m - 1)x + (m^2 - 7)y - 5 = 0$, and (m - 2)x + (2m - 5)y = 0 are a) concurrent for three values of m b) concurrent for one value of m c) concurrent for no value of m d) parallel for m = 3.

164. In triangle ABC , the equation of the right bisectors of the sides AB and AC are x+y=0 and y-x=0 , respectively. If $A\equiv(5,7)$, then find the equation of side BC.

Watch Video Solution

165. If
$$\left(\frac{x}{a}\right) + \left(\frac{y}{b}\right) = 1$$
 and $\left(\frac{x}{c}\right) + \left(\frac{y}{d}\right) = 1$ intersect the axes at four concylic points and $a^2 + c^2 = b^2 + d^2$, then these lines can intersect at, $(a, b, c, d > 0) (1, 1)$ (b) $(1, -1) (2, -2)$ (d) $(3, 3)$

Watch Video Solution

166. Show that the straight lines given by x(a + 2b) + y(a + 3b) = a for

different values of a and b pass through a fixed point.

167. The straight line 3x + 4y - 12 = 0 meets the coordinate axes at AandB . An equilateral triangle ABC is constructed. The possible coordinates of vertex C (a) $\left(2\left(1 - \frac{3\sqrt{3}}{4}\right), \frac{3}{2}\left(1 - \frac{4}{\sqrt{3}}\right)\right)$ (b) $\left(-2(1+\sqrt{3}), \frac{3}{2}(1-\sqrt{3})\right)$ (c) $\left(2(1+\sqrt{3}), \frac{3}{2}(1+\sqrt{3})\right)$ (d) $\left(2\left(1 + \frac{3\sqrt{3}}{4}\right), \frac{3}{2}\left(1 + \frac{4}{\sqrt{3}}\right)\right)$

Watch Video Solution

168. Let ax + by + c = 0 be a variable straight line, whre a, bandc are the 1st, 3rd, and 7th terms of an increasing AP, respectively. Then prove that the variable straight line always passes through a fixed point. Find that point.

169. Angle made with the x-axis by a straight line drawn through (1, 2) so that it intersects x + y = 4 at a distance $\frac{\sqrt{6}}{3}$ from (1, 2) is 105° (b) 75°

Watch Video Solution

170. Prove that all the lines having the sum of the interceps on the axes equal to half of the product of the intercepts pass through the point. Find the fixed point.



171. Given three straight lines 2x + 11y - 5 = 0, 24x + 7y - 20 = 0,

and 4x - 3y - 2 = 0. Then, a)they form a triangle b)one line bisects the

angle between the other two c) two of them are parallel



172. Find the straight line passing through the point of intersection of

2x + 3y + 5 = 0, 5x - 2y - 16 = 0, and through the point (-1, 3).

173. The lines x + 2y + 3 = 0, x + 2y - 7 = 0, and2x - y - 4 = 0 are the sides of a square. The equation of the remaining side of the square can be (a) 2x - y + 6 = 0 (b) 2x - y + 8 = 0 (c) 2x - y - 10 = 0 (d) 2x - y - 14 = 0

Watch Video Solution

174. Consider a family of straight lines $(x+y)+\lambda(2x-y+1)=0$. Find the equation of the straight line belonging to this family that is farthest from (1, -3).

Watch Video Solution

175. Determine all the values of α for which the point (α, α^2) lies inside the triangle formed by the lines. 2x + 3y - 1 = 0 x + 2y - 3 = 05x - 6y - 1 = 0 176. If 5a + 5b + 20c = t, then find the value of t for which the line ax + by + c - 1 = 0 always passes through a fixed point.

Watch Video Solution

177. If the y = mx + 1, of the circle $x^2 + y^2 = 1$ subtends an angle of measure 45° of the major segment of the circle then value of m is -

Watch Video Solution

178. If $\frac{x}{l} + \frac{y}{m} = 1$ is any line passing through the intersection point of the lines $\frac{x}{a} + \frac{y}{b} = 1$ and $\frac{x}{b} + \frac{y}{a} = 1$ then prove that $\frac{1}{l} + \frac{1}{m} = \frac{1}{a} + \frac{1}{b}$

179. Two sides of a rhombus OABC (lying entirely in first quadrant or fourth quadrant) of area equal to 2 sq. units, are $y = \frac{x}{\sqrt{3}}$, $y = \sqrt{3}x$ Then possible coordinates of B is/are (O being the origin).

180. The equation of straight line belonging to both the families of lines $(x - y + 1) + \lambda_1(2x - y - 2) = 0$ and $(5x + 3y - 2) + \lambda_2(3x - y - 4) = 0$ where λ_1, λ_2 are arbitrary numbers is (A) 5x - 2y - 7 = 0 (B)2x + 5y - 7 = 0 (C) 5x + 2y - 7 = 0(D) 2x - 5y - 7 = 0

Watch Video Solution

181. If m_1 and m_2 are the roots of the equation $x^2 - ax - a - 1 = 0$, then the area of the triangle formed by the three straight lines $y = m_1 x, y = m_2 x$, and $y = a(a \neq -1)$ is 1. $\frac{a^2(a+2)}{2(a+1)}$ if a > -1

Watch Video Solution

182. If the algebraic sum of the distances of a variable line from the points (2, 0), (0, 2), and (-2, -2) is zero, then the line passes through the fixed point. (a) (-1, -1) (b) (0, 0) (c) (1, 1) (d) (2, 2)

Watch Video Solution

183. If the points
$$\left(\frac{a^3}{(a-1)}\right)$$
, $\left(\frac{(a^2-3)}{(a-1)}\right)$, $\left(\frac{b^3}{b-1}\right)$, $\left(\frac{b^2-3}{(b-1)}\right)$, $\left(\frac{c^3}{(b-1)}\right)$, and $\left(\frac{(c^2-3)}{(c-1)}\right)$, where a, b, c are different from 1, lie on the $lx + my + n = 0$, then (a) $a + b + c = -\frac{m}{l}$ (b) $ab + bc + ca = \frac{n}{l}$ (c) $abc = \frac{(m+n)}{l}$ (d) $abc - (bc + ca + ab) + 3(a + b + c) = 0$

184. If a, b, c are in harmonic progression, then the straight line $\left(\left(\frac{x}{a}\right)\right) + \left(\frac{y}{b}\right) + \left(\frac{l}{c}\right) = 0$ always passes through a fixed point. Find that point.

Watch Video Solution

185. A variable line cuts n given concurrent straight lines at $A_1, A_2...A_n$

such that $\sum_{i=1}^{n} \frac{1}{OA_{i}}$ is a constant. Show that it always passes through a

fixed point, O being the point of intersection of the lines

Watch Video Solution

186. Prove that the area of the parallelogram formed by the lines

 $3x - 4y + a = 0, \; 3x - 4y + 3a = 0, \; 4x - 3y - a = 0 \; and \; 4x - 3y - 2a = 0$

187. Two sides of a rhombus lying in the first quadrant are given by 3x - 4y = 0 and 12x - 5y = 0. If the length of the longer diagonal is 12, then find the equations of the other two sides of the rhombus.

188. The equation of straight line passing through (-2,-7) and having an intercept of length 3 between the straight lines : 4x + 3y = 12, 4x + 3y = 3 are : (A) 7x + 24y + 182 = 0 (B) 7x + 24y + 18 = 0 (C) x + 2 = 0 (D) x - 2 = 0



189. Let ABC be a given isosceles triangle with AB = AC. Sides ABandAC are extended up to EandF, respectively, such that $BExCF = AB^2$. Prove that the line EF always passes through a fixed point.

190. ABC is an equilateral triangle with A(0, 0) and B(a, 0), (a>0). L, M and N are the foot of the perpendiculars drawn from a point P to the side AB, BC, andCA, respectively. If P lies inside the triangle and satisfies the condition $PL^2 = PM\dot{P}N$, then find the locus of P.

Watch Video Solution

191. Let $L_1 = 0$ and $L_2 = 0$ be two fixed lines. A variable line is drawn through the origin to cut the two lines at R and $S\dot{P}$ is a point on the line AB such that $\frac{(m+n)}{OP} = \frac{m}{OR} + \frac{n}{OS}$. Show that the locus of P is a straight line passing through the point of intersection of the given lines R, S, R are on the same side of O).

Watch Video Solution

192. Find the points on y-axis whose perpendicular distance from the line

4x - 3y - 12 = 0 is 3.

193. Find all the values of θ for which the point $(\sin^2 \theta, \sin \theta)$ lies inside

the square formed by the line xy = 0 and 4xy - 2x - 2y + 1 = 0.

194. If p and q are the lengths of perpendiculars from the origin to the

lines $x\cos heta-y\sin heta=k\cos2 heta$ and $x\sec heta+y\ \csc heta=k$,

respectively, prove that $p^2+4q^2=k^2$.

Watch Video Solution

195. The equations of two sides of a triangle are 3y - x - 2 = 0 and y + x - 2 = 0. The third side, which is variable, always passes through the point (5, -1). Find the range of the values of the slope of the third side, so that the origin is an interior point of the triangle.



196. Prove that the lengths of the perpendiculars from the points $(m^2,2m), (mm',m+m'),$ and $(m^{\,'2},2m')$ to the line x+y+1=0 are in GP.

Watch Video Solution

197. A triangle has two sides $y = m_1 x$ and $y = m_2 x$ where m_1 and m_2 are the roots of the equation $b\alpha^2 + 2h\alpha + a = 0$. If (a, b) be the orthocenter of the triangle, then find the equation of the third side in terms of a, b and h.



198. The ratio in which the line 3x+4y+2=0 divides the distance between

3x+4y+5=0 and 3x+4y-5=0 is?

199. Let $A \equiv (6,7), B \equiv (2,3) and C \equiv (-2,1)$ be the vertices of a triangle. Find the point P in the interior of the triangle such that PBC is an equilateral triangle.

Watch Video Solution

200. Find the equations of lines parallel to 3x - 4y - 5 = 0 at a unit

distance from it.

Watch Video Solution

201. Let P (sin theta, cos theta) (0 le theta le 2pi) be a point and let OAB be a triangle with vertices (0,0) , (sqrt(3/2),0) and (0,sqrt(3/2)) Find theta if P lies inside triangle OAB.

202. Find the equation of a straight line passing through the point (-5,4) and which cuts off an intercept fo $\sqrt{2}$ units between the lines x+y+1=0 and x+y-1=0



203. If $A(x_1, y_1)$, $B(x_2, y_2)$, $C(x_3, y_3)$ are the vertices of a triangle, then the equation $|xy1x_1y_11x_2y_31| + |xy1x_1y_11x_3y_31| = 0$ represents (a)the median through A (b)the altitude through A (c)the perpendicular bisector of BC (d)the line joining the centroid with a vertex

Watch Video Solution

204. Are the points (3, 4) and (2, -6) on the same or opposite sides of

the line 3x - 4y = 8?



205. Consider the equation $y - y_1 = m(x - x_1)$. If m and x_1 are fixed and different lines are drawn for different values of y_1 , then

A. (a) the lines will pass through a fixed point

B. (b) there will be a set of parallel lines

C. (c) all the lines intersect the line $x=x_1$

D. (d) all the lines will be parallel to the line $y=x_1$

Answer: null

Watch Video Solution

206. एक रेखा ऐसी है कि लाइनों 5x - y + 4 = 0 तथा 3x + 4y - 4 = 0के बीच इसका खंड बिंदु (1,5) पर विभाजित है (1,5)। इसके समीकरण प्राप्त करें।

207. If the straight line ax + cy = 2b, where a, b, c > 0, makes a triangle of area 2 sq. units with the coordinate axes, then (a)a, b, c are in GP (b)a, -b; c are in GP (c)a, 2b, c are in GP (d) a, -2b, c are in GP



208. ABCD is a square whose vertices are A(0, 0), B(2, 0), C(2, 2), and D(0, 2). The square is rotated in the XY - plane through an angle 30^0 in the anticlockwise sense about an axis passing though Aperpendicular to the XY - plane. Find the equation of the diagonal BD of this rotated square.

Watch Video Solution

209. The x-coordinates of the vertices of a square of unit area are the roots of the equation $x^2 - 3|x| + 2 = 0$. The y-coordinates of the vertices are the roots of the equation $y^2 - 3y + 2 = 0$. Then the possible vertices of the square is/are (a)(1, 1), (2, 1), (2, 2), (1, 2)


210. Consider a triangle with vertices A(1, 2), B(3, 1) and C(-3, 0). Find the equation of altitude through vertex A.

Watch Video Solution

211. If (x, y) is a variable point on the line y = 2x lying between the lines

$$2(x+1)+y=0 ext{ and } x+3(y-1)=0$$
 , then (a) $x\in\left(-rac{1}{2},rac{6}{7}
ight)$ (b) $x\in\left(-rac{1}{2},rac{3}{7}
ight)$ (c) $y\in\left(-1,rac{3}{7}
ight)$ (d) $y\in\left(-1,rac{6}{7}
ight)$

212. A rectangle has two opposite vertices at the points (1, 2) and (5,5). It these vertices lie on the line x = 3, find the other vertices of the rectangle.

213. If D, E, and F are three points on the sides BC, AC, and AB of a triangle ABC such that AD, BE, and CF are concurrent, then show that $BD \cdot CE \cdot AF = DC \cdot EA \cdot FB$.



214. Find the coordinates of the foot of the perpendicular drawn from the

point (1, -2) on the line y = 2x + 1.

215. Let the sides of a parallelogram be U = a, U = b, V = a' and V = b', where U = lx + my + n, V = l'x + m'y + n'. Show that the equation of the diagonal through the point of intersection of U = a, V = a' and U = b, V = b' is given by det $\begin{bmatrix} U & a & b \\ V & a' & b' \\ 1 & 1 & 1 \end{bmatrix} = 0.$

Watch Video Solution

216. Find the image of the point (-8, 12) with respect to line mirror 4x + 7y + 13 = 0.

Watch Video Solution

217. One side of a rectangle lies along the line 4x + 7y + 5 = 0. Two of its vertices are (-3, 1) and (1, 1). Find the equations of the other three sides.

218. In a triangle ABC, side AB has equation 2x + 3y = 29 and side AC has equation x + 2y = 16. If the midpoint of BC is (5, 6), then find the equation of BC



219. The foot of the perpendicular on the line $3x + y = \lambda$ drawn from the origin is C. If the line cuts the x and the y-axis at AandB, respectively, then BC:CA is 1:3 (b) 3:1 (c) 1:9 (d) 9:1



220. Two consecutive sides of a parallelogram are 4x + 5y = 0 and 7x + 2y = 0. If the equation of one diagonal is 11x + 7y = 9, find the equation of the other diagonal.

221. The real value of a for which the value of m satisfying the equation $(a^2-1)m^2-(2a-3)m+a=0$ given the slope of a line parallel to
the y-axis is(a) $\frac{3}{2}$ (b) 0 (c) 1 (d) ± 1

Watch Video Solution

222. If one of the sides of a square is 3x-4y-12 = 0 and the center is (0,0), then find the equations of the diagonals of the square.

Watch Video Solution

223. If the quadrilateral formed by the lines

$$ax + by + c = 0$$
, $a'x + b'y + c = 0$, $ax + by + c' = 0$, $a'x + b'y + c' =$
has perpendicular diagonals, then (a) $b^2 + c^2 = b'^2 + c'^2$ (b)
 $c^2 + a^2 = c'^2 + a'^2$ (c) $a^2 + b^2 = a'^2 + b'^2$ (d) none of these

224. The vertex P of an equilateral triangle $\triangle PQR$ is at (2, 3) and the equation of the opposite side QR is given by x + y = 2. Find the possible equations of the side PQ.

Watch Video Solution

225. The straight lines 7x - 2y + 10 = 0 and 7x + 2y - 10 = 0 form an isosceles triangle with the line y = 2. The area of this triangle is equal to $\frac{15}{7}squants$ (b) $\frac{10}{7}squants$ $\frac{18}{7}squants$ (d) none of these Watch Video Solution

226. Find the least value of $(x-2)^2 + (y-2)^2$ under the condition 3x + 4y - 2 = 0.

227. θ_1 and θ_2 are the inclination of lines $L_1 and L_2$ with the x-axis. If $L_1 and L_2$ pass through $P(x_1, y_1)$, then the equation of one of the angle bisector of these lines is

(a)
$$\frac{x - x_1}{\cos\left(\frac{\theta_1 - \theta_2}{2}\right)} = \frac{y - y_1}{\sin\left(\frac{\theta_1 - \theta_2}{2}\right)} \quad \text{(b)} \frac{x - x_1}{-\sin\left(\frac{\theta_1 - \theta_2}{2}\right)} = \frac{y - y_1}{\cos\left(\frac{\theta_1 - \theta_2}{2}\right)} \quad \text{(c)}$$
$$\frac{x - x_1}{\sin\left(\frac{\theta_1 - \theta_2}{2}\right)} = \frac{y - y_1}{\cos\left(\frac{\theta_1 - \theta_2}{2}\right)} \quad \text{(d)} \frac{x - x_1}{-\sin\left(\frac{\theta_1 + \theta_2}{2}\right)} = \frac{y - y_1}{\cos\left(\frac{\theta_1 + \theta_2}{2}\right)}$$

Watch Video Solution

228. Find the least and greatest values of the distance of the point $(\cos heta, \sin heta), heta \in R$, from the line 3x - 4y + 10 = 0.

Watch Video Solution

229. A light ray coming along the line 3x + 4y = 5 gets reflected from the line ax + by = 1 and goes along the line 5x - 12y = 10. Then, (A) $a = \frac{64}{115}, b = \frac{112}{15}$ (B) $a = \frac{14}{15}, b = -\frac{8}{115}$ (C) $a = \frac{64}{115}, b = -\frac{8}{115}$ (D) $a = \frac{64}{15}, b = \frac{14}{15}$ **230.** Prove that the product of the lengths of the perpendiculars drawn from the points $\left(\sqrt{a^2 - b^2}, 0\right)$ and $\left(-\sqrt{a^2 - b^2}, 0\right)$ to the line $\frac{x}{a}$ $\cos \theta + \frac{y}{b} \sin \theta = 1$ is b^2 .

Watch Video Solution

231. Line ax + by + p = 0 makes angle $\frac{\pi}{4}$ with $x \cos \alpha + y \sin \alpha = p, p \in R^+$. If these lines and the line $x \sin \alpha - y \cos \alpha = 0$ are concurrent, then $a^2 + b^2 = 1$ (b) $a^2 + b^2 = 2$ $2(a^2 + b^2) = 1$ (d) none of these

Watch Video Solution

232. Two sides of a square lie on the lines x + y = 1 and x + y + 2 = 0.

What is its area?



233. A line is drawn perpendicular to line y = 5x, meeting the coordinate axes at AandB. If the area of triangle OAB is 10 sq. units, where O is the origin, then the equation of drawn line is 3x - y - 9 (b) $x - 5y = 10 \ x + 4y = 10$ (d) x - 4y = 10

> Watch Video Solution

234. Find the coordinates of a point on x + y + 3 = 0, whose distance from x + 2y + 2 = 0 is $\sqrt{5}$.

Watch Video Solution

235. If x - 2y + 4 = 0 and 2x + y - 5 = 0 are the sides of an isosceles triangle having area 10squares, the equation of the third side is (a) 3x - y = -9 (b) 3x - y + 11 = 0 (c) x - 3y = 19 (d) 3x - y + 15 = 0 **236.** If p is the length of perpendicular from the origin to the line whose intercepts on the axes are a and b, then show that $\frac{1}{p^2} = \frac{1}{a^2} + \frac{1}{b^2}$.

237. The number of values of a for which the lines 2x + y - 1 = 0, ax + 3y - 3 = 0, and 3x + 2y - 2 = 0 are concurrent is (a).0 (b) 1 (c) 2 (d) infinite

Watch Video Solution

238. The center of a square is at the origin and its one vertex is A(2, 1).

Find the coordinates of the other vertices of the square.

239. ABCD is a square $A \equiv (1, 2), B \equiv (3, -4)$. If line CD passes through (3, 8), then the midpoint of CD is (a) (2, 6) (b) (6, 2) (c) (2, 5) (d) $\left(\frac{28}{5}, \frac{1}{5}\right)$

Watch Video Solution

240. Find the distance between A(2, 3) on the line of gradient 3/4 and the point of intersection P of this line with 5x + 7y + 40 = 0.

Watch Video Solution

241. The equation of the straight line which passes through the point (-4, 3) such that the portion of the line between the axes is divided internally be the point in the ratio 5:3 is (A)9x - 20y + 96 = 0 (B) 9x + 20y = 24 (C) 20x + 9y + 53 = 0 (D) None of these



242. If one side of the square is 2x - y + 6 = 0, then one of the vertices

is (2, 1) . Find the other sides of the square.



243. The equation of the bisector of the acute angle between the lines

2x-y+4=0 and x-2y=1 is (a) x-y+5=0 (b)x-y+1=0

(c)x-y=5 (d) none of these

Watch Video Solution

244. Find equation of the line which is equidistant from parallel lines

9x + 6y - 7 = 0 and 3x + 2y + 6 = 0.

Watch Video Solution

245. If the equations y=mx+c and $x\coslpha+y\sinlpha=p$ represent the same straight line, then (a) $p=c\sqrt{1+m^2}$ (b) $c=p\sqrt{1+m^2}$ (c)

$$cp=\sqrt{1+m^2}$$
 (d) $p^2+c^2+m^2=1$

Watch Video Solution

246. Find the equation of the line passing through (2, 3) which is parallel to the x-axis.

Watch Video Solution

247. Consider three lines as follows. $L_1: 5x - y + 4 = 0$ $L_2: 3x - y + 5 = 0$ $L_3: x + y + 8 = 0$ If these lines enclose a triangle *ABC* and the sum of the squares of the tangent to the interior angles can be expressed in the form $\frac{p}{q}$, where *pandq* are relatively prime numbers, then the value of p + q is (a)500 (b)450 (c)230 (d) 465

248. Find the equation of a straight line cutting off an intercept-1 from the y-axis and being equally inclined to the axes.

Watch Video Solution

249. The line $L_1 \equiv 4x + 3y - 12 = 0$ intersects the x-and y-axies at Aand B, respectively. A variable line perpendicular to L_1 intersects the xand the y-axis at P and Q, respectively. Then the locus of the circumcenter of triangle ABQ is (a) 3x - 4y + 2 = 0 (b) 4x + 3y + 7 = 0 (c) 6x - 8y + 7 = 0 (d) none of these

Watch Video Solution

250. Find the equation of the line which intersects the y-axis at a distance of 2 units above the origin and makes an angle of 30^0 with the positive direction of the x-axis.

251. Find the locus of the point at which two given portions of the straight line subtend equal angle.



252. Find the equation of the perpendicular bisector of the line segment

joining the points A(2,3) and B(6, -5).

Watch Video Solution

253. Having given the bases and the sum of the areas of a number of triangles which have a common vertex, show that the locus of the vertex is a straight line.



254. Find the equation of a line that has y-intercept 4 and is a perpendicular to the line joining (2, -3) and (4, 2).



255. Find the equations of the diagonals of the square formed by the

lines x = 0, y = 0, x = 1 and y = 1.

Watch Video Solution

256. Find the equation of the straight line that passes through the point

(3,4) and is perpendicular to the line 3x + 2y + 5 = 0



257. Find the equation of the line which is parallel to 3x - 2y + 5 = 0and passes through the point (5, -6).



258. Consider two lines L_1andL_2 given by $a_1x + b_1y + c_1 = 0anda_2x + b_2y + c_2 = 0$ respectivelywherec1 and $c2 \neq$ intersecting at point $P\dot{A}$ line L_3 is drawn through the origin meeting the lines L_1andL_2 at AandB, respectively, such that PA = PB. Similarly, one more line L_4 is drawn through the origin meeting the lines L_1andL_2 at AandB, respectively. Such that PA = PB. Similarly, at A_1andB_2 , respectively, such that $PA_1 = PB_1$. Obtain the combined equation of lines L_3andL_4 .



259. Find the locus of a point P which moves such that its distance from the line $y = \sqrt{3}x - 7$ is the same as its distance from $(2\sqrt{3}, -1)$

260. Consider two lines L_1andL_2 given by x - y = 0 and x + y = 0, respectively, and a moving point P(x, y). Let $d(P, L_1), i = 1, 2$, represents the distance of point P from the line L_i . If point P moves in a certain region R in such a way that $2 \leq d(P, L_1) + d(P, L_2) \leq 4$, find the area of region R.

Watch Video Solution

Watch Video Solution

261. In what ratio does the line joining the points (2, 3) and (4, 1) divide the segment joining the points (1, 2) and (4, 3)?

262. Show that the lines 4x + y - 9 = 0, x - 2y + 3 = 0, 5x - y - 6 = 0 make equal intercepts on any line of slope 2. **Watch Video Solution**

263. Find the equation of the bisector of the obtuse angle between the

lines 3x - 4y + 7 = 0 and 12x + 5y - 2 = 0.



264. Show that if any line through the variable point A(k + 1, 2k) meets the lines 7x + y - 16 = 0, 5x - y - 8 = 0, x - 5y + 8 = 0 at B, C, D, respectively, the AC, AB, andAD are in harmonic progression. (The three lines lie on the same side of point A).

Watch Video Solution

265. The incident ray is along the line 3x - 4y - 3 = 0 and the reflected

ray is along the line 24x + 7y + 5 = 0. Find the equation of mirrors.

266. If the line $y = \sqrt{3}x$ cuts the curve $x^3 + y^3 + 3xy + 5x^2 + 3y^2 + 4x + 5y - 1 = 0$ at the point A, B, C, then $OA\dot{O}B\dot{O}C$ is equal to $\left(\frac{k}{13}\right)(3\sqrt{3}-1)$. The value of k is_____

Watch Video Solution

267. about to only mathematics

Watch Video Solution

268. The area of a parallelogram formed by the lines $ax \pm by \pm c = 0$ is

(a)
$$rac{c^2}{(ab)}$$
 (b) $rac{2c^2}{(ab)}$ (c) $rac{c^2}{2ab}$ (d) none of these

Watch Video Solution

269. The vertices BandC of a triangle ABC lie on the lines 3y = 4xandy = 0, respectively, and the side BC passes through the

point $\left(\frac{2}{3}, \frac{2}{3}\right)$. If *ABOC* is a rhombus lying in the first quadrant, *O*

being the origin, find the equation of the line BC_{\cdot}

Watch Video Solution

270. If each of the points $(x_1, 4)$, $(-2, y_1)$ lies on the line joining the points (2, -1)and(5, -3), then the point $P(x_1, y_1)$ lies on the line. (a) 6(x + y) - 25 = 0 (b) 2x + 6y + 1 = 0 (c)2x + 3y - 6 = 0 (d) 6(x + y) + 25 = 0

Watch Video Solution

271. If the lines

concurrent, show that the point $(a_1, b_1), (a_1, b_2)$ and (a_3, b_3) are collinear.

272. The diagonals of a parallelogram PQRS are along the lines x+3y =4

and 6x-2y = 7, Then PQRS must be :



274. A straight line segment of length/moves with its ends on two mutually perpendicular lines. Find the locus of the point which divides the line segment in the ratio 1:2



275. Find the foot of the perpendicular from the point (2, 4) upon x + y = 1.

276. The lines x + y - 1 = 0, $(m - 1)x + (m^2 - 7)y - 5 = 0$, and (m - 2)x + (2m - 5)y = 0 are a.) concurrent for three values of m b.) concurrent for no value of m c.) parallel for one value of m d.) parallel for two value of m

Watch Video Solution

277. In riangle ABC, vertex A is (1, 2). If the internal angle bisector of $\angle B$ is 2x - y + 10 = 0 and the perpendicular bisector of AC is y = x, then find the equation of BC



278. The equation of the line which bisects the obtuse angle between the

line x - 2y + 4 = 0 and 4x - 3y + 2 = 0 is



279. If the line ax + by = 1 passes through the point of intersection of

 $y = x an lpha + p \sec lpha, y \sin(30^\circ - lpha) - x \cos(30^\circ - lpha) = p$, and is

inclined at 30° with y= anlpha x , then prove that $a^2+b^2=rac{3}{4p^2}$.

Watch Video Solution

280. A straight line L is perpendicular to the line 5x - y = 1. The area of the triangle formed by line L, and the coordinate axes is 5. Find the equation of line L.

Watch Video Solution

281. The reflection of the point (4,-13) about the line 5x + y + 6 = 0 is (-1, -14) b. (3, 4) c. (0, -0) d. (1, 2)

282. Triangle ABC with AB = 13, BC = 5, and AC = 12 slides on the coordinates axes with AandB on the positive x-axis and positive y-axis respectively. The locus of vertex C is a line 12x - ky = 0. Then the value of k is_____

Watch Video Solution

283. The line
$$y = \frac{3x}{4}$$
 meets the lines $x - y + 1 = 0$ and $2x - y = 5$ at A and B respectively. Coordinates of P on $y = \frac{3x}{4}$ such that $PA \cdot PB = 25$.

Watch Video Solution

284. In a plane there are two families of lines y = x + r, y = -x + r, where $r \in \{0, 1, 2, 3, 4\}$. The number of squares of diagonals of length 2 formed by the lines is:

285. Line $\frac{x}{a} + \frac{y}{b} = 1$ cuts the co-ordinate axes at A(a,0) and B(0,b) and the line $\frac{x}{a}' + \frac{y}{b}' = -1$ at A'(-a', 0) and B'(0, -b'). If the points

A,B,A',B' are concyclic then the orthocentre of triangle ABA' is



287. The points (1, 3) and (5, 1) are two opposite vertices of a rectangle.

The other two vertices lie on the line the y=2x+c. Find c

A.-4

 $\mathsf{B.8}$

C. 24

Answer: A



288. The ends A and B of a straight line segment of constant length c slide upon the fixed rectangular axes OX and OY, respectively. If the rectangle OAPB be completed, then the locus of the foot of the perpendicular drawn from P to AB is

Watch Video Solution

289. All points lying inside the triangle formed by the points (1. 3). (5, 0) and (-1, 2) satisfy (A) $3x+2y\geq 0$ (B) $2x+y-13\geq 0$ (C) $2x-3y-12\leq 0$ (D) $-2x+y\geq 0$

290. The equation of the straight line passing through the point $(a\cos^3\theta, a\sin^3\theta)$ and perpendicular to the line $x \sec \theta + y \cos ec\theta = a$ is (A) $x\cos\theta - y\sin\theta = a\cos 2\theta$ (B) $x\cos\theta + y\sin\theta = a\cos 2\theta$ (C) $x\sin\theta + y\cos\theta = a\cos 2\theta$ (D) none of these

Watch Video Solution

291. The equation of a straight line on which the length of perpendicular from the origin is four units and the line makes an angle of 120^0 with the x-axis is (A) $x\sqrt{3} + y + 8 = 0$ (B) $x\sqrt{3} - y = 8$ (C) $x\sqrt{3} - y = 8$ (D) $x - \sqrt{3}y + 8 = 0$

Watch Video Solution

292. The number of integral values of m for which the x-coordinate of the point of intersection of the lines 3x + 4y = 9 and y = mx + 1 is also an integer is (a) 2 (b) 0 (c) 4 (d) 1

293. If the equation of base of an equilateral triangle is 2x - y = 1 and the vertex is (-1, 2), then the length of the sides of the triangle is $\sqrt{\frac{20}{3}}$ (b) $\frac{2}{\sqrt{15}} \sqrt{\frac{8}{15}}$ (d) $\sqrt{\frac{15}{2}}$

Watch Video Solution

294. The equation of straight line passing through (-a, 0) and making a triangle with the axes of area T is (a) $2Tx + a^2y + 2aT = 0$ (b) $2Tx - a^2y + 2aT = 0$ (c) $2Tx - a^2y - 2aT = 0$ (d)none of these

Watch Video Solution

295. The line PQ whose equation is x - y = 2 cuts the x-axis at P ,and Q is (4,2). The line PQ is rotated about P through 45° in the anticlockwise direction. The equation of the line PQ in the new position is (A) $y = -\sqrt{2}$ (B) y = 2 (C) x = 2 (D) x = -2 **296.** If the equation of the locus of a point equidistant from the points (a_1, b_1) and (a_2, b_2) is $(a_1 - a_2)x + (b_1 - b_2)y + c = 0$, then find the value of c.

Watch Video Solution

297. The extremities of the base of an isosceles triangle are (2, 0)and(0, 2). If the equation of one of the equal side is x = 2, then the equation of other equal side is (a) x + y = 2 (b) x - y + 2 = 0 (c) y = 2 (d) 2x + y = 2

Watch Video Solution

298. A triangle is formed by the lines x + y = 0, x - y = 0, and lx + my = 1. If l and m vary subject to the condition $l^2 + m^2 = 1$, then the locus of its circumcenter is (a) $(x^2 - y^2)^2 = x^2 + y^2$ (b)

$$ig(x^2+y^2ig)^2 = ig(x^2-y^2ig)$$
 (c) $ig(x^2+y^2ig)^2 = 4x^2y^2$ (d) $ig(x^2-y^2ig)^2 = ig(x^2+y^2ig)^2$

Watch Video Solution

299. The line x + y = p meets the x- and y-axes at AandB, respectively. A triangle APQ is inscribed in triangle OAB, O being the origin, with right angle at $Q\dot{P}$ and Q lie, respectively, on OBandAB. If the area of triangle APQ is $\frac{3}{8}th$ of the are of triangle OAB, the $\frac{AQ}{BQ}$ is equal to (a)2(b) $\frac{2}{3}$ (c) $\frac{1}{3}$ (d)3

Watch Video Solution

300. A is a point on either of two lines $y + \sqrt{3}|x| = 2$ at a distance of $4\sqrt{3}$ units from their point of intersection. The coordinates of the foot of perpendicular from A on the bisector of the angle between them are (a)

$$\left(-rac{2}{\sqrt{3}},2
ight)$$
 (b) $(0,0)$ (c) $\left(rac{2}{\sqrt{3}},2
ight)$ (d) $(0,4)$

301. A pair of perpendicular straight lines is drawn through the origin forming with the line 2x + 3y = 6 an isosceles triangle right-angled at the origin. The equation to the line pair is (a) $5x^2 - 24xy - 5y^2 = 0$ (b) $5x^2 - 26xy - 5y^2 = 0$ (c) $5x^2 + 24xy - 5y^2 = 0$ (d) $5x^2 + 26xy - 5y^2 = 0$

Watch Video Solution

302. If the vertices PandQ of a triangle PQR are given by (2, 5) and (4, -11), respectively, and the point R moves along the line N given by 9x + 7y + 4 = 0, then the locus of the centroid of triangle PQR is a straight line parallel to PQ (b) QR (c) RP (d) N

303. Given $A \equiv (1, 1)$ and AB is any line through it cutting the x-axis at B. If AC is perpendicular to AB and meets the y-axis in C, then the equation of the locus of midpoint P of BC is (a) x + y = 1 (b) x + y = 2 (c) x + y = 2xy (d) 2x + 2y = 1

Watch Video Solution

304. The straight lines 4ax + 3by + c = 0, where a + b + c are concurrent at the point a) (4,3) b) $\left(\frac{1}{4},\frac{1}{3}\right)$ c) $\left(\frac{1}{2},\frac{1}{3}\right)$ d) none of these

Watch Video Solution

305. The line parallel to the x-axis and passing through the intersection of the lines ax + 2by + 3b = 0 and bx - 2y - 3a = 0, where $(a, b) \neq (0, 0)$, is (a)above the x-axis at a distance of 3/2 units from it (b)above the x-axis at a distance of 2/3 units from it (c)below the x-axis at a distance of 3/2 units from it (d)below the x-axis at a distance of 2/3 units from it

Watch Video Solution

306. The line $L_1: y - x = 0$ and $L_2: 2x + y = 0$ intersect the line $L_3: y + 2 = 0$ at P and Q respectively. The bisector of the acute angle between L_1 and L_2 intersects L_3 at R. Statement-1 : The ratio PR: RQ equals $2\sqrt{2}: \sqrt{5}$ Statement-2 : In any triangle, bisector of an angle divides the triangle into two similar triangles. Statement-1 is true, Statement-2 is true ; Statement-2 is correct explanation for Statement-1 Statement-1 is true, Statement-2 is true ; Statement-1 is true, Statement-2 is true ; Statement-2 is false Statement-1 is false, Statement-2 is true

Watch Video Solution

307. If the lines ax + y + 1 = 0, x + by + 1 = 0, and x + y + c = 0(a, b, c being distinct and different from 1) are concurrent,



308. Two sides of a rhombus ABCD are parallel to the lines y = x + 2 and y =

7x + 3 If the diagonals of the rhombus intersect at the point (1, 2) and the

vertex A is on the y-axis, then vertex A can be

Watch Video Solution

309. Equation(s) of the straight line(s), inclined at 30° to the x-axis such that the length of its (each of their) line segment(s) between the coordinates axes is 10 units, is (are)

310. If a pair of perpendicular straight lines drawn through the origin forms an isosceles triangle with the line 2x + 3y = 6, then area of the triangle so formed is (a) $\frac{36}{13}$ (b) $\frac{12}{17}$ (c) $\frac{13}{5}$ (d) $\frac{17}{14}$



311. The sides of a rhombus are parallel to the lines x + y - 1 = 0 and 7x - y - 5 = 0. It is given that the diagonals of the rhombus intersect at (1, 3) and one vertex, A of the rhombus lies on the line y = 2x. Then the coordinates of vertex A are $(a)\left(\frac{8}{5}, \frac{16}{5}\right)$ (b) $\left(\frac{7}{15}, \frac{14}{15}\right)$ (c) $\left(\frac{6}{5}, \frac{12}{5}\right)$ (d) $\left(\frac{4}{15}, \frac{8}{15}\right)$

Watch Video Solution

312. The image of P(a, b) on the line y = -x is Q and the image of Q on the line y = x is R. Then the midpoint of PR is (a) (a + b, b + a) (b) $\left(\frac{a+b}{2}, \frac{b+2}{2}\right)$ (c) (a - b, b - a) (d) (0, 0)
313. Consider a $\triangle ABC$ whose sides AB, BC and CA are represented by the straight lines 2x + y = 0, x + py = q and x - y = 3respectively. The point P is (2, 3). If P is orthocentre, then find the value of (p+q) is

Watch Video Solution

314. The equations of two equal sides ABandAC of an isosceles triangle ABC are x + y = 5 and 7x - y = 3, respectively. Then the equation of side BC if $ar(ABC) = 5unit^2$ is x - 3y + 1 = 0 (b) x - 3y - 21 = 03x + y + 2 = 0 (d) 3x + y - 12 = 0

Watch Video Solution

315. Find the area of the triangle formed by the line x + y = 3 and the angle bisectors of the pair of lines $x^2 - y^2 + 2y = 1$

316. The sides of a triangle have the combined equation $x^2 - 3y^2 - 2xy + 8y - 4 = 0$. The third side, which is variable, always passes through the point (-5, -1). Find the range of values of the slope of the third line such that the origin is an interior point of the triangle.

Watch Video Solution

317. The equation of the lines passing through the point (1, 0) and at a distance $\frac{\sqrt{3}}{2}$ from the origin is (a). $\sqrt{3} + y - \sqrt{3} = 0$ (b). $x + \sqrt{3}y - \sqrt{3} = 0$ (c). $\sqrt{3}x - y - \sqrt{3} = 0$ (d). $x - \sqrt{3}y - \sqrt{3} = 0$ **Watch Video Solution**

318. The number of values of k for which the lines (k+1)x + 8y = 4k and kx + (k+3)y = 3k - 1 are coincident is



(2a+b)x+(a+3b)y+(b-3a)=0 and mx+2y+6=0 are

concurrent. Then |m| is equal to_____

Watch Video Solution

320. The line x = c cuts the triangle with corners (0, 0), (1, 1) and (9, 1)into two region. two regions to be the same c must be equal to (A) $\frac{5}{2}$ (B) 3 (C) $\frac{7}{2}$ (D) 5 or 15

Watch Video Solution

321. The absolute value of the sum of the abscissas of all the points on the line x + y = 4 that lie at a unit distance from the line



322. The point (x,y) lies on the line 2x+3y=6. The smallest value of the quantity $\sqrt{x^2+y^2}$ is m then the value of $\sqrt{13} \ m$ is_____

Watch Video Solution

323. The equations of the perpendicular bisectors of the sides ABandAC of triangle ABC are x - y + 5 = 0 and x + 2y = 0, respectively. If the point A is (1, -2), then find the equation of the line BC.

Watch Video Solution

324. One of the diagonals of a square is the portion of the line $\frac{x}{2} + \frac{y}{3} = 2$ intercepted between the axes. Then the extremities of the

other diagonal are: (a) (5,5), (-1,1) (b) (0,0), (4,6), (0,0), (-1,1)(d) (5,5), 4, 6)

Watch Video Solution

325. Two sides of a triangle are along the coordinate axes and the medians through the vertices (other than the origin) are mutually perpendicular. The number of such triangles is/are (a) zero (b) two (c) four (d) infinite

Watch Video Solution

326. The graph of $y^2 + 2xy + 40|x| = 400$ divides the plane into regions. Then the area of the bounded region is (a)200squnits (b) 400squnits (c)

800squnits (d) 500squnits

327. In a triangle ABC, $A = (\alpha, \beta)B = (2, 3)$, andC = (1, 3). Point A lies on line y = 2x + 3, where $\alpha \in I$. The area of ABC, , is such that $[\Delta] = 5$. The possible coordinates of A are (where [.] represents greatest integer function). (a)(2, 3) (b) (5, 13) (-5, -7) (d) (-3, -5)

Watch Video Solution

328. If the straight lines 2x + 3y - 1 = 0, x + 2y - 1 = 0, and ax + by - 1 = 0 form a triangle with the origin as orthocentre, then (a, b) is given by (a) (6, 4) (b) (-3, 3) (c) (-8, 8) (d) (0, 7)

Watch Video Solution

329. Let *O* be the origin. If A(1, 0)andB(0, 1)andP(x, y) are points such that xy > 0andx + y < 1, then (a)*P* lies either inside the triangle *OAB* or in the third quadrant. (b)*P* cannot lie inside the triangle *OAB* (c)*P* lies inside the triangle *OAB* (d)*P* lies in the first quadrant only

330. If the area of the rhombus enclosed by the lines $lx\pm my\pm n=0$ is

2 sq. units, then, a) l,m,n are in G.P b) l,n,m are in G.P. c) lm=n d) ln=m

Watch Video Solution

331. In a triangle ABC, the bisectors of angles BandC lies along the lines x = yandy = 0. If A is (1, 2), then the equation of line BC is (a)

2x+y=1 (b) 3x-y=5 (c) x-2y=3 (d) x+3y=1

332. If
$$\frac{a}{bc} - 2 = \sqrt{\frac{b}{c}} + \sqrt{\frac{c}{b}}$$
, where $a, b, c > 0$, then the family of lines $\sqrt{ax} + \sqrt{by} + \sqrt{c} = 0$ passes though the fixed point given by (a) (1, 1)
(b) $(1, -2) (-1, 2)$ (d) $(-1, 1)$

333. P(m, n) (where m, n are natural numbers) is any point in the interior of the quadrilateral formed by the pair of lines xy = 0 and the lines 2x + y - 2 = 0 and 4x + 5y = 20. The possible number of positions of the point P is. 7 (b) 5 (c) 4 (d) 6

Watch Video Solution

334. A diagonal of rhombus ABCD is member of both the families of lines $(x + y - 1) + \lambda(2x + 3y - 2) = 0$ and $(x - y + 2) + \lambda(2x - 3y + 5) = 0$ and rhombus is (3, 2). If the area of the rhombus is $12\sqrt{5}$ sq. units, then find the remaining vertices of the rhombus.

335. A regular polygon has two of its consecutive diagonals as lines $\sqrt{3}x + y = \sqrt{3}$ and $2y = \sqrt{3}$. Point (1,c) is one of its vertices. Find the

equation of the sides of the polygon and also find the coordinates of the vertices.



336. Find the locus of the circumcenter of a triangle whose two sides are along the coordinate axes and the third side passes through the point of intersection of the line ax + by + c = 0 and lx + my + n = 0.

Watch Video Solution

337. A line $L_1 \equiv 3y - 2x - 6 = 0$ is rotated about its point of intersection with the y-axis in the clockwise direction to make it L_2 such that the are formed by L_1 , L_2 the x-axis, and line x = 5 is $\frac{49}{3}$ squarits if its point of intersection with x = 5 lies below the x-axis. Find the equation of L_2 .

338. Straight lines $y = mx + c_1$ and $y = mx + c_2$ where $m \in R^+$, meet the x-axis at A_1andA_2 , respectively, and the y-axis at B_1andB_2 , respectively. It is given that points A_1, A_2, B_1 , and B_2 are concylic. Find the locus of the intersection of lines A_1B_2 and A_2B_1 .

Watch Video Solution

339. Show that the reflection of the line ax + by + c = 0 on the line

x+y+1=0 is the line b+ay+(a+b-c)=0 where a
eq b

Watch Video Solution

340. Two equal sides of an isosceles triangle are 7x-y+3=0 and x+y-3=0. Its

third side passes the point (1,-10).

Determine the equation of the third side.

341. The number of possible straight lines passing through point(2,3) and forming a triangle with coordiante axes whose area is 12 sq. unit is: a. one b. two c. three d. four

Watch Video Solution

342. In a triangle ABC, if A is (2, -1), and7x - 10y + 1 = 0 and 3x - 2y + 5 = 0 are the equations of an altitude and an angle bisector, respectively, drawn from B, then the equation of BC is (a) a + y + 1 = 0 (b)5x + y + 17 = 0 (c)4x + 9y + 30 = 0 (d) x - 5y - 7 = 0

Watch Video Solution

343. The sides of a triangle are the straight line x+y=1, 7y=x, and $\sqrt{3}y + x = 0$. Then which of the following is an interior point of the triangle?

344. One of the diameter of a circle circumscribing the rectangle ABCD is

4y = x + 7, If A and B are the points (-3,4) and (5,4) respectively,

then the area of rectangle is

Watch Video Solution

345. The coordinates of two consecutive vertices A and B of a regular hexagon ABCDEF are (1, 0) and (2, 0), respectively. The equation of the diagonal CE is (a) $\sqrt{3}x + y = 4$ (b) $x + \sqrt{3}y + 4 = 0$ (c) $x + \sqrt{3}y = 4$ (d) none of these

Watch Video Solution

346. P is a point on the line y + 2x = 1, and Q and R are two points on the line 3y + 6x = 6 such that triangle PQR is an equilateral triangle.



348. In $\triangle ABC$, the coordinates of the vertex A are, (4, -1) and lines x - y - 1 = 0 and 2x - y = 3 are the internal bisectors of angles B and C. Then the radius of the circles of triangle AbC is



349. If the equation of any two diagonals of a regular pentagon belongs to the family of lines $(1+2\lambda)y - (2+\lambda)x + 1 - \lambda = 0$ and their

lengths are sin 36^0 , then the locus of the center of circle circumscribing the given pentagon (the triangles formed by these diagonals with the sides of pentagon side have common) no is (a) $x^2 + y^2 - 2x - 2y + 1 + \sin^2 72^0 = 0$ (b) $x^2 + y^2 - 2x - 2y + \cos^2 72^0 = 0$ (c) $x^2 + y^2 - 2x - 2y + 1 + \cos^2 72^0 = 0$ (d) $x^2 + y^2 - 2x - 2y + \sin^2 72^0 = 0$

Watch Video Solution

350. If it is possible to draw a line which belongs to all the given family of lines

$$y-2x+1+\lambda_1(2y-x-1)=0,$$
 $3y-x-6+\lambda_2(y-3x+6)=0,$
 $ax+y-2+\lambda_3(6x+ay-a)=0$, then
(a) $a=4$ (b) $a=3$ (c) $a=-2$ (d) $a=2$

351. The locus of the image of the point
$$(2,3)$$
 in the line
 $(x-2y+3)+\lambda(2x-3y+4)=0$ is $(\lambda \in R)$ (a)
 $x^2+y^2-3x-4y-4=0$ (b) $2x^2+3y^2+2x+4y-7=0$ (c)
 $x^2+y^2-2x-4y+4=0$ (d) none of these

Watch Video Solution

352. ABC is a variable triangle such that A is (1,2) and B and C lie on line $y = x + \lambda$ (where λ is a variable). Then the locus of the orthocentre of triangle ABC is (a) $(x - 1)^2 + y^2 = 4$ (b) x + y = 3 (c) 2x - y = 0 (d) none of these

Watch Video Solution

353. If $P\left(1+\frac{t}{\sqrt{2}},2+\frac{t}{\sqrt{2}}\right)$ is any point on a line, then the range of

the values of t for which the point P lies between the parallel lines



355. If the extremities of the base of an isosceles triangle are the points

(2a,0) and (0,a), and the equation of one of the sides x=2a, then the area

of the triangle is



356. The coordinates of the foot of the perpendicular from the point (2,3) on the line -y + 3x + 4 = 0 are given by (a) $\left(\frac{37}{10}, -\frac{1}{10}\right)$ (b)

$$(-\frac{1}{10},\frac{37}{10})$$
 (c) $\left(\frac{10}{37},\ -10\right)$ (d) $\left(\frac{2}{3},\ -\frac{1}{3}\right)$

Watch Video Solution

357. The straight lines x + 2y - 9 = 0, 3x + 5y - 5 = 0, and ax + by - 1 = 0 are concurrent, if the straight line 35x - 22y + 1 = 0 passes through the point (a).(a, b) (b) (b, a) (-a, -b) (d) none of these

Watch Video Solution

358. If lines x + 2y - 1 = 0, ax + y + 3 = 0, and bx - y + 2 = 0 are concurrent, and S is the curve denoting the locus of (a, b), then the least distance of S from the origin is (a) $\frac{5}{\sqrt{57}}$ (b) $\frac{5}{\sqrt{51}}$ (c) $\frac{5}{\sqrt{58}}$ (d) $\frac{5}{\sqrt{59}}$

359. $L_1 and L_2$ are two lines. If the reflection of $L_1 on L_2$ and the reflection of L_2 on L_1 coincide, then the angle between the lines is (a) 30^0 (b) 60^0 45^0 (d) 90^0

Watch Video Solution

360. $A \equiv (-4, 0), B \equiv (4, 0) \dot{M} and N$ are the variable points of the yaxis such that M lies below NandMN = 4. Lines AMandBN intersect at P. The locus of P is (a) $2xy - 16 - x^2 = 0$ (b) $2xy + 16 - x^2 = 0$ (c) $2xy + 16 + x^2 = 0$ (d) $2xy - 16 + x^2 = 0$

Watch Video Solution

361. If $\sin(\alpha + \beta)\sin(\alpha - \beta) = \sin\gamma(2\sin\beta + \sin\gamma)$, where $0 < \alpha, \beta, \gamma < \pi$, then the straight line whose equation is $x\sin\alpha + y\sin\beta - \sin\gamma = 0$ passes through point (a) (1, 1) (b) (-1, 1) (c) (1, -1) (d) none of these 362. Let P be (5,3) and a point R on y = x and Q on the X - axis be such that

PQ + QR + RP is minimum ,then the coordinates of Q are

Watch Video Solution

363. Given A(0,0) and B(x,y) with $x \in (0,1)$ and y>0. Let the slope of line AB be m_1 . Point C lies on line x = 1 such that the slope of BC is equal to m_2 where $0 < m_2 < m_1$. If the area of triangle ABC can be expressed as $(m_1 - m_2)f(x)$ then the largest possible value of x is

Watch Video Solution

364. If the straight lines x + y - 2 = 0, 2x - y + 1 = 0 and ax + by - c = 0 are concurrent, then the family of lines 2ax + 3by + c = 0(a, b, c are nonzero) is concurrent at (a) (2, 3) (b) $\left(\frac{1}{2}, \frac{1}{3}\right)$ (c) $\left(-\frac{1}{6}, -\frac{5}{9}\right)$ (d) $\left(\frac{2}{3}, -\frac{7}{5}\right)$

365. The equaiton of the lines through the point (2, 3) and making an intercept of length 2 units between the lines y + 2x = 3 and y + 2x = 5 are (A) x + 3 = 0, 3x + 4y = 12 (B) y - 2 = 0, 4x - 3y = 6 (C) x - 2 = 0, 3x + 4y = 18 (D) none of these Watch Video Solution

366. A beam of light is sent along the line x - y = 1, which after refracting from the x-axis enters the opposite side by turning through 30^0 towards the normal at the point of incidence on the x-axis. Then the equation of the refracted ray is (a) $(2 - \sqrt{3})x - y = 2 + \sqrt{3}$ (b) $(2 + \sqrt{3})x - y = 2 + \sqrt{3}$ (c) $(2 - \sqrt{3})x + y = (2 + \sqrt{3})$ (d) $y = (2 - \sqrt{3})(x - 1)$

367. Determine all the values of α for which the point (α, α^2) lies inside the triangle formed by the lines. 2x + 3y - 1 = 0 x + 2y - 3 = 05x - 6y - 1 = 0

368. A line through A(-5, -4) meets the lines x + 3y + 2 = 0, 2x + y + 4 = 0 and x - y - 5 = 0 at the points B, CandD respectively, if $\left(\frac{15}{AB}\right)^2 + \left(\frac{10}{AC}\right)^2 = \left(\frac{6}{AD}\right)^2$ find the equation of the line.

Watch Video Solution

369. If $u = a_1x + b_1y + c_1 = 0$, $v = a_2x + b_2y + c_2 = 0$, and $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, then the curve u + kv = 0 is (a)the same straight line u (b)different straight line (c)not a straight line (d)none of these

370. The point A(2, 1) is translated parallel to the line x - y = 3 by a distance of 4 units. If the new position A' is in the third quadrant, then the coordinates of A' are (A) $(2 + 2\sqrt{2}, 1 + 2\sqrt{2})$ (B) $(-2 + \sqrt{2}, -1 - 2\sqrt{2})$ (C) $(2 - 2\sqrt{2}, 1 - 2\sqrt{2})$ (D) none of these

Watch Video Solution

371. Let ABC be a triangle. Let A be the point (1, 2), y = x be the perpendicular bisector of AB, and x - 2y + 1 = 0 be the angle bisector of $\angle C$. If the equation of BC is given by ax + by - 5 = 0, then the value of a + b is (a)1(b) 2(c) 3 (d) 4

Watch Video Solution

372. The area enclosed by $2|x|+3|y|\leq 6$ is (a)3 sq. units (b) 4 sq. units

(c)12 sq. units (d) 24 sq. units

373. The lines $y=m_1x, y=m_2xandy=m_3x$ make equal intercepts on

the line
$$x + y = 1$$
. Then (a)
 $2(1 + m_1)(1 + m_3) = (1 + m_2)(2 + m_1 + m_3)$ (b)
 $(1 + m_1)(1 + m_3) = (1 + m_2)(1 + m_1 + m_3)$ (c)
 $(1 + m_1)(1 + m_2) = (1 + m_3)(2 + m_1 + m_3)$ (d)
 $2(1 + m_1)(1 + m_3) = (1 + m_2)(1 + m_1 + m_3)$

Watch Video Solution

374. The condition on a and b, such that the portion of the line ax + by - 1 = 0 intercepted between the lines ax + y = 0 and x + by = 0 subtends a right angle at the origin, is a = b (b) a + b = 0a = 2b (d) 2a = b

375. One diagonal of a square is along the line 8x - 15y = 0 and one of its vertex is (1, 2). Then the equations of the sides of the square passing through this vertex are (a)23x + 7y = 9, 7x + 23y = 53(b)23x - 7y + 9 = 0, 7x + 23y + 53 = 0(c)23x - 7y - 9 = 0, 7x + 23y - 53 = 0 (d)none of these

Watch Video Solution

376. The straight line ax + by + c = 0, where $abc \neq 0$, will pass through the first quadrant if (a) ac > 0, bc > 0 (b) ac > 0 or bc < 0 (c) bc > 0 or ac > 0 (d) ac < 0 or bc < 0

Watch Video Solution

377. A square of side a lies above the x-axis and has one vertex at the origin. The side passing through the origin makes an angle $\alpha \left(0 < \alpha < \frac{\pi}{4}\right)$ with the positive direction of x-axis. equation its diagonal not passing through origin is (a)





378. If the sum of the distances of a point from two perpendicular lines in

a plane is 1, then its locus is

(a)a square (b) a circle (c) a straight line (d) two intersecting lines

Watch Video Solution

379. ABC is a variable triangle such that A is (1, 2), and B and C on the line $y = x + \lambda(\lambda)$ is a variable). Then the locus of the orthocentre of ΔABC is (a)x + y = 0 (b) x - y = 0 $x^2 + y^2 = 4$ (d) x + y = 3

380. Consider a ABC is which side ABandAC are perpendicular to x - y - 4 = 0and2x - y - 5 = 0, respectively. Vertex A is (-2, 3) and the circumcenter of ABC is $\left(\frac{3}{2}, \frac{5}{2}\right)$. The equation of line in column I is of the form ax + by + c = 0, where $a, b, c \in I$. Match it with the corresponding value of c in column II Column I|Column II Equation of the perpendicular bisector of side $BA \mid p. -1$ Equation of the perpendicular bisector of side $AC \mid q. 1$ Equation of side $AC \mid r. -16$ Equation of the median through $A \mid s. -4$

Watch Video Solution

381. Column I|Column II Two vertices of a triangle are (5, -1)and(-2, 3). If the orthocentre is the origin, then the coordinates of the third vertex are|p. (-4, -7) A point on the line x + y = 4 which lies at a unit distance from the line 4x + 3y = 10 is|q. (-7, 11) The orthocentre of the triangle formed by the lines x + y - 1 = 0, x - y + 3 = 0, 2x + y = 7 is|r. (2, -2) If 2a, b, c are in AP, then lines ax + by = c are concurrent at|s. (-1, 2)

lines

 $(a+b)x+(a-b)y-2ab=0,\,(a-b)x+(a+b)y-2ab=0\,\, ext{and}\,\,x+y$

form an isosceles triangle whose vertical angle is

Watch Video Solution

383. Each equation contains statements given in two columns which have to be matched. Statements (a,b,c,d) in column I have to be matched with Statements (p, q, r, s) in column II. If the correct match are $a\overrightarrow{p}, a\overrightarrow{s}, b\overrightarrow{q}, b\overrightarrow{r}, c\overrightarrow{p}, c\overrightarrow{q}$, and $d\overrightarrow{s}$, then the correctly bubbled 4x4 matrix should be as follows: Figure Consider the lines represented by equation $(x^2 + xy - x)x(x - y) = 0$, forming a triangle. Then match the following: Column II Orthocenter of triangle |p. $(\frac{1}{6}, \frac{1}{2})$ Circumcenter|q. $(1(2 + 2\sqrt{2}), \frac{1}{2})$ Centroid|r. $(0, \frac{1}{2})$ Incenter|s. $(\frac{1}{2}, \frac{1}{2})$

384. The st. lines 3x + 4y = 5 and 4x - 3y = 15 interrect at a point A(3, -1). On these linepoints B and C are chosen so that AB = AC. Find the possible eqns of the line BC pathrough the point (1, 2)

Watch Video Solution

385. The area of the triangular region in first quadrant bounded on the left by the line 7x + 4y = 168, and bounded below by the line 5x + 3y = 121 is A. Then the value of $\frac{3A}{10}$ is_____

> Watch Video Solution

386. Find the area enclosed by the graph of
$$x^2y^2 - 9x^2 - 25y^2 + 225 = 0.$$

387. Lines $L_1 \equiv ax + by + c = 0$ and $L_2 \equiv lx + my + n = 0$ intersect at the point P and make an angle θ with each other. Find the equation of a line different from L_2 which passes through P and makes the same angle θ with L_1 .

Watch Video Solution

388. Let ABC be a triangle with AB = AC. If D is the midpoint of BC, E is the foot of the perpendicular drawn from D to AC, andF is the midpoint of DE, then prove that AF is perpendicular to BE.

Watch Video Solution

389. For a > b > c > 0, the distance between (1, 1) and the point of

intersection of the lines ax+by+c =0 and bx+ay+c=0 is les than $2\sqrt{2}$. Then

390. A straight line L through the point (3,-2) is inclined at an angle 60° to the line $\sqrt{3}x + y = 1$. If L also intersects the x-axis then find the equation of L.

Watch Video Solution

391. The locus of the orthocentre of the triangle formed by the lines (1+p)x - py + p(1+p) = 0, (1+q)x - qy + q(1+q) = 0 and y = 0, where $p \neq \cdot q$, is (A) a hyperbola (B) a parabola (C) an ellipse (D) a straight line

Watch Video Solution

392. The vertices of a triangle are (A(-1, -7), B(5, 1), and C(1, 4)). The equation of the bisector of $\angle ABC$ is

393. Let the algebraic sum of the perpendicular distance from the points (2, 0), (0,2), and (1, 1) to a variable straight line be zero. Then the line passes through a fixed point whose coordinates are___

394. A straight line through the origin 'O' meets the parallel lines 4x + 2y = 9 and 2x + y = -6 at points P and Q respectively. Then the point 'O' divides the segment PQ in the ratio



395. A straight line I with negative slope passes through (8,2) and cuts the coordinate axes at P and Q. Find absolute minimum value of "OP+OQ where O is the origin-



396. A straight line L through the origin meets the lines x + y = 1 and x + y = 3 at P and Q respectively. Through P and Q two straight lines L_1 , and L_2 are drawn, parallel to 2x - y - 5 and 3x + y5 respectively. Lines L_1 and L_2 intersect at R. Locus of R, as L varies, is

Watch Video Solution

397. A rectangle PQRS has its side PQ parallel to the line y = mx and vertices P, Q, and S on the lines y = a, x = b, and x = -b, respectively. Find the locus of the vertex R.

> Watch Video Solution

398. The area of the triangle formed by the intersection of a line parallel to x-axis and passing through P (h, k) with the lines y = x and x + y = 2 is $4h^2$. Find the locus of the point P.

399. The lines ax + by + c = 0, where 3a + 2b + 4c = 0, are concurrent at the

point (a)
$$\left(rac{1}{2},rac{3}{4}
ight)$$
 (b) $(1,3)$ (c) $(3,1)$ (d) $\left(rac{3}{4},rac{1}{2}
ight)$

Watch Video Solution

400. The area enclosed within the curve |x|+|y|=1 is



401. Find the orthocentre of the triangle the equations of whose sides

are x + y = 1, 2x + 3y = 6 and 4x - y + 4 = 0.

Watch Video Solution

402. If a, b and c are in AP, then the straight line ax + by + c = 0 will always pass through a fixed point whose coordinates are (a) (1,2) (b) (1,-2) (c) (2,3) (d) (0,0)



403. Statement-I: If the diagonals of the quadrilateral formed by the lines px + gy + r = 0, p'x + gy + r' = 0, p'x + q'y + r' = 0, are at right angles, then $p^2 + q^2 = p^2 + q^2$. Statement-2: Diagonals of a rhombus are bisected and perpendicular to each other.



404. Statement :Two different lines can be drawn passing through two given points.

Watch Video Solution

405. Statement 1: The internal angle bisector of angle C of a triangle ABC with sides AB, AC, and BC as y = 0, 3x + 2y = 0, and 2x + 3y + 6 = 0, respectively, is 5x + 5y + 6 = 0 Statement 2: The

image of point A with respect to 5x+5y+6=0 lies on the side BC of the triangle.



406. The joint equation of lines y = xandy = -x is $y^2 = -x^2$, i.e., $x^2 + y^2 = 0$ Statement 2: The joint equation of lines ax + by = 0 and cx + dy = 0 is (ax + by)(cx + dy) = 0, wher a, b, c, d are constant.

Watch Video Solution

407. Statement I If sum of algebraic distances from points A(1,2),B(2,3),C(6,1) is zero on the line ax + by + c = 0 then 2a + 3b + c = 0,

Statement II The centroid of the triangle is (3,2)

408. Each question has four choice: a, b, c and d, out of which only one is correct. Each question contains Statement 1 and Statement 2. Find the correct answer. Both the Statements are true but Statement 2 is the correct explanation of Statement 1. Both the Statement are True but Statement 2 is not the correct explanation of Statement 1. Statement 1. Statement 1 is True and Statement 2 is False. Statement 1 is False and Statement 2 is True Statement 1: The lines (a + b)x + (a - 2b)y = a are concurrent at the point $(\frac{2}{3}, \frac{1}{3})$. Statement 2: The lines x + y - 1 = 0 and x - 2y = 0 intersect at the point $(\frac{2}{3}, \frac{1}{3})$.

Watch Video Solution

409. Statement 1:If the point $(2a - 5, a^2)$ is on the same side of the line x + y - 3 = 0 as that of the origin, then $a \in (2, 4)$ Statement 2: The points $(x_1, y_1)and(x_2, y_2)$ lie on the same or opposite sides of the line ax + by + c = 0, as $ax_1 + by_1 + c$ and $ax_2 + by_2 + c$ have the same or opposite signs.
410. Statement 1: Each point on the line y - x + 12 = 0 is equidistant from the lines 4y + 3x - 12 = 0, 3y + 4x - 24 = 0 Statement 2: The locus of a point which is equidistant from two given lines is the angular bisector of the two lines.

Watch Video Solution

411. If lines px + qy + r = 0, qx + ry + p = 0 and rx + py + q = 0 are

concurrent, then prove that p + q + r = 0 (where p, q, r are distinct).

Watch Video Solution

412. the diagonals of the parallelogram formed by the the lines $a_1x + b_1y + c_1 = 0$, $a_1x + b_1y + c_1' = 0$, $a_2x + b_2y + c_1 = 0$, $a_$

413. If the lines joining the origin and the point of intersection of curves $ax^2 + 2hxy + by^2 + 2gx + 0$ and $a_1x^2 + 2h_1xy + b_1y^2 + 2g_1x = 0$ are mutually perpendicular, then prove that $g(a_1 + b_1) = g_1(a + b)$.

Watch Video Solution

414. Prove that the angle between the lines joining the origin to the points of intersection of the straight line y = 3x + 2 with the curve

$$x^2+2xy+3y^2+4x+8y-11=0 ext{ is } an^{-1}igg(rac{2\sqrt{2}}{3}igg)$$

Watch Video Solution

415. Prove that the straight lines joining the origin to the point of intersection of the straight line hx + ky = 2hk and the curve $(x - k)^2 + (y - h)^2 = c^2$ are perpendicular to each other if $h^2 + k^2 = c^2$.

416. If
$$x^2 - 2pxy - y^2 = 0$$
 and $x^2 - 2qxy - y^2 = 0$ bisect angles

between each other, then find the condition.



 $ax^2+5xy+2y^2=0$ are mutually perpendicular.

Watch Video Solution

418. Find the acute angle between the pair of lines represented by $(x\coslpha-y\sinlpha)^2=ig(x^2+y^2ig)\sin^2lpha.$

419. If the angle between the two lines represented by $2x^2 + 5xy + 3y^2 + 6x + 7y + 4 = 0$ is $\tan^{-1}(m)$, then find the value of m.

420. If the pair of straight lines $ax^2 + 2hxy + by^2 = 0$ is rotated about the origin through 90^0 , then find the equations in the new position.

Watch Video Solution

421. The orthocentre of the triangle formed by the lines xy=0 and

$$x+y=1$$
 is $\left(rac{1}{2},rac{1}{2}
ight)$ (b) $\left(rac{1}{3},rac{1}{3}
ight)$ (0, 0) (d) $\left(rac{1}{4},rac{1}{4}
ight)$

422. The lines joining the origin to the point of intersection of $3x^2 + mxy - 4x + 1 = 0$ and 2x + y - 1 = 0 are at right angles. Then which of the following is a possible value of m? -4 (b) 4 (c) 7 (d) 3

423. If the slope of one line is double the slope of another line and the

combined equation of the pair of lines is $\left(\frac{x^2}{a}\right) + \left(\frac{2xy}{h}\right) + \left(\frac{y^2}{b}\right) = 0$, then find the ratio ab : h^2 .

Watch Video Solution

424. Find the combined equation of the pair of lines through the point (1,

0) and parallel to the lines represented by $2x^2-xy-y^2=0$

425. The value k for which $4x^2 + 8xy + ky^2 = 9$ is the equation of a pair of straight lines is



426. The two lines represented by $3ax^2 + 5xy + (a^2 - 2)y^2 = 0$ are perpendicular to each other for two values of a (b) a for one value of a (d) for no values of a

Watch Video Solution

427. If two lines represented by $x^4 + x^3y + cx^2y^2 - xy^3 + y^4 = 0$

bisector of the angle between the other two, then the value of c is (a)0 (b)

-1 (c) 1 (d) -6

428. The straight lines represented by $x^2 + mxy - 2y^2 + 3y - 1 = 0$ meet at (a) $\left(-\frac{1}{3}, \frac{2}{3}\right)$ (b) $\left(-\frac{1}{3}, -\frac{2}{3}\right)$ (c) $\left(\frac{1}{3}, \frac{2}{3}\right)$ (d) none of these

Watch Video Solution

429. The straight lines represented by the equation $135x^2 - 136xy + 33y^2 = 0$ are equally inclined to the line (a) x - 2y = 7 (b) x+2y=7 (c) x - 2y = 4 (d) 3x + 2y = 4

Watch Video Solution

430. If one of the lines of $my^2+ig(1-m^2ig)xy-mx^2=0$ is a bisector of the angle between the lines xy=0 , then m is (a)1 (b) 2 (c) $-rac{1}{2}$ (d) -1

431. Statement 1 : If -2h = a + b, then one line of the pair of lines $ax^2 + 2hxy + by^2 = 0$ bisects the angle between the coordinate axes in the positive quadrant. Statement 2 : If ax + y(2h + a) = 0 is a factor of $ax^2 + 2hxy + by^2 = 0$, then b + 2h + a = 0 Both the statements are true but statement 2 is the correct explanation of statement 1. Both the statements are true but statement 2 is not the correct explanation of statement 1 is true and statement 2 is false. Statement 1 is false and statement 2 is true.

Watch Video Solution

432. Show that all chords of the curve $3x^2 - y^2 - 2x + 4y = 0$, which subtend a right angle at the origin, pass through a fixed point. Find the coordinates of the point.



434. The distance between the lines $\left(x+7y
ight)^2+4\sqrt{2}(x+7y)-42=0$

is_____

Watch Video Solution

435. x+y=7 and $ax^2+2hxy+ay^2=0,\,(a
eq 0)$, are three real

distinct lines forming a triangle. Then the triangle is

(a)isosceles (b) scalene equilateral (d) right angled

436. If the slope of one of the lines represented by $ax^2 + 2hxy + by^2 = 0$ is the square of the other, then $\frac{a+b}{h} + \frac{8h^2}{ab} = 4$ (b) 6 (c) 8 (d) none of these

Watch Video Solution

437.
$$\int \left\{ \frac{2-3\sin x}{\cos^2 x} \right\}$$

Watch Video Solution

438. The sides of a triangle have the combined equation $x^2 - 3y^2 - 2xy + 8y - 4 = 0$. The third side, which is variable, always passes through the point (-5, -1). Find the range of values of the slope of the third line such that the origin is an interior point of the triangle.

439. Let PQR be a right-angled isosceles triangle, right angled at P(2, 1). If the equation of the line QR is 2x + y = 3, then the equation representing the pair of lines PQ and PR is $3x^2 - 3y^2 + 8xy + 20x + 10y + 25 = 0$ $3x^2 - 3y^2 + 8xy - 20x - 10y + 25 = 0$ $3x^2 - 3y^2 + 8xy + 10x + 15y + 20 = 0$ $3x^2 - 3y^2 - 8xy - 15y - 20 = 0$

Watch Video Solution

440. The combined equation of three sides of a triangle is $(x^2 - y^2)(2x + 3y - 6) = 0$. If (-2, a) is an interior point and (b, 1)

is an exterior point of the triangle, then

Watch Video Solution

441. Find the equation of the bisectors of the angles between the lines joining the origin to the point of intersection of the straight line

x-y=2 with the curve $5x^2+11xy-8y^2+8x-4y+12=0$

Watch Video Solution

442. If θ is the angle between the lines given by the equation $6x^2 + 5xy - 4y^2 + 7x + 13y - 3 = 0$, then find the equation of the line passing through the point of intersection of these lines and making an angle θ with the positive x-axis.

Watch Video Solution

443. The distance of a point (x_1, y_1) from two straight lines which pass through the origin of coordinates is p. Find the combined equation of these straight lines.



444. Prove that the product of the perpendiculars from (α, β) to the pair

of lines
$$ax^2+2hxy+by^2=0$$
 is $\displaystyle rac{alpha^2+2hlphaeta+beta^2}{\sqrt{\left(a-b
ight)^2+4h^2}}$

Watch Video Solution



446. Show that the pairs of straight lines $2x^2 + 6xy + y^2 = 0$ and $4x^2 + 18xy + y^2 = 0$ are equally inclined,then b is equal to

Watch Video Solution

447. The product of the perpendiculars from origin to the pair of lines

$$ax^2+2hxy+by^2+2gx+2fy+c=0$$
 is

448. Find the angle between the straight lines joining the origin to the point of intersection of $3x^2 + 5xy - 3y^2 + 2x + 3y = 0$ and 3x - 2y = 1

Watch Video Solution

449. Through a point A(2,0) on the x-axis, a straight line is drawn parallel to the y-axis so as to meet the pair of straight lines $ax^2 + 2hxy + by^2 = 0$ at B and C. If AB = BC, then (a) $h^2 = 4ab$ (b) $8h^2 = 9ab$ (c) $9h^2 = 8ab$ (d) $4h^2 = ab$



451. Does equation $x^2 + 2y^2 - 2\sqrt{3}x - 4y + 5 = 0$ satisfies the condition $abc + 2gh - af^2 - bg^2 - ch^2 = 0$? Does it represent a pair of straight lines?

Watch Video Solution

452. Find the value of λ if $2x^2 + 7xy + 3y^2 + 8x + 14y + \lambda = 0$ represents a pair of straight lines.

Watch Video Solution

453. Find the distance between the pair of parallel lines $x^2 + 4xy + 4y^2 + 3x + 6y - 4 = 0$

454. If the pair of lines $ax^2+2hxy+by^2+2gx+2fy+c=0$ intersect on the y-axis, then prove that $2fgh=bg^2+ch^2$

455. Find the lines whose combined equation is $6x^2 + 5xy - 4y^2 + 7x + 13y - 3 = 0$ using the concept of parallel lines through the origin.

Watch Video Solution

456. If the component lines whose combined equation is $px^2 - qxy - y^2 = 0$ make the angles α and β with x-axis, then find the value of tan $(\alpha + \beta)$.







four times their product, then the value of c is_____

Watch Video Solution

459. The distance between the two lines represented by the sides of an

equilateral triangle a right-angled triangle an isosceles triangle



460. If the gradient one of the lines $x^2 + hxy + 2y^2 = 0$ is twice that of

the other, then $h = _$ __ _

Watch Video Solution

461. If one of the lines of $my^2 + ig(1-m^2ig)xy - mx^2 = 0$ is a bisector of

the angle between the lines xy = 0, then m is

Watch Video Solution

462. Two pairs of straight lines have the equations $y^2 + xy - 12x^2 = 0$ and $ax^2 + 2hxy + by^2 = 0$. One line will be

common among them if

463. If the equation of the pair of straight lines passing through the point (1, 1), one making an angle θ with the positive direction of the x-axis and the other making the same angle with the positive direction of the y-axis, is $x^2 - (a+2)xy + y^2 + a(x+y-1) = 0, a \neq 2$, then the value of $\sin 2\theta$ is a - 2 (b) a + 22(a + 2) (d) $\frac{2}{a}$

Watch Video Solution

464. If one of the lines given by the equation $2x^2 + pxy + 3y^2 = 0$ coincide with one of those given by $2x^2 + qxy - 3y^2 = 0$ and the other lines represented by them are perpendicular, then value of p + q is

Watch Video Solution

465. If $x^2 + 2hxy + y^2 = 0$ represents the equation of the straight lines through the origin which make an angle α with the straight line y + x = 0

(a)
$$\sec 2\alpha = h$$
 (b) $= \sqrt{\frac{(1+h)}{(2h)}}$ (c) $2\sin \alpha = \sqrt{\frac{(1+h)}{h}}$ (d) $\cot \alpha$
 $= \sqrt{\frac{(1+h)}{(h-1)}}$

Watch Video Solution

466. The equation to a pair of opposite sides of a parallelogram are $x^2 - 5x + 6 = 0$ and $y^2 - 6y + 5 = 0$. The equations to its diagonals are x + 4y = 13, y = 4x - 7 (b) 4x + y = 13, 4y = x - 74x + y = 13, y = 4x - 7 (d) y - 4x = 13, y + 4x - 7

Watch Video Solution

467. The equation $a^2x^2+2h(a+b)xy+b^2y^2=0$ and $ax^2+2hxy+by^2=0$ represent

A. two pairs of perpendicular straight lines

B. two pairs of parallel straight lines

C. two pairs of straight lines which are equally inclined to each other

D. none of these

Answer: null

Watch Video Solution

468. The equation $x^3 + x^2y - xy^2 = y^3$ represents (a)three real straight lines (b)lines in which two of them are perpendicular to each other (c)lines in which two of them are coincident (d)none of these

Watch Video Solution

469. The image of the pair of lines represented by $ax^2 + 2hxy + by^2 = 0$ by the line mirror y = 0 is $ax^2 - 2hxy - by^2 = 0$ $bx^2 - 2hxy + ay^2 = 0$ $bx^2 + 2hxy + ay^2 = 0$ $ax^2 - 2hxy + by^2 = 0$

470. The combined equation of the lines l_1andl_2 is $2x^2 + 6xy + y^2 = 0$ and that of the lines m_1andm_2 is $4x^2 + 18xy + y^2 = 0$. If the angle between l_1 and m_2 is α then the angle between l_2andm_1 will be

471. If the equation $ax^2 - 6xy + y^2 = 0$ represents a pair of lines whose slopes are m and m^2 , then the value(s) of a is/are

Watch Video Solution

472. The equation of a line which is parallel to the line common to the pair of lines given by $6x^2 - xy - 12y^2 = 0$ and $15x^2 + 14xy - 8y^2 = 0$ and at a distance of 7 units from it is 3x - 4y = -35 5x - 2y = 73x + 4y = 35 2x - 3y = 7

473. If the sum of the slopes of the lines given by $x^2 - 2cxy - 7y^2 = 0$ is

four times their product, then the value of c is_____



474. Area of the triangle formed by the line x + y = 3 and angle bisectors of the pair of straight lines $x^2 - y^2 + 2y = 1$ is 2squares b. 4squares c. 6squares d. 8squares

Watch Video Solution

475. The equation $x^2y^2 - 9y^2 - 6x^2y + 54y = 0$ represents (a)a pair of straight lines and a circle (b)a pair of straight lines and a parabola (c)a set of four straight lines forming a square (d)none of these

476. The straight lines represented by $(y-mx)^2=a^2\bigl(1+m^2\bigr)$ and $(y-nx)^2=a^2\bigl(1+n^2\bigr)$ from a rectangle (b) rhombus trapezium (d) none of these

Watch Video Solution

477. If the pairs of lines $x^2 + 2xy + ay^2 = 0$ and $ax^2 + 2xy + y^2 = 0$ have exactly one line in common, then the joint equation of the other two lines is given by $3x^2 + 8xy - 3y^2 = 0$ $3x^2 + 10xy + 3y^2 = 0$ $y^2 + 2xy - 3x^2 = 0$ $x^2 + 2xy - 3y^2 = 0$

Watch Video Solution

478. The condition that one of the straight lines given by the equation $ax^2 + 2hxy + by^2 = 0$ may coincide with one of those given by the equation $a'x^2 + 2h'xy + b'y^2 = 0$ is $(ab' - a'b)^2 = 4(ha' - h'a)(bh' - b'h)$ $(ab' - a'b)^2 = (ha' - h'a)(bh' - b'h)$

$$ig(ha'-h'aig)^2 = 4ig(ab'-a'big)ig(bh'-b'hig) \ ig(bh'-b'hig)^2 = 4ig(ab'-a'big)ig(ha'-h'aig) \ ig(ha'-h'aig)$$

Watch Video Solution

479. If the represented by the equation $3y^2 - x^2 + 2\sqrt{3}x - 3 = 0$ are rotated about the point $(\sqrt{3}, 0)$ through an angle of 15^0 , on in clockwise direction and the other in anticlockwise direction, so that they become perpendicular, then the equation of the pair of lines in the new position

is $y^2-x^2+2\sqrt{3}x+3=0$ $y^2-x^2+2\sqrt{3}x-3=0$ $y^2-x^2-2\sqrt{3}x+3=0$ $y^2-x^2+3=0$

Watch Video Solution

480. A point moves so that the distance between the foot of perpendiculars from it on the lines $ax^2 + 2hxy + by^2 = 0$ is a constant 2d. Show that the equation to its locus is $(x^2 + y^2)(h^2 - ab) = d^2 \Big\{ (a - b)^2 + 4h^2 \Big\}$.



481. The angle between the pair of lines whose equation is $4x^2 + 10xy + my^2 + 5x + 10y = 0$ is(a) $\tan^{-1}\left(\frac{3}{8}\right)$ (b) $\tan^{-1}\left(\frac{3}{4}\right)$ (c) $\tan^{-1}\left\{2\frac{\sqrt{25-4m}}{m+4}\right\}, m \in R$ (d) none of these **Vatch Video Solution**

482. Find the point of intersection of the pair of straight lines represented by the equation $6x^2 + 5xy - 21y^2 + 13x + 38y - 5 = 0$.

Watch Video Solution

483. Find the angle between the lines represented by
$$x^2 + 2xy\sec heta + y^2 = 0.$$

484. If the pair of lines $\sqrt{3}x^2 - 4xy + \sqrt{3}y^2 = 0$ is rotated about the origin by $\frac{\pi}{6}$ in the anticlockwise sense, then find the equation of the pair in the new position.

Watch Video Solution

485. If the equation $2x^2 + kxy + 2y^2 = 0$ represents a pair of real and

distinct lines, then find the values of k.

Watch Video Solution

486. If the equation $x^2 + (\lambda + \mu)xy + \lambda uy^2 + x + \mu y = 0$ represents two parallel straight lines, then prove that $\lambda = \mu$.

Watch Video Solution

487. If one of the lines of the pair $ax^2 + 2hxy + by^2 = 0$ bisects the angle between the positive direction of the axes. Then find the relation

for a, b and h.

Watch Video Solution

488. Prove that the equation $2x^2 + 5xy + 3y^2 + 6x + 7y + 4 = 0$ respresents a pair of straight lines .Find the coordinates of their point of intersection and also the angle between them.

Watch Video Solution

489. A line L passing through the point (2, 1) intersects the curve $4x^2 + y^2 - x + 4y - 2 = 0$ at the point AandB. If the lines joining the origin and the points A, B are such that the coordinate axes are the bisectors between them, then find the equation of line L.



491. If one of the lines denoted by the line pair $ax^2 + 2hxy + by^2 = 0$ bisects the angle between the coordinate axes, then prove that $(a+b)^2 = 4h^2$