



India's Number 1 Education App

## MATHS

### BOOKS - CENGAGE MATHS (ENGLISH)

#### DIFFERENTIATION

##### Illustration

1. If  $y = \sqrt{\frac{1 - \cos 2x}{1 + \cos 2x}}$ ,  $x \in \left(0, \frac{\pi}{2}\right) \cup \left(\frac{\pi}{2}, \pi\right)$ , then find  $\frac{dy}{dx}$ .



Watch Video Solution

2. Find the derivative of  $e^{\sqrt{x}}$  w.r.t.  $x$  using the first principle.



Watch Video Solution

3. If  $f(x) = x \tan^{-1} x$ , find  $f'(\sqrt{3})$  using the first principle.



Watch Video Solution

4. If  $f(x) = [2x] \sin 3\pi x$  then prove that  $f(k^+) = 6k\pi(-1)^k$ , (where  $[.]$  denotes the greatest integer function and  $k \in N$ ).



Watch Video Solution

5. Let  $f: R \rightarrow$  satisfying  $|f(x)| \leq x^2 \forall x \in R$  be differentiable at  $x = 0$ . Then find  $f'(0)$



Watch Video Solution

6. A function  $f: R \rightarrow R$  satisfies the equation  $f(x+y) = f(x)f(y)$  for all  $x, y \in R$  and  $f(x) \neq 0$  for all  $x \in R$ . If  $f(x)$  is differentiable at  $x = 0$  and  $f'(0) = 2$ , then prove that  $f'(x) = 2f(x)$



Watch Video Solution

7. If  $y = \left(1 + x^{\frac{1}{4}}\right) \left(1 + x^{\frac{1}{2}}\right) \left(1 - x^{\frac{1}{4}}\right)$ , then find  $\frac{dy}{dx}$ .



Watch Video Solution

8. If  $f(x) = x|x|$ , then prove that  $f'(x) = 2|x|$



Watch Video Solution

9. If  $y = 1 + \frac{x}{1!} + \frac{x^2}{2!} + \frac{x^3}{3!} + \dots + \frac{x^n}{n!}$ , show that  $\frac{dy}{dx} - y + \frac{x^n}{n!} = 0$ .



Watch Video Solution

10. Find  $\frac{dy}{dx}$  for  $y = \sin^{-1}(\cos x)$ , where  $x \in (0, 2\pi)$



Watch Video Solution

11. Differentiate  $\sin^{-1}\left(2x\sqrt{1-x^2}\right)$  with respect to  $x$ , if  $-1/\sqrt{2}$



Watch Video Solution

12.  $y = \tan^{-1}((a\cos x - b\sin x)/(b\cos x + a\sin x))$ , where  $e^{-\pi/2} < 1$



Watch Video Solution

13.  $y = \sin^{-1}\left(\frac{x}{\sqrt{1+x^2}}\right) + \cos^{-1}\left(\frac{1}{\sqrt{1+x^2}}\right)$ , where  $0 < x < \infty$  Find  $\frac{dy}{dx}$



Watch Video Solution

14. Find  $\frac{dy}{dx}$  for  $y = \sin(x^2 + 1)$



Watch Video Solution

$$15. \text{ If } y = \sqrt{\log \left\{ \sin \left( \frac{x^2}{3} - 1 \right) \right\}}, \text{ then } f \in d \frac{dy}{dx}$$



Watch Video Solution

$$16. \text{ Differentiate the function } f(x) = \sec(\tan(\sqrt{x})) \text{ with respect to } x$$



Watch Video Solution

$$17. \text{ Find } \frac{dy}{dx} \text{ for } y = \log \left( x + \sqrt{a^2 + x^2} \right).$$



Watch Video Solution

$$18. y = \tan^{-1} \left( \frac{\sqrt{1+x^2} + \sqrt{1-x^2}}{\sqrt{1+x^2} - \sqrt{1-x^2}} \right),$$



Watch Video Solution

19. Find  $\frac{dy}{dx}$  for  $y = \tan^{-1} \sqrt{\frac{a-x}{a+x}}$ ,  $-a < x < a$



Watch Video Solution

20. If  $y = \sin^{-1} \left[ x\sqrt{1-x} - \sqrt{x}\sqrt{1-x^2} \right]$  and '0



Watch Video Solution

21. If  $y = \frac{\tan^{-1} 1}{1+x+x^2} + \frac{\tan^{-1} 1}{x^2+3x+3} + \frac{\tan^{-1} 1}{x^2+5x+7} + \dots$  upto  $n$  terms, then

find the value of  $y'(0)$



Watch Video Solution

22. Let  $f: \vec{R} \rightarrow \vec{R}$  be a one-one onto differentiable function, such that

$f(2) = 1$  and  $f'(2) = 3$ . Then find the value of  $\left( \left( \frac{d}{dx} (f^{-1}(x)) \right) \right)_{x=1}$



Watch Video Solution

23. If  $f(x) = \cos x \cos 2x \cos 4x \cos(8x) \cdot \cos 16x$  then find  $f\left(\frac{\pi}{4}\right)$



Watch Video Solution

24. If  $\cos y = x \cos(a + y)$ , with  $\cos a \neq \pm 1$ , prove that  $\frac{dy}{dx} = \frac{\cos^2(a + y)}{\sin a}$



Watch Video Solution

25. Find  $\frac{dy}{dx}$  or  $y = x \sin x \log x$



Watch Video Solution

26. Evaluate  $(\lim)_{h \rightarrow 0} \frac{(a + h)^2 \sin^{-1}(a + h) - a^2 \sin^{-1}a}{h}$



Watch Video Solution

27. Differentiate  $y = \frac{e^x}{1 + \sin x}$



[Watch Video Solution](#)

28. If  $f(x) = \cos x \cdot \cos 2x \cdot \cos 4x \cdot \cos 8x \cdot \cos 16x$ , then find  $f\left(\frac{\pi}{4}\right)$ .



[Watch Video Solution](#)

29. If  $\cos y = x \cos(a + y)$ , with  $\cos a \neq \pm 1$ , prove that  $\frac{dy}{dx} = \frac{\cos^2(a + y)}{\sin a}$



[Watch Video Solution](#)

30. If  $y = \sqrt{\frac{1-x}{1+x}}$ , prove that  $(1-x^2)\frac{dy}{dx} + y = 0$ .



[Watch Video Solution](#)

31. Find the sum of the series  $(1 + 2x + 3x^2 + (n - 1)x^{n-2})$  using differentiation.



Watch Video Solution

32. If  $\sqrt{x} + \sqrt{y} = 4$ , then find  $\frac{dy}{dx}$ .



Watch Video Solution

33. If  $xy + y^2 = \tan x + y$ , then find  $\frac{dy}{dx}$ .



Watch Video Solution

34. If  $y = x + \frac{1}{x + \frac{1}{x + \frac{1}{x + \dots}}}$ , prove that  $\frac{dy}{dx} = \frac{y}{2y - x}$ .



Watch Video Solution

**35.** If  $\sec(x + y) = xy$  find  $dy/dx$

 **Watch Video Solution**

**36.** If  $y = \sqrt{\sin x + \sqrt{\sin x + \sqrt{\sin x + \dots}}}$ , prove that  $\frac{dy}{dx} = \frac{\cos x}{2y - 1}$

 **Watch Video Solution**

**37.** If  $\log(x^2 + y^2) = 2\tan^{-1}\left(\frac{y}{x}\right)$ , show that  $\frac{dy}{dx} = \frac{x + y}{x - y}$

 **Watch Video Solution**

**38.** If  $y = y(x)$  and it follows the relation  $4xe^{xy} = y + 5\sin^2 x$ , then  $y'(0)$  is equal to \_\_\_\_\_

 **Watch Video Solution**

39. Find  $\frac{dy}{dx}$  if  $x = a(\theta - \sin\theta)$  and  $y = a(1 - \cos\theta)$



Watch Video Solution

40. If  $x = a \sec^3\theta$  and  $y = a \tan^3\theta$ , find  $\frac{dy}{dx}$  at  $\theta = \frac{\pi}{3}$



Watch Video Solution

41. If  $x = e^{\cos 2t}$  and  $y = e^{\sin 2t}$ , then move that  $\frac{dy}{dx} = -\frac{y \log x}{x \log y}$ .



Watch Video Solution

42. Let  $y = x^3 - 8x + 7$  and  $x = f(t) \cdot \frac{f(dy)}{dx} = 2$  and  $x = 3a = 0$ , then find the value of  $\frac{dx}{dt}$  when  $a = 0$ .



Watch Video Solution

43. Find the derivative of  $\frac{\sqrt{x}(x+4)^{\frac{3}{2}}}{(4x-3)^{\frac{4}{3}}}$



Watch Video Solution

44. If  $x^m y^n = (x+y)^{m+n}$ , prove that  $\frac{dy}{dx} = \frac{y}{x}$



Watch Video Solution

45. Differentiate  $(\log x)^{\cos x}$  with respect to  $x$



Watch Video Solution

46. If  $f(x) = |x|^{\sin x}|$ , then find  $f' \left( -\frac{\pi}{4} \right)$



Watch Video Solution

47. If  $y = x^x \wedge x \in ((\infty))$ , find  $\frac{dy}{dx}$ .



[Watch Video Solution](#)

48. If  $f(x) = \lim_{h \rightarrow 0} \frac{(\sin(x+h))^{\log_e(x+h)} - (\sin x)^{\log_e x}}{h}$  then find  $f(\pi/2)$ .



[Watch Video Solution](#)

49. If  $x < 1$ , prove that  $\frac{1}{1+x} + \frac{2x}{1+x^2} + \frac{4x^3}{1+x^4} + \dots = \frac{1}{1-x}$



[Watch Video Solution](#)

50. Differentiate  $\log \sin x$  w.r.t.  $\sqrt{x}$ .



[Watch Video Solution](#)

51. Differentiate  $\tan^{-1}\left(\frac{\sqrt{1+x^2}-1}{x}\right)$  with respect to  $\sin^{-1}\left(\frac{2x}{1+x^2}\right)$ ,



Watch Video Solution

52. Find the derivative of  $f(\tan x) \text{ wrt } g(\sec x) \text{ at } x = \frac{\pi}{4}$ , where  $f'(1) = 2$  and  $g'(\sqrt{2}) = 4$ .



Watch Video Solution

53. Let  $= \begin{vmatrix} \cos x & \sin x & \cos x \\ \cos 2x & \sin 2x & 2\cos 2x \\ \cos 3x & \sin 3x & 3\cos 3x \end{vmatrix}$  then find the values of  $f(0)$  and  $f'(\pi/2)$ .



Watch Video Solution

54.  $f(x) = \begin{vmatrix} \cos x & x & 1 \\ 2\sin x & x^2 & 2x \\ \tan x & x & 1 \end{vmatrix}$ . Then find the value of  $\lim_{x \rightarrow 0} \frac{f(x)}{x}$ .



**Watch Video Solution**

55. If  $y = \cos^{-1}x$ , find  $\frac{d^2y}{dx^2}$  in terms of  $y$  alone.



**Watch Video Solution**

56. about to only mathematics



**Watch Video Solution**

57. If  $y = x \log \left\{ \frac{x}{(a + bx)} \right\}$ , then show that  $x^3 \frac{d^2y}{dx^2} = \left( x \frac{dy}{dx} - y \right)^2$



**Watch Video Solution**

$$\left[ 1 + \left( \frac{dy}{dx} \right)^2 \right]^{\frac{3}{2}}$$

58. If  $(x - a)^2 + (y - b)^2 = c^2$ , for some  $c > 0$ , prove that  $\frac{\frac{d^2y}{dx^2}}{\left[ 1 + \left( \frac{dy}{dx} \right)^2 \right]^{\frac{3}{2}}}$  is a constant independent of a and b.



**Watch Video Solution**

59. If  $y = e^a \cos^{(-1)x}$ ,  $-1 \leq x < 1$ , show that  $\left(1 - x^2\right) \frac{d^2y}{dx^2} - x \frac{dy}{dx} - a^2 y = 0$



**Watch Video Solution**

60. If  $x = a(\cos t + t \sin t)$  and  $y = a(\sin t - t \cos t)$ , find  $(d^2y)/(dx^2)$ .



**Watch Video Solution**

61. If g is inverse of f then prove that  $f'(g(x)) = -g''(x)(f(g(x)))^3$ .



**Watch Video Solution**

**62.** Let  $f(x)$  and  $g(x)$  be real valued functions such that  $f(x)g(x)=1$ ,  $\forall x \in R$ . If  $f'(x)$  and  $g''(x)$  exists  $\forall x \in R$  and  $f'(x)$  and  $g'(x)$  are never zero, then prove that  $\frac{f'(x)}{f(x)} - \frac{g''(x)}{g'(x)} = \frac{2f(x)}{f'(x)}$

 Watch Video Solution

**63.** If  $f(x)$ ,  $g(x)$  and  $h(x)$  are three polynomial of degree 2, then prove that

$$\phi(x) = \begin{vmatrix} f(x) & g(x) & h(x) \\ f'(x) & g'(x) & h'(x) \\ f''(x) & g''(x) & h''(x) \end{vmatrix}$$
 is a constant polynomial.

 Watch Video Solution

**64.** If  $f\left(\frac{x+y}{3}\right) = \frac{2+f(x)+f(y)}{3}$  for all real  $x$  and  $y$  and  $f'(2)=2$ , then determine  $y=f(x)$ .

 Watch Video Solution

**65.**

If  $f(x) + f(y) = f\left(\frac{x+y}{1-xy}\right)$  for all  $x, y \in R$ , ( $xy \neq 1$ ), and  $\lim_{x \rightarrow 0} \frac{f(x)}{x} = 2$  then



**Watch Video Solution**

**66.** Let  $f: R - \{0\} \rightarrow R$  be a function which is differentiable in its domain

and satisfying the equation  $f(x+y) = f(x) + f(y) + \frac{x+y}{xy} - \frac{1}{x+y}$ , also

$f'(1)=2$ . Then find the function.



**Watch Video Solution**

**67.** Find function  $f(x)$  which is differentiable and satisfy the relation

$f(x+y) = f(x) + f(y) + (e^x - 1)(e^y - 1) \forall x, y \in R$ , and  $f(0) = 2$ .



**Watch Video Solution**

68. If  $f\left(\frac{x}{y}\right) = \frac{f(x)}{f(y)}$ ,  $\forall y, f(y) \neq 0$  and  $f(1) = 2$ , find  $f(x)$ .

 Watch Video Solution

69. Let  $f: \vec{R} \rightarrow R$  be a function satisfying condition  
 $f(x + y^3) = f(x) + [f(y)]^3$  for all  $x, y \in R$ . If  $f'(0) \geq 0$ , find  $f(10)$ .

 Watch Video Solution

70. Let  $f(x + y) = f(x) + f(y) + 2xy - 1$  for all real  $x$  and  $y$  and  $f(x)$  be a differentiable function. If  $f'(0) = \cos\alpha$ , prove that  $f(x) > 0 \forall x \in R$ .

 Watch Video Solution

71. If  $y = \sqrt{\frac{1 - \cos 2x}{1 + \cos 2x}}$ ,  $x \in \left(0, \frac{\pi}{2}\right) \cup \left(\frac{\pi}{2}, \pi\right)$ , then find  $\frac{dy}{dx}$ .

 Watch Video Solution

72. Find the derivative of  $e^{\sqrt{x}}$  w.r.t.  $x$  using the first principle.



Watch Video Solution

73. If  $f(x) = x \tan^{-1}x$ , find  $f'(\sqrt{3})$  using the first principle.



Watch Video Solution

74. If  $f(x) = [2x]\sin 3\pi x$  then prove that  $f'(k^+) = 6k\pi(-1)^k$ , (where  $[.]$  denotes the greatest integer function and  $k \in \mathbb{N}$ ).



Watch Video Solution

75. Let  $f: R \rightarrow R$  satisfying  $|f(x)| \leq x^2$ ,  $\forall x \in R$ , differentiable at  $x = 0$  then find  $f'(0)$



Watch Video Solution

**76.** A function  $f: R \rightarrow R$  satisfies the equation  $f(x+y)=f(x)f(y)$  for all  $x, y \in R$  and  $f(x) \neq 0$  for all  $x \in R$ . If  $f(x)$  is differentiable at  $x=0$  and  $f'(0)=2$ , then prove that  $f'(x) = 2f(x)$ .



**Watch Video Solution**

**77.** If  $y = \left(1 + x^{\frac{1}{4}}\right)\left(1 + x^{\frac{1}{2}}\right)\left(1 - x^{\frac{1}{4}}\right)$ , then find  $\frac{dy}{dx}$ .



**Watch Video Solution**

**78.** If  $f(x) = x|x|$ , then prove that  $f'(x) = 2|x|$



**Watch Video Solution**

**79.** If  $y = 1 + \frac{x}{1!} + \frac{x^2}{2!} + \frac{x^3}{3!} + \dots + \frac{x^n}{n!}$ , show that  $\frac{dy}{dx} - y + \frac{x^n}{n!} = 0$ .



**Watch Video Solution**



Watch Video Solution

80. Find  $\frac{dy}{dx}$  for  $y = \sin^{-1}(\cos x)$ ,  $x \in (0, \pi) \cup (\pi, 2\pi)$ .



Watch Video Solution

81. Differentiate  $\sin^{-1}\left(2x\sqrt{1-x^2}\right)$  with respect to  $x$  if  $\frac{1}{\sqrt{2}} < x < 1$



Watch Video Solution

82.

If  $y = \tan^{-1}\left(\frac{a\cos x - b\sin x}{b\cos x + a\sin x}\right)$ , where  $-\frac{\pi}{2} < x < \frac{\pi}{2}$  and  $\frac{a}{b}\tan x > -1$ , then find



Watch Video Solution

83.  $y = \sin^{-1}\left(\frac{x}{\sqrt{1+x^2}}\right) + \cos^{-1}\left(\frac{1}{\sqrt{1+x^2}}\right)$ , Find  $dy/dx$



Watch Video Solution

84. Find  $\frac{dy}{dx}$  for  $y = \sin(x^2 + 1)$ .



Watch Video Solution

85. If  $y = \sqrt{\log\left\{\sin\left(\frac{x^3}{3} - 1\right)\right\}}$ , then find  $\frac{dy}{dx}$ .



Watch Video Solution

86. Differentiate the function  $f(x) = \sec(\tan\sqrt{x})$  with respect to  $x$ ,



Watch Video Solution

87. Find  $\frac{dy}{dx}$  for  $y = \log\left(x + \sqrt{a^2 + x^2}\right)$ .



Watch Video Solution

$$88. y = \tan^{-1} \left( \frac{\sqrt{1+x^2} + \sqrt{1-x^2}}{\sqrt{1+x^2} - \sqrt{1-x^2}} \right), \text{ where } -1$$



Watch Video Solution

$$89. \text{Find } \frac{dy}{dx} \text{ for } y = \tan^{-1} \left\{ \sqrt{\frac{a-x}{a+x}} \right\}, \text{ where } -a < x < a$$



Watch Video Solution

$$90. \text{if } y = \sin^{-1} \left[ x\sqrt{1-x} - \sqrt{x}\sqrt{1-x^2} \right] \text{ and } 0 < x < 1, \text{ then find } \frac{dy}{dx}.$$



Watch Video Solution

$$91. \text{If } y = \tan^{-1} \frac{1}{1+x+x^2} + \tan^{-1} \frac{1}{x^2+3x+3} + \tan^{-1} \frac{1}{x^2+5x+7} + \dots \text{ upto n terms, then find the value of } y'(0).$$



Watch Video Solution

92. Let  $f: \mathbb{R} \rightarrow \mathbb{R}$  be a one-one onto differentiable function, such that

$f(2) = 1$  and  $f'(2) = 3$ . Then find the value of  $\left( \left( \frac{d}{dx} (f^{-1}(x)) \right) \right)_{x=1}$



Watch Video Solution

93. If  $f(x) = \cos x \cos 2x \cos 4x \cos(8x) \dots \cos 16x$  then find  $f' \left( \frac{\pi}{4} \right)$



Watch Video Solution

94. If  $\cos y = x \cos(a + y)$ , with  $\cos a \neq \pm 1$ , prove that  $\frac{dy}{dx} = \frac{\cos^2(a + y)}{\sin a}$



Watch Video Solution

95. Find  $\frac{dy}{dx}$  for  $y = x \sin x \log x$



Watch Video Solution

96. Evaluate (  $\lim_{h \rightarrow 0}$  )  $\frac{(a + h)^2 \sin^{-1}(a + h) - a^2 \sin^{-1}a}{h}$



Watch Video Solution

97. Differentiate  $y = \frac{e^x}{1 + \sin x}$



Watch Video Solution

98. If  $f(x) = \cos x \cos 2x \cos 4x \cos 8x \cos 16x$  then find  $f\left(\frac{\pi}{4}\right)$



Watch Video Solution

99. If  $\cos y = x \cos(a + y)$ , with  $\cos a \neq \pm 1$ , prove that  $\frac{dy}{dx} = \frac{\cos^2(a + y)}{\sin a}$



Watch Video Solution

100. If  $y = \sqrt{\frac{1-x}{1+x}}$ , prove that  $(1-x^2)\frac{dy}{dx} + y = 0$



**Watch Video Solution**

101. Find the sum of the series  $1 + 2x + 3x^2 + (n-1)x^{n-2}$  using differentiation.



**Watch Video Solution**

102. If  $\sqrt{x} + \sqrt{y} = 4$ , then find  $\frac{dy}{dx}$ .



**Watch Video Solution**

103. If  $xy + y^2 = \tan x + y$ , then find  $\frac{dy}{dx}$ .



**Watch Video Solution**

**104.** If  $y = x + \frac{1}{x + \frac{1}{x + \frac{1}{x^+}}}$ , prove that  $\frac{dy}{dx} = \frac{y}{2y - x}$ .



**Watch Video Solution**

**105.** If  $\sec(x+y) = xy$ , then find  $\frac{dy}{dx}$



**Watch Video Solution**

**106.** If  $y = \sqrt{\sin x + \sqrt{\sin x + \sqrt{\sin x + \dots}}}$ , prove that  $\frac{dy}{dx} = \frac{\cos x}{2y - 1}$



**Watch Video Solution**

**107.** If  $\log(x^2 + y^2) = 2\tan^{-1}\left(\frac{y}{x}\right)$ , show that  $\frac{dy}{dx} = \frac{x+y}{x-y}$



**Watch Video Solution**

**108.** If  $y = y(x)$  and it follows the relation  $4xe^{xy} = y + 5\sin^2x$ , then  $y'(0)$  is equal to \_\_\_\_\_

 Watch Video Solution

**109.** Find  $\frac{dy}{dx}$  if  $x = a(\theta - \sin\theta)$  and  $y = a(1 - \cos\theta)$ .

 Watch Video Solution

**110.** If  $x = a\sec^3\theta$  and  $y = a\tan^3\theta$ ,  $f \in d\frac{dy}{dx} \text{ at } \eta = \frac{\pi}{3}$

 Watch Video Solution

**111.** If  $x = e^{\cos 2t}$  and  $y = e^{\sin 2t}$ , then move that  $\frac{dy}{dx} = -\frac{y \log x}{x \log y}$ .

 Watch Video Solution

**112.** Let  $y = x^3 - 8x + 7$  and  $x = f(t)$ . If  $\frac{dy}{dt} = 2$  and  $x = 3$  at  $t = 0$ , then find the value of  $\frac{dx}{dt}$  at  $t = 0$ .

 Watch Video Solution

**113.** Find the derivative of  $\frac{\sqrt{x}(x+4)^{\frac{3}{2}}}{(4x-3)^{\frac{4}{3}}}$

 Watch Video Solution

**114.** If  $x^m y^n = (x+y)^{m+n}$ , prove that  $\frac{dy}{dx} = \frac{y}{x}$ .

 Watch Video Solution

**115.** Differentiate  $(\log x)^{\cos x}$  with respect to  $x$ .

 Watch Video Solution

**116.** If  $f(x) = |x|^{\sin x}$ , then find  $f' \left( -\frac{\pi}{4} \right)$



**Watch Video Solution**

**117.** If  $y = x^x \wedge x \in ((\infty))$ , find  $\frac{dy}{dx}$ .



**Watch Video Solution**

**118.** If  $f(x) = \lim_{h \rightarrow 0} \frac{(\sin(x+h))^{\log_e(x+h)} - (\sin x)^{\log_e x}}{h}$  then find  $f(\pi/2)$ .



**Watch Video Solution**

**119.** If  $0 < x < 1$ , prove that  $\frac{1}{1+x} + \frac{2x}{1+x^2} + \frac{4x^3}{1+x^4} + \dots^\infty = \frac{1}{1-x}$



**Watch Video Solution**

120. Differentiate  $\log \sin x$  w.r.t.  $\sqrt{x}$ .



[Watch Video Solution](#)

121. Find the derivative of  $f(\tan x) \text{ wrt } g(\sec x)$  at  $x = \frac{\pi}{4}$ , where  $f'(1) = 2$  and  $g'(\sqrt{2}) = 4$ .



[Watch Video Solution](#)

122. Let  $= \begin{vmatrix} \cos x & \sin x & \cos x \\ \cos 2x & \sin 2x & 2\cos 2x \\ \cos 3x & \sin 3x & 3\cos 3x \end{vmatrix}$  then find the values of  $f(0)$  and  $f'(\pi/2)$ .



[Watch Video Solution](#)

$$123. f(x) = \begin{vmatrix} \cos x & x & 1 \\ 2\sin x & x^2 & 2x \\ \tan x & x & 1 \end{vmatrix} \text{ then find the value of}$$

$$\lim_{x \rightarrow 0} \frac{f(x)}{x}$$



[Watch Video Solution](#)

$$124. \text{ If } y = \cos^{-1} x, \text{ find } \frac{d^2y}{dx^2} \text{ in terms of } y \text{ alone.}$$



[Watch Video Solution](#)

$$125. \text{ if } y = (x^2 - 1)^m, \text{ then the } (2m)\text{th differential coefficient of } y \text{ is}$$



[Watch Video Solution](#)

$$126. \text{ If } y = x \log \left\{ \frac{x}{(a + bx)} \right\}, \text{ then show that } x^3 \frac{d^2y}{dx^2} = \left( x \frac{dy}{dx} - y \right)^2.$$



[Watch Video Solution](#)

$$\left[ 1 + \left( \frac{dy}{dx} \right)^2 \right]^{\frac{3}{2}}$$

127. If  $(x - a)^2 + (y - b)^2 = c^2$ , for some  $c > 0$ , prove that  $\frac{\frac{d^2y}{dx^2}}{\left( 1 + \left( \frac{dy}{dx} \right)^2 \right)^{\frac{3}{2}}}$  is a

constant independent of a and b.



**Watch Video Solution**

128. If  $y = e^{a\cos^{-1}x}$ ,  $-1 \leq x \leq 1$ , show that  $\left(1 - x^2\right) \frac{d^2y}{dx^2} - x \frac{dy}{dx} - a^2y = 0$ .



**Watch Video Solution**

129. If  $x=a(\cos t + t \sin t)$  and  $y=a(\sin t - t \cos t)$ , find  $\frac{d^2y}{dx^2}$



**Watch Video Solution**

130. If g is inverse of f then prove that  $f'(g(x)) = -g''(x)(f(g(x)))^3$ .



Watch Video Solution

131. Let  $f(x)$  and  $g(x)$  be real valued functions such that  $f(x)g(x)=1$ ,  $\forall x \in R$ . If  $f'(x)$  and  $g'(x)$  exists  $\forall x \in R$  and  $f'(x)$  and  $g'(x)$  are never zero, then prove that  $\frac{f'(x)}{f(x)} - \frac{g''(x)}{g'(x)} = \frac{2f'(x)}{f(x)}$



Watch Video Solution

132. if  $f(x)$   $g(x)$  and  $h(x)$  are three polynomials of degree 2,

then prove that  $\phi(x) = \begin{vmatrix} f(x) & g(x) & h(x) \\ f'(x) & g'(x) & h'(x) \\ f''(x) & g''(x) & h''(x) \end{vmatrix}$  is a

constant polynomial.



Watch Video Solution

133. Let  $f\left(\frac{x+y}{2}\right) = \frac{f(x) + f(y)}{2}$  for all real  $x$  and  $y$ . If  $f'(0)$  exists and equals  $-1$  and  $f(0) = 1$ , then  $f \in df(2)$

 Watch Video Solution

134.  $f(x) + f\left(\frac{x+y}{1-xy}\right) \quad \text{for} \quad \forall x, y \in R$

( $xy \neq 1$ ), and ( $\lim_{x \rightarrow 0} \frac{f(x)}{x} = 2$ ).  $F \in df\left(\frac{1}{\sqrt{3}}\right)$  and  $f'(1)$

 Watch Video Solution

135. Find function  $f(x)$  which is differentiable and satisfy the relation

$$f(x+y) = f(x) + f(y) + (e^x - 1)(e^y - 1) \quad \forall x, y \in R, \text{ and } f(0) = 2.$$

 Watch Video Solution

136. If  $f\left(\frac{x}{y}\right) = \frac{f(x)}{f(y)}$ ,  $\forall y, f(y) \neq 0$  and  $f'(1) = 2$ , find  $f(x)$ .



[Watch Video Solution](#)

137. Let  $f: R \rightarrow R$  be a function satisfying condition  
 $f(x + y^3) = f(x) + [f(y)]^3$  for all  $x, y \in R$ . If  $f'(0) \geq 0$ , find  $f(10)$



[Watch Video Solution](#)

138. Let  $f(x + y) = f(x) + f(y) + 2xy - 1$  for all real  $x$  and  $y$  and  $f(x)$  be a differentiable function. If  $f'(0) = \cos\alpha$ , prove that  $f(x) > 0 \forall x \in R$



[Watch Video Solution](#)

**Solved Examples**

1. If  $f(x) = (\log)_x^2(\log x)$ , then  $f'(x)$  at  $x = e$  is 0 (b) 1 (c)  $\frac{1}{e}$  (d)  $\frac{1}{2}e$



Watch Video Solution

2. Given that  $\cos\left(\frac{x}{2}\right) \cdot \cos\left(\frac{x}{4}\right) \cdot \cos\left(\frac{x}{8}\right) \dots = \frac{\sin x}{x}$  Prove that  
 $\left(\frac{1}{2^2}\right)\sec^2\left(\frac{x}{2}\right) + \left(\frac{1}{2^4}\right)\sec^2\left(\frac{x}{4}\right) + \dots = \operatorname{cosec}^2 x - \frac{1}{x^2}$



Watch Video Solution

3. If  $y = f(a^x)$  and  $f'(\sin x) = (\log)_e x$ , then  $f \in d \frac{dy}{dx}$ , if it exists, where  $\pi/2$



Watch Video Solution

4. If  $P_n$  is the sum of a GP upto  $n$  terms ( $n \geq 3$ ), then prove that

$(1 - r) \frac{dP_n}{dr} = (1 - n)P_n + nP_{n-1}$ , where  $r$  is the common ratio of GP



Watch Video Solution



Watch Video Solution

5. If  $g(x) = \frac{f(x)}{(x - a)(x - b)(x - c)}$ , where  $f(x)$  is a polynomial of degree  $< 3$ ,

then

$$\int g(x)dx = |1af(a)\log|x - a|1bf(b)\log|x - b|1cf(c)\log|x - c|| \div |1aa^21 \wedge 21 \wedge 2| + k$$

$$\frac{dg(x)}{dx} = \left| 1af(a)\log(x - a)^21bf(b)\log(x - b)^21cf(c)\log(x - c)^2 \right| \div |a^2a1b^2b1c^2c1|$$

$$\frac{dg(x)}{dx} = \left| 1af(a)\log(x - a)^{-2}1bf(b)\log(x - b)^{-2}1cf(c)\log(x - c)^{-2} \right| \div |1aa^21 \wedge 21 \wedge 2|$$

$$\int g(x)dx = |1af(a)\log|x - a|1bf(b)\log|x - b|1cf(c)\log|x - c|| \div |a^2a1b^2b1c^2c1| + k$$



Watch Video Solution

6. If  $x = \operatorname{cosec}\theta - \sin\theta$  and  $y = \operatorname{cosec}^n\theta - \sin^n\theta$ , then show that

$$(x^2 + 4) \left( \frac{dy}{dx} \right)^2 = n^2(y^2 + 4).$$



Watch Video Solution

7. If  $y = \frac{ax^2}{(x-a)(x-b)(x-c)} + \frac{bx}{(x-b)(x-c)} + \frac{c}{x-c} + 1$ , then prove that

$$\frac{y'}{y} = \frac{1}{x} \left[ \frac{a}{a-x} + \frac{b}{b-x} + \frac{c}{c-x} \right]$$

 Watch Video Solution

8. Find the differential equation of the family of curves  $y = Ae^{2x} + Be^{-2x}$ , where A and B are arbitrary constants.

 Watch Video Solution

9. If  $y = \left(\frac{1}{2}\right)^{n-1} \cos(n\cos^{-1}x)$ , then prove that y satisfies the differential equation  $(1-x^2) \frac{d^2y}{dx^2} - x \frac{dy}{dx} + n^2y = 0$

 Watch Video Solution

10. Let  $f(x)$  and  $g(x)$  be two functions having finite nonzero third-order derivatives  $f'''$  and  $g'''$  for all  $x \in R$ . If  $f(x)g(x) = 1$  for all  $x \in R$ , then prove that  $f'''(x)f' - g'''(x)g' = 3\left(f''/f - g''/g\right)$



**Watch Video Solution**

11. If a curve is represented parametrically by the equation  $x = f(t)$  and  $y = g(t)$  then prove that  $\frac{d^2y}{dx^2} = - \left[ \frac{g'(t)}{f'(t)} \right]^3 \left( \frac{d^2x}{dy^2} \right)$



**Watch Video Solution**

12. If  $f\left(\frac{x+y}{3}\right) = \frac{2 + f(x) + f(y)}{3}$  for all real  $x$  and  $y$  and  $f'(2) = 2$ , then determine  $y = f(x)$



**Watch Video Solution**

13. If  $f(xy) = \frac{f(x)}{y} + \frac{f(y)}{x}$  holds for all real  $x$  and  $y$  greater than 0 and  $f(x)$  is a differentiable function for all  $x > 0$  such that  $f(e) = \frac{1}{e}$ , then find  $f(x)$

 Watch Video Solution

14. If  $|a_1 \sin x + a_2 \sin 2x + \dots + a_n \sin nx| \leq |\sin x|$  for  $x \in R$ , then prove that  $|a_1 + 2a_2 + 3a_3 + na_n| \leq 1$

 Watch Video Solution

15. Suppose  $p(x) = a_0 + a_1x + a_2x^2 + \dots + a_nx^n$ . If  $|p(x)| \leq e^{x-1} - 1$  for all  $x \geq 0$ , prove that  $|a_1 + 2a_2 + \dots + na_n| \leq 1$ .

 Watch Video Solution

16. If  $f(x) = \log_x(\log x)$ , then find  $f'(x)$  at  $x = e$

 Watch Video Solution

17. Given that  $\frac{\cos x}{2} \cdot \frac{\cos x}{4} \cdot \frac{\cos x}{8} \dots = \frac{\sin x}{x}$  Then find the sum  
 $\frac{1}{2^2} \frac{\sec^2 x}{2} + \frac{1}{2^4} \frac{\sec^2 x}{4} + \dots$



[Watch Video Solution](#)

18. If  $y = f(a^x)$  and  $f'(\sin x) = (\log)_e x$ , then  $f \in d\frac{dy}{dx}$ , if it exists, where  $\pi/2 < x < \pi$



[Watch Video Solution](#)

19. If  $P_n$  is the sum of a GP upto  $n$  terms ( $n \geq 3$ ), then prove that

$$(1 - r) \frac{dP_n}{dr} = (1 - n)P_n + nP_{n-1}, \text{ where } r \text{ is the common ratio of GP}$$



[Watch Video Solution](#)

20. If  $x = \operatorname{cosec}\theta - \sin\theta$  and  $y = \operatorname{cosec}^n\theta - \sin^n\theta$ , then show that

$$(x^2 + 4) \left( \frac{dy}{dx} \right)^2 = n^2 (y^2 + 4).$$



[Watch Video Solution](#)

21. If  $y = \frac{ax^2}{(x-a)(x-b)(x-c)} + \frac{bx}{(x-b)(x-c)} + \frac{c}{x-c} + 1$ , then prove that

$$\frac{y'}{y} = \frac{1}{x} \left[ \frac{a}{a-x} + \frac{b}{b-x} + \frac{c}{c-x} \right]$$



[Watch Video Solution](#)

22.

If  $y = \frac{2}{\sqrt{a^2 - b^2}} \left\{ \tan^{-1} \left( \sqrt{\frac{a-b}{a+b}} \tan \frac{x}{2} \right) \right\}$ , then show that  $\frac{d^2y}{dx^2} = \frac{bs \sin x}{(a + b \cos x)^2}$ .



[Watch Video Solution](#)

23. If  $y = \left(\frac{1}{2}\right)^{n-1} \cos(n \cos^{-1} x)$ , then prove that  $y$  satisfies the differential equation  $(1 - x^2) \frac{d^2y}{dx^2} - x \frac{dy}{dx} + n^2 y = 0$



Watch Video Solution

24. If a curve is represented parametrically by the equation

$x = f(t)$  and  $y = g(t)$  then prove that  $\frac{d^2y}{dx^2} = - \left[ \frac{g'(t)}{f'(t)} \right]^3 \left( \frac{d^2x}{dy^2} \right)$



Watch Video Solution

25. If  $f\left(\frac{x+y}{3}\right) = \frac{2 + f(x) + f(y)}{3}$  for all real  $x$  and  $y$  and  $f'(2) = 2$ , then determine  $y = f(x)$



Watch Video Solution

26. If  $f(x) = \frac{f(x)}{y} + \frac{f(y)}{x}$  holds for all real  $x$  and  $y$  greater than 0 and  $f(x)$  is a differentiable function for all  $x > 0$  such that  $f(e) = \frac{1}{e}$ , then  $f \in df(x)$ .

 Watch Video Solution

27. If  $|a_1 \sin x + a_2 \sin 2x + \dots + a_n \sin nx| \leq |\sin x|$  for  $x \in R$ , then prove that  $|a_1 + 2a_2 + 3a_3 + na_n| \leq 1$

 Watch Video Solution

28. Suppose  $p(x) = a_0 + a_1x + a_2x^2 + \dots + a_nx^n$ . If  $|p(x)| \leq e^{x-1} - 1$  for all  $x \geq 0$ , prove that  $|a_1 + 2a_2 + \dots + na_n| \leq 1$ .

 Watch Video Solution

Concept Application 3 1

1. Using the definition of derivative find the derivative of  $\sqrt{\sin x}$



Watch Video Solution

### Concept Application 3 2

1. Find the derivative of  $\sqrt{4 - x}$  w.r.t.  $x$  using the first principle.



Watch Video Solution

2. If  $y = \sin^{-1} \left( \frac{2x}{1 + x^2} \right)$ , then find  $\frac{dy}{dx}$



Watch Video Solution

3.  $y = \tan^{-1} \frac{3x - x^3}{2x^2 - 1}$ ,  $-\frac{1}{\sqrt{3}} < x < \frac{1}{\sqrt{3}}$



Watch Video Solution

4. If  $y = \sec^{-1}(1/(2x^2 - 1))$ ; 0



**Watch Video Solution**

5. Find  $\frac{dy}{dx}$  if  $y = \frac{\tan^{-1}(4x)}{1 + 5x^2} + \frac{\tan^{-1}(2 + 3x)}{3 - 2x}$



**Watch Video Solution**

6. Find  $\frac{dy}{dx}$  if  $y = \tan^{-1}\left(\frac{\sqrt{1+x^2} - 1}{x}\right)$ , where  $x \neq 0$



**Watch Video Solution**

7.  $y = \tan^{-1}\left(\frac{x}{1 + \sqrt{1-x^2}}\right)$



**Watch Video Solution**

8. Find  $\frac{dy}{dx}$  for the function:  $y = \sin^{-1}\sqrt{(1-x)} + \cos^{-1}\sqrt{x}$



**Watch Video Solution**

9.  $y = \sqrt{\sin\sqrt{x}}$



**Watch Video Solution**

10.  $y = e^{\sin x^3}$  find  $\frac{dy}{dx}$



**Watch Video Solution**

11.  $y = \log\sqrt{\sin\sqrt{e^x}}$



**Watch Video Solution**

12. Find  $\frac{dy}{dx}$  for the function:  $y = a^{\sin^{-1}x} \wedge (2)$



Watch Video Solution

13. Find  $\frac{dy}{dx}$  if  $y = \log \left\{ e^x \left( \frac{x-2}{x+2} \right)^{\frac{3}{4}} \right\}$



Watch Video Solution

14.  $y = \sin^{-1} \left[ \sqrt{x - ax} - \sqrt{a - ax} \right]$



Watch Video Solution

15. Find  $\frac{dy}{dx}$  for the functions:  $y = x^3 e^x \sin x$



Watch Video Solution

16. Find  $\frac{dy}{dx}$  for the function:  $y = (\log)_e \sqrt{\frac{1 + \sin x}{1 - \sin x}}$ , where  $x = \frac{\pi}{3}$



Watch Video Solution

17. Find  $\frac{dy}{dx}$  for the functions:  $y = \frac{x + \sin x}{x + \cos x}$



Watch Video Solution

18. If  $y = (1 + x)(1 + x^2)(1 + x^4) \dots (1 + x^{2^n})$  then  $\frac{dy}{dx}$  at  $x = 0$  is



Watch Video Solution

19.  $x\sqrt{1+y} + y\sqrt{1+x} = 0$  then  $\frac{dy}{dx} =$



Watch Video Solution

**20.** If  $g$  is the inverse function of and  $f'(x) = \sin x$  then prove that  $g'(x) = \operatorname{cosec}(g(x))$

 **Watch Video Solution**

**21.** Find the derivative of  $\sqrt{4 - x}$  w.r.t.  $x$  using the first principle.

 **Watch Video Solution**

**22.** Find  $\frac{dy}{dx}$ ,  $y = \sin^{-1} \frac{2x}{1+x^2}$ ,  $-1 \leq x \leq 1$

 **Watch Video Solution**

**23.** Find  $\frac{dy}{dx}$ ,  $y = \tan^{-1} \left[ \frac{3x - x^3}{1 - 3x^2} \right]$ ,  $-\frac{1}{\sqrt{3}} < x < \frac{1}{\sqrt{3}}$

 **Watch Video Solution**

$$24. y = \sec^{-1} \frac{1}{2x^2 - 1}, 0 < x < \frac{1}{\sqrt{2}}$$



Watch Video Solution

$$25. \text{Find } \frac{dy}{dx} \text{ if } y = \frac{\tan^{-1}(4x)}{1 + 5x^2} + \frac{\tan^{-1}(2 + 3x)}{3 - 2x}$$



Watch Video Solution

$$26. \text{Find } \frac{dy}{dx} \text{ if } y = \tan^{-1} \left( \frac{\sqrt{1+x^2} - 1}{x} \right), \text{ where } x \neq 0$$



Watch Video Solution

$$27. y = \tan^{-1} \left( \frac{x}{1 + \sqrt{1-x^2}} \right)$$



Watch Video Solution

28. Find  $\frac{dy}{dx}$  for the function:  $y = \sin^{-1}\sqrt{(1-x)} + \cos^{-1}\sqrt{x}$



Watch Video Solution

29.  $y = \sqrt{\sin\sqrt{x}}$



Watch Video Solution

30.  $y = e^{\sin x^3}$  find  $\frac{dy}{dx}$



Watch Video Solution

31.  $y = \log\sqrt{\sin\sqrt{e^x}}$



Watch Video Solution

32. Find  $\frac{dy}{dx}$  for the function:  $y = a(\sin^{-1}x)^2$



Watch Video Solution

33. Find  $\frac{dy}{dx}$  if  $y = \log \left\{ e^x \left( \frac{x-2}{x+2} \right)^{\frac{3}{4}} \right\}$



Watch Video Solution

34.  $y = \sin^{-1} [\sqrt{x - ax} - \sqrt{a - ax}]$  provethat  $\frac{dy}{dx}$  is  $\frac{1}{2\sqrt{x(1-x)}}$



Watch Video Solution

35. Find  $\frac{dy}{dx}$  for the functions:  $y = x^3 e^x \sin x$



Watch Video Solution

36. Find  $\frac{dy}{dx}$  for the function:  $y = (\log_e) \sqrt{\frac{1 + \sin x}{1 - \sin x}}$ , where  $x = \frac{\pi}{3}$



Watch Video Solution

37.  $y = \frac{x + \sin x}{x + \cos x}$ , find  $\frac{dy}{dx}$



Watch Video Solution

38. If  $y = (1 + x)(1 + x^2)(1 + x^4) \dots (1 + x^{2^n})$  then  $\frac{dy}{dx}$  at  $x = 0$  is



Watch Video Solution

39. If  $x\sqrt{1+y} + y\sqrt{1+x} = 0$ , prove that  $\frac{dy}{dx} = -\frac{1}{(x+1)^2}$ .



Watch Video Solution

40. If  $g$  is the inverse function of  $f(x) = \sin x$  then prove that  $g'(x) = \operatorname{cosec}(g(x))$



Watch Video Solution

## Concept Application 3 3

1. Let  $f(x) = \frac{(2^x + 2^{-x}) \sin x \sqrt{\tan^{-1}(x^2 - x + 1)}}{(7x^2 + 3x + 1)^3}$ . Then find the value of  $f'(0)$ .



Watch Video Solution

2. If  $x^3 + y^3 = 3axy$ , find  $\frac{dy}{dx}$



Watch Video Solution

3. If  $y = b \tan^{-1}(x/a) + \tan^{-1}(y/x)$ , find  $(dy)/(dx)$



Watch Video Solution

4. If  $\log_e(\log_e x - \log_e y) = e^{x^2y}(1 - \log_e x)$ , then find the value of  $y'(e)$ .



**Watch Video Solution**

5. If  $y = \sqrt{x + \sqrt{y + \sqrt{x + \sqrt{y + \dots}}}}$ , then prove that  $\frac{dy}{dx} = \frac{y^2 - x}{2y^3 - 2xy - 1}$



**Watch Video Solution**

6. Let  $f(x) = \frac{(2^x + 2^{-x})\sin x \sqrt{\tan^{-1}(x^2 - x + 1)}}{(7x^2 + 3x + 1)^3}$ . Then find the value of  $f'(0)$ .



**Watch Video Solution**

7. If  $x^3 + y^3 = 3axy$ , find  $\frac{dy}{dx}$ .



**Watch Video Solution**

8. If  $y = b \tan^{-1} \left( \frac{x}{a} + \tan^{-1} y/x \right)$ , find  $\frac{dy}{dx}$ .



**Watch Video Solution**

9. If  $\log_e (\log_e x - \log_e y) = e^{x^2y} (1 - \log_e x)$ , then find the value of  $y'(e)$ .



**Watch Video Solution**

10. If  $y = \sqrt{x + \sqrt{y + \sqrt{x + \sqrt{y + \dots}}}}$  then prove that  $\frac{dy}{dx} = \frac{y^2 - x}{2y^3 - 2xy - 1}$



**Watch Video Solution**

### Concept Application 3 4

1. Statement 1: Let  $f: \mathbb{R} \rightarrow \mathbb{R}$  be a real-valued function  $\forall x, y \in \mathbb{R}$  such that  $|f(x) - f(y)| \leq |x - y|^3$ . Then  $f(x)$  is a constant function. Statement 2: If the derivative of the function w.r.t.  $x$  is zero, then function is constant.



Watch Video Solution

2. If  $x = \frac{2t}{1+t^2}$ ,  $y = \frac{1-t^2}{1+t^2}$ , then if  $\in d\frac{dy}{dx} a = 2$ .



Watch Video Solution

3. If  $x = \sqrt{a^{\sin^{-1}t}}$ ,  $y = \sqrt{a^{\cos^{-1}t}}$  then show that,  $\frac{dy}{dx} = -\frac{y}{x}$ .



Watch Video Solution

4. Find  $\frac{dy}{dx}$  if  $x = 3 \cos \theta - \cos 2\theta$  and  $y = \sin \theta - \sin 2\theta$ .



Watch Video Solution

5. If  $x = 3\cos\theta - 2\cos^3\theta$ ,  $y = 3\sin\theta - 2\sin^3\theta$ , then  $\frac{dy}{dx}$  is



Watch Video Solution

6. If  $x = a \left( \cos t + \frac{1}{2} \log \tan' \frac{t}{2} \right)$  and  $y = \sin t$  then  $f \in d(dy)/(dx)$  at  $t = \pi/4$



**Watch Video Solution**

7. Statement 1: Let  $f: \vec{R} \rightarrow \vec{R}$  be a real-valued function  $\forall x, y \in R$  such that  $|f(x) - f(y)| \leq |x - y|^3$ . Then  $f(x)$  is a constant function. Statement 2: If the derivative of the function w.r.t.  $x$  is zero, then function is constant.



**Watch Video Solution**

8. If  $x = \frac{2t}{1+t^2}, y = \frac{1-t^2}{1+t^2}$ , then find  $\frac{dy}{dx}$  at  $t = 2$ .



**Watch Video Solution**

9. If  $x = \sqrt{a^{\sin(-1)t}}, y = \sqrt{a^{\cos(( - 1)t)}}$ , show that  $\frac{dy}{dx} = -\frac{y}{x}$



**Watch Video Solution**

10. Find  $\frac{dy}{dx}$  if  $x = \cos\theta - \cos 2\theta$

and  $y = \sin\theta - \sin 2\theta$



Watch Video Solution

11. Find  $\frac{dy}{dx}$  if  $x = 3\cos\theta - 2\cos^3\theta, y = 3\sin\theta - 2\sin^3\theta.$



Watch Video Solution

12. about to only mathematics



Watch Video Solution

Concept Application 3 5

1. Differentiate  $\sqrt{\frac{(x - 1)(x - 2)}{(x - 3)(x - 4)(x - 5)}}$  with respect to  $x$



Watch Video Solution

2. If  $x^y = e^{x-y}$ , prove that  $\frac{dy}{dx} = \frac{\log x}{(1 + \log x)^2}$



Watch Video Solution

3. If  $xy = e^{(x-y)}$ , then find  $\frac{dy}{dx}$



Watch Video Solution

4. If  $y^x = x^y$ , then find  $\frac{dy}{dx}$



Watch Video Solution

5. If  $x = e^y + e^{(y+ \rightarrow \infty)}$ , where  $x > 0$ , then  $\frac{dy}{dx}$



Watch Video Solution

6. Find  $\frac{dy}{dx}$  if  $y = x^x$



Watch Video Solution

7. Differentiate  $(x \cos x)^x$  with respect to  $x$



Watch Video Solution

8. If  $y = (\tan x)^{(\tan x)^{\tan x}}$ , then find  $\frac{dy}{dx}$ .



Watch Video Solution

9. Differentiate  $\sqrt{\frac{(x-1)(x-2)}{(x-3)(x-4)(x-5)}}$  with respect to  $x$



Watch Video Solution

10. If  $y \log x = x - y$ , prove that  $\frac{dy}{dx} = \frac{\log x}{(1 + \log x)^2}$



Watch Video Solution

11. If  $xy = e^{(x-y)}$ , then find  $\frac{dy}{dx}$ .



Watch Video Solution

12. If  $y^x = x^y$ , then find  $\frac{dy}{dx}$ .



Watch Video Solution

13. If  $x = e^y + e^{(y+ \rightarrow \infty)}$ , where  $x > 0$ , then find  $\frac{dy}{dx}$



Watch Video Solution

14. Find  $\frac{dy}{dx}$  or  $y = x^x$



Watch Video Solution

15. Differentiate  $(x \cos x)^x$  with respect to  $x$



Watch Video Solution

16. If  $y = (\tan x)^{(\tan x)^{\tan x}}$ , then find  $\frac{dy}{dx}$ .



Watch Video Solution

1. Find the derivative of  $\frac{\tan^{-1}(2x)}{1-x^2} \cdot \frac{\sin^{-1}(2x)}{1+x^2}$



Watch Video Solution

2. The differential coefficient of  $\sec^{-1}\left(\frac{1}{2x^2 - 1}\right)$  w.r.t  $\sqrt{1 - x^2}$  is-



Watch Video Solution

3. Differentiate  $\frac{x}{\sin x}$  w.r.t .  $\sin x$ .



Watch Video Solution

4.

If  $y = f(x^3)$ ,  $z = g(x^5)$ ,  $f'(x) = \tan x$ , and  $g'(x) = \sec x$ , then find the value of of



Watch Video Solution

5. Find the derivative of  $\tan^{-1} \frac{2x}{1-x^2}$  w.r.t.  $\sin^{-1} \frac{2x}{1+x^2}$ ,  $|x| < 1$ .



Watch Video Solution

6. Find the derivative of  $\sec^{-1} \left( \frac{1}{2x^2 - 1} \right)$  w.r.t.  $\sqrt{1 - x^2}$  at  $x = \frac{1}{2}$ .



Watch Video Solution

7. Differentiate  $\frac{x}{\sin x}$  w.r.t .  $\sin x$ .



Watch Video Solution

8.

If  $y = f(x^3)$ ,  $z = g(x^5)$ ,  $f'(x) = \tan x$ , and  $g'(x) = \sec x$ , then find the value of of



Watch Video Solution

## Concept Application 3 7

1. If  $f(x) = \begin{vmatrix} x + a^2 & ab & ac \\ ab & x + b^2 & bc \\ ac & bc & x + c^2 \end{vmatrix}$ , then prove that

$$f'(x) = 3x^2 + 2x(a^2 + b^2 + c^2).$$



Watch Video Solution

2. Let  $f(x) = \begin{vmatrix} \cos(x + x^2) & \sin(x + x^2) & -\cos(x + x^2) \\ \sin(x - x^2) & \cos(x - x^2) & \sin(x - x^2) \\ \sin 2x & 0 & \sin(2x^2) \end{vmatrix}$ .

Find the value of  $f'(0)$ .



Watch Video Solution

$$3. \text{ Let } g(x) = \begin{vmatrix} f(x+c) & f(x+2c) & f(x+3c) \\ f(c) & f(2c) & f(3c) \\ f(c) & f(2c) & f(3c) \end{vmatrix},$$

where c is constant, then find  $\lim_{x \rightarrow 0} \frac{g(x)}{x}$ .



**Watch Video Solution**

$$4. \text{ If } f(x) = \begin{vmatrix} x+a^2 & ab & ac \\ ab & x+b^2 & bc \\ ac & bc & x+c^2 \end{vmatrix}, \text{ then prove that}$$

$$f'(x) = 3x^2 + 2x(a^2 + b^2 + c^2).$$



**Watch Video Solution**

$$5. \text{ Let } f(x) = \begin{vmatrix} \cos(x+x^2) & \sin(x+x^2) & -\cos(x+x^2) \\ \sin(x-x^2) & \cos(x-x^2) & \sin(x-x^2) \\ \sin 2x & 0 & \sin(2x^2) \end{vmatrix}.$$

Find the value of  $f'(0)$ .



Watch Video Solution

6. Let  $g(x) = \begin{vmatrix} f(x+c) & f(x+2c) & f(x+3c) \\ f(c) & f(2c) & f(3c) \\ f'(c) & f'(2c) & f'(3c) \end{vmatrix}$ ,

where c is constant, then find  $\lim_{x \rightarrow 0} \frac{g(x)}{x}$ .



Watch Video Solution

### Concept Application 3 8

1. If  $f(x) = (1+x)^2$ , then the value of  $f(x_0) + f'(0) + \frac{f^0}{2!} + \frac{f^0}{3!} + \frac{f^n(0)}{n!} \cdot$



Watch Video Solution

2. If  $e^y(x+1) = 1$ , show that  $\frac{d^2y}{dx^2} = \left(\frac{dy}{dx}\right)^2$



Watch Video Solution

3. Prove that  $\frac{d^n}{dx^n} \left( e^{2x} + e^{-2x} \right) = 2^n \left[ e^{2x} + (-1)^n e^{-2x} \right]$



Watch Video Solution

4. If  $y = \sin(\sin x)$  and  $\frac{d^2y}{dx^2} + \frac{dy}{dx} \tan x + f(x) = 0$ , then find  $f(x)$ .



Watch Video Solution

5. If  $y = \log(1 + \sin x)$ , prove that  $y_4 + y_3 y_1 + y_2^2 = 0$ .



Watch Video Solution

6. If  $f(x) = \begin{vmatrix} x^n & n! & 2; \cos x & \cos\left(\frac{n\pi}{2}\right) & 4; \sin x & \sin\left(\frac{n\pi}{2}\right) & 8 \end{vmatrix}$  then find the value of  $\frac{d^n}{dx^n} ([f(x)])_{x=0} \quad n \in \mathbb{Z}$



Watch Video Solution

7. If  $x = a\cos\theta$ ,  $y = b\sin\theta$ , then prove that  $\frac{d^3y}{dx^3} = \frac{3b}{a^3}\cosec^4\theta\cot\theta$ .



[Watch Video Solution](#)

8. If  $x = a\cos^3\theta$ ,  $y = b\sin^3\theta$ ,  $f \in d\frac{d^3y}{dx^3}$  at  $\theta = 0$ .



[Watch Video Solution](#)

9. If  $f(x) = (1 + x)^2$ , then the value of  $f(x_0) + f'(0) + \frac{f^0}{2!} + \frac{f^0}{3!} + \frac{f^n(0)}{n!}$



[Watch Video Solution](#)

10. If  $e^y(x + 1) = 1$ , show that  $\frac{d^2y}{dx^2} = \left(\frac{dy}{dx}\right)^2$



[Watch Video Solution](#)

11. Prove that  $\frac{d^n}{dx^n} \left( e^{2x} + e^{-2x} \right) = 2^n \left[ e^{2x} + (-1)^n e^{-2x} \right]$



Watch Video Solution

12. If  $y \sin(\sin x)$  and  $\frac{d^2y}{dx^2} + \frac{dy}{dx} \tan x + f(x) = 0$ , then find  $f(x)$ .



Watch Video Solution

13. If  $y = \log(1 + \sin x)$ , prove that  $y_4 + y_3 y_1 + y_2^2 = 0$ .



Watch Video Solution

14. If  $x = a\cos\theta$ ,  $y = b\sin\theta$ , then prove that  $\frac{d^3y}{dx^3} = \frac{3b}{a^3} \operatorname{cosec}^4\theta \cot\theta$ .



Watch Video Solution

15. If  $x = a\cos^3\theta$ ,  $y = b\sin^3\theta$ ,  $f \in d\frac{d^3y}{dx^3}$  at  $\theta = 0$ .



Watch Video Solution

### Concept Application 3 9

1. Let  $f(x + y) = f(x)f(y)$  for all  $x$  and  $y$ . Suppose  $f(5) = 2$  and  $f'(0) = 3$ . Find  $f'(5)$



Watch Video Solution

2. Let  $f(xy) = f(x)f(y) \forall x, y \in R$  and  $f$  is differentiable at  $x = 1$  such that  $f'(1) = 1$ . Also,  $f(1) \neq 0$ ,  $f(2) = 3$ . Then find  $f'(2)$



Watch Video Solution

3. Let  $f$  be a function such that  $f(x + y) = f(x) + f(y)$  for all  $x$  and  $y$  and  $f(x) = (2x^2 + 3x)g(x)$  for all  $x$ , where  $g$  is continuous and  $g(0) = 3$ . Then find  $f'(x)$



Watch Video Solution

4. Let  $g: R \rightarrow R$  be a differentiable function satisfying  $g(x) = g(y)g(x - y) \forall x, y \in R$  and  $g'(0) = a$  and  $g'(3) = b$ , Then find the value of



Watch Video Solution

5. Let  $f(x^m y^n) = mf(x) + nf(y)$  for all  $x, y \in R^+$  and for all  $m, n \in R$ . If  $f'(x)$  exists and has the value  $\frac{e}{x}$ , then find  $(\lim)_{x \rightarrow 0} \frac{f(1+x)}{x}$



Watch Video Solution

6. If  $f\left(\frac{x+2y}{3}\right) = \frac{f(x) + 2f(y)}{3} \forall x, y \in R$  and  $f'(0) = 1, f(0) = 2$ , then find  $f(x)$



Watch Video Solution

7.

Prove that  $\lim_{h \rightarrow 0} \frac{f(x + h) + f(x - h) - 2f(x)}{h^2} = f'(x)$  (without using L'Hospital's rule)



[Watch Video Solution](#)

8. Let  $f(x + y) = f(x)f(y)$  for all  $x$  and  $y$ . Suppose  $f(5) = 2$  and  $f'(0) = 3$ . Find  $f'(5)$



[Watch Video Solution](#)

9. Let  $f(xy) = f(x)f(y) \forall x, y \in R$  and  $f$  is differentiable at  $x = 1$  such that  $f'(1) = 1$ . Also,  $f(1) \neq 0$ ,  $f(2) = 3$ . Then find  $f'(2)$



[Watch Video Solution](#)

10. Let  $f$  be a function such that  $f(x + y) = f(x) + f(y)$  for all  $x$  and  $y$  and  $f(x) = (2x^2 + 3x)g(x)$  for all  $x$ , where  $g(x)$  is continuous and

$g(0) = 3$ . Then find  $f'(x)$



**Watch Video Solution**

11. Let  $g: \vec{R} \rightarrow \vec{R}$  be a differentiable function satisfying  $g(x) = g(y)g(x - y) \forall x, y \in R$  and  $g'(0) = a$  and  $g'(3) = b$ . Then find the value of  $g'(-3)$



**Watch Video Solution**

12. Let  $f(x^m y^n) = mf(x) + nf(y)$  for all  $x, y \in R^+$  and for all  $m, n \in R$ . If  $f'(x)$  exists and has the value  $\frac{e}{x}$ , then find  $(\lim)_{x \rightarrow 0} \frac{f(1+x)}{x}$



**Watch Video Solution**

13. If  $f\left(\frac{x+2y}{3}\right) = \frac{f(x) + 2f(y)}{3} \forall x, y \in R$  and  $f'(0) = 1, f(0) = 2$ , then find  $f(x)$



Watch Video Solution

14.

Prove

that

$$\lim_{h \rightarrow 0} \frac{f(x + h) + f(x - h) - 2f(x)}{h^2} = f'(x) \text{ (without using L'Hospital's rule).}$$



Watch Video Solution

## Exercises

1. If  $y = a\sin x + b\cos x$ , then  $\left(\frac{dy}{dx}\right)^2 + y^2$  is

- A. function of  $x$
- B. function of  $y$
- C. function of  $x$  and  $y$
- D. constant

**Answer:**



Watch Video Solution

2.  $\frac{d}{dx} \sqrt{\frac{1-\sin 2x}{1+\sin 2x}}$  is equal to,

A.  $\sec^2 x$

B.  $-\sec^2\left(\frac{\pi}{4} - x\right)$

C.  $\sec^2\left(\frac{\pi}{4} + x\right)$

D.  $\sec^2\left(\frac{\pi}{4} - x\right)$

**Answer:**



Watch Video Solution

3. If  $f(x) = |\cos x| + |\sin x|$ , then  $\frac{dy}{dx}$  at  $x = \frac{2\pi}{3}$  is equal to

A.  $\frac{1 - \sqrt{3}}{2}$

B. 0

C.  $\frac{1}{2}(\sqrt{3} - 1)$

D. none of these

**Answer:**



**Watch Video Solution**

4. If  $f(x) = |\log_e|x||$ , then  $f'(x)$  equals

A.  $\frac{1}{|x|}$ , where  $x \neq 0$

B.  $\frac{1}{x}$  for  $|x| > 1$  and  $-\frac{1}{x}$  for  $|x| < 1$

C.  $-\frac{1}{x}$  for  $|x| > 1$  and  $\frac{1}{x}$  for  $|x| < 1$

D.  $\frac{1}{x}$  for  $x > 0$  and  $-\frac{1}{x}$  for  $x < 0$

**Answer:**



**Watch Video Solution**

5. If  $f(x) = \sqrt{1 - \sin 2x}$ , then  $f'(x)$  is equal to

(a)  $-(\cos x + \sin x)$ , for  $x \in \left(\frac{\pi}{4}, \frac{\pi}{2}\right)$

(b)  $\cos x + \sin x$ , for  $x \in \left(0, \frac{\pi}{4}\right)$

(c)  $-(\cos x + \sin x)$ , for  $x \in \left(0, \frac{\pi}{4}\right)$

(d)  $\cos x - \sin x$ , for  $x \in \left(\frac{\pi}{4}, \frac{\pi}{2}\right)$

A.  $-(\cos x + \sin x)$ , for  $x \in (\pi/4, \pi/2)$

B.  $\cos x + \sin x$  for  $x \in (0, \pi/4)$

C.  $-(\cos x + \sin x)$ , for  $x \in (0, \pi/4)$

D.  $\cos x - \sin x$ , for  $x \in (\pi/4, \pi/2)$

**Answer:**



**Watch Video Solution**

6. Instead of the usual definition of derivative  $Df(x)$ , if we define a new

kind of derivative  $D^*F(x)$  by the formula

$$D^*(x) = (\lim_{h \rightarrow 0}) \frac{f^2(x+h) - f^2(x)}{h}, \text{ where } f^2(x)$$

$[f(x)]^2$  and  $f(x) = x \log x$ , then  $D^*f(x)(|)_{x=e}$  has the value e (B) 2e (c) 4e (d)

none of these

A. e

B. 2e

C. 4e

D. none of these

Answer:



Watch Video Solution

7. If  $y = \cot^{-1} \left[ \frac{\sqrt{1 + \sin x} + \sqrt{1 - \sin x}}{\sqrt{1 + \sin x} - \sqrt{1 - \sin x}} \right] (0 < x < \pi/2)$ , then  $\frac{dy}{dx} =$

A.  $\frac{1}{2}$

B.  $\frac{2}{3}$

C. 3

D. 1

**Answer:**



**Watch Video Solution**

8. if  $y = \frac{(a - x)\sqrt{a - x} - (b - x)\sqrt{x - b}}{\sqrt{a - x} + \sqrt{x - b}}$  then  $\frac{dy}{dx}$  wherever it is defined is equal

to :

A.  $\frac{x + (a + b)}{\sqrt{(a - x)(x - b)}}$

B.  $\frac{2x - a - b}{2\sqrt{a - x}\sqrt{x - b}}$

C.  $-\frac{(a + b)}{2\sqrt{(a - x)(x - b)}}$

D.  $\frac{2x + (a + b)}{2\sqrt{(a - x)(x - b)}}$

**Answer:**



**Watch Video Solution**

**9.** the derivative of  $y = (1 - x)(2 - x) \dots (n - x)$  at  $x = 1$  is equal to

A. 0

B.  $(-1)(n - 1)!$

C.  $n! - 1$

D.  $(-1)^{n-1}(n - 1)!$

**Answer:**



**Watch Video Solution**

**10.** If  $y = \sqrt{\frac{1-x}{1+x}}$ , then  $\left(1-x^2\right) \frac{dy}{dx}$  is equal to

A.  $y^2$

B.  $1/y$

C.  $-y$

D.  $-y/x$

**Answer:**



**Watch Video Solution**

11. If  $y = \frac{\sqrt{a+x} - \sqrt{a-x}}{\sqrt{a+x} + \sqrt{a-x}}$ , then  $\frac{dy}{dx}$  is equal to (a)  $\frac{ay}{x\sqrt{a^2 - x^2}}$  (b)  $\frac{ay}{\sqrt{a^2 - x^2}}$

(c)  $\frac{ay}{x\sqrt{a^2 - x^2}}$  (d) none of these

A.  $\frac{ay}{x\sqrt{a^2 - x^2}}$

B.  $\frac{ay}{\sqrt{a^2 - x^2}}$

C.  $\frac{ay}{x\sqrt{x^2 - a^2}}$

D. none of these

**Answer:**



Watch Video Solution

12. Let  $u(x)$  and  $v(x)$  be differentiable functions such that

$\frac{u(x)}{v(x)} = 7$ .  $\frac{u'(x)}{v'(x)} = p$  and  $\left(\frac{u(x)}{v(x)}\right)' = q$ , then  $\frac{p+q}{p-q}$  has the value of to 1 (b) 0

(c) 7 (d) -7

A. 1

B. 0

C. 7

D. -7

**Answer:**



Watch Video Solution

13. If  $\sin^{-1}\left(\frac{x^2 - y^2}{x^2 + y^2}\right) = \log a$ , then  $\frac{dy}{dx}$  is equal to  $\frac{x}{y}$  (b)  $\frac{y}{x^2}$   $\frac{x^2 - y^2}{x^2 + y^2}$  (d)  $\frac{y}{x}$

A.  $\frac{x}{y}$

B.  $\frac{y}{x^2}$

C.  $\frac{x^2 - y^2}{x^2 + y^2}$

D.  $\frac{y}{x}$

**Answer:**



**Watch Video Solution**

14. Let  $h(x)$  be differentiable for all  $x$  and let  $f(x) = (kx + e^x)h(x)$ , where  $k$  is some constant. If  $h(0) = 5$ ,  $h'(0) = -2$ , and  $f'(0) = 18$ , then the value of  $k$  is

A. 5

B. 4

C. 3

D. 2.2

**Answer:**



**Watch Video Solution**

**15.** If  $\lim_{t \rightarrow x} \frac{e^t f(x) - e^x f(t)}{(t - x)(f(x))^2} = 2$  and  $f(0) = \frac{1}{2}$ , then find the value of  $f'(0)$ .

A. 4

B. 2

C. 0

D. 1

**Answer:**



**Watch Video Solution**

**16.** If  $f(0) = 0$ ,  $f'(0) = 2$ , then the derivative of  $y = f(f(f(x)))$  at  $x = 0$  is 2 (b)

8 (c) 16 (d) 4

A. 2

B. 8

C. 16

D. 4

**Answer:**



**Watch Video Solution**

17. If  $f(x) = \sqrt{1 + \cos^2(x^2)}$ , then  $f' \left( \frac{\sqrt{\pi}}{2} \right)$  is  $\frac{\sqrt{\pi}}{6}$  (b)  $-\sqrt{\pi/6}$  (c)  $1/\sqrt{6}$  (d)  $\pi/\sqrt{6}$

A.  $\sqrt{\pi/6}$

B.  $-\sqrt{\pi/6}$

C.  $1/\sqrt{6}$

D.  $\pi/\sqrt{6}$

**Answer:**



[Watch Video Solution](#)



Watch Video Solution

18.  $\frac{d}{dx} \cos^{-1} \sqrt{\cos x}$  is equal to

A.  $\frac{1}{2} \sqrt{1 + \sec x}$

B.  $\sqrt{1 + \sec x}$

C.  $-\frac{1}{2} \sqrt{1 + \sec x}$

D.  $-\sqrt{1 + \sec x}$

**Answer:**



Watch Video Solution

19. if  $y = \tan^{-1} \left( \frac{2^x}{1 + 2^{2x+1}} \right)$  then  $\frac{dy}{dx} \text{at } x = 0$  is

A. 1

B. 2

C. In 2

D.  $-\frac{1}{10} \ln 2$

**Answer:**



**Watch Video Solution**

20. If  $y = \left( x + \sqrt{x^2 + a^2} \right)^n$ , then  $\frac{dy}{dx}$  is (b)  $-\frac{ny}{\sqrt{x^2 + a^2}}$  (d)  $\frac{ny}{\sqrt{x^2 + a^2}} \frac{nx}{\sqrt{x^2 + a^2}}$

-  $\frac{nx}{\sqrt{x^2 + a^2}}$

A.  $\frac{ny}{\sqrt{x^2 + a^2}}$

B.  $-\frac{ny}{\sqrt{x^2 + a^2}}$

C.  $\frac{nx}{\sqrt{x^2 + a^2}}$

D.  $-\frac{nx}{\sqrt{x^2 + a^2}}$

**Answer:**



**Watch Video Solution**

21. if  $y = \log_{\sin x} \tan x$  then  $\left(\frac{dy}{dx}\right)_{\frac{\pi}{4}}$  is

A.  $\frac{4}{\log 2}$

B.  $-4\log 2$

C.  $\frac{-4}{\log 2}$

D. none of these

**Answer:**



**Watch Video Solution**

22.  $\frac{d}{dx} \left[ \sin^2 \cot^{-1} \sqrt{\frac{1-x}{1+x}} \right]$  is

A. -1

B.  $\frac{1}{2}$

C.  $-\frac{1}{2}$

D. 1

**Answer:**



**Watch Video Solution**

23. The differential coefficient of  $f(\log_e x)$  w. r. t.  $x$ , where  $f(x) = \log_e x$ , is (i)

(i)  $\frac{x}{\ln x}$  (ii)  $\frac{\ln x}{x}$  (iii)  $\frac{1}{x \ln x}$  (iv)  $x \ln x$

A.  $\frac{x}{\log_e x}$

B.  $\frac{1}{x} \log_e x$

C.  $\frac{1}{x \log_e x}$

D. none of these

**Answer:**



**Watch Video Solution**

24. if  $f(x) = \sqrt{2x^2 - 1}$  and  $y = f(x^2)$  then  $\frac{dy}{dx}$  at  $x = 1$  is:

- A. 2
- B. 1
- C. -2
- D. none of these

**Answer:**



**Watch Video Solution**

25. If  $u = f(x^3)$ ,  $v = g(x^2)$ ,  $f'(x) = \cos x$ , and  $g'(x) = \sin x$ , then  $\frac{du}{dv}$  is

a)  $\frac{3}{2}x\cos x^3 \operatorname{cosec} x^2$  b)  $\frac{2}{3}\sin x^3 \sec x^2$  c)  $\tan x$  (d) none of these

A.  $\frac{3}{2}x \cos x^2 \operatorname{cosec} x^2$

B.  $\frac{3}{2} \sin x^3 \sec x^2$

C.  $\tan x$

D. none of these

**Answer:**



**Watch Video Solution**

26. A function  $f$ , defined for all positive real numbers, satisfies the equation  $f(x^2) = x^3$  for every  $x > 0$ . Then the value of  $f'(4)$  is 12 (b) 3 (c)  $3/2$  (d) cannot be determined

A. 12

B. 3

C.  $3/2$

D. cannot be determined

**Answer:**



**Watch Video Solution**

27. Let  $f: (-5, 5) \rightarrow R$  be a differentiable function of  $x$  with  $f(4) = 1, f'(4) = 1, f(0) = -1$  and  $f'(0) = 1$ . If  $g(x) = \left(f\left(2f^2(x) + 2\right)\right)^2$ , then  $g'(0)$  equals

A. 4

B. -4

C. 8

D. -8

**Answer:**



**Watch Video Solution**

28. The function  $f(x) = e^x + x$ , being differentiable and one-to-one, has a differentiable inverse  $f^{-1}(x)$ . The value of  $\frac{d}{dx}\left(f^{-1}\right)$  at the point  $f(\log 2)$  is  $\frac{1}{\ln 2}$  (b)  $\frac{1}{3}$  (c)  $\frac{1}{4}$  (d) none of these

A.  $\frac{1}{\ln 2}$

B.  $\frac{1}{3}$

C.  $\frac{1}{4}$

D. none of these

**Answer:**



**Watch Video Solution**

**29.** If  $f(x) = x + \tan x$  and  $f$  is the inverse of  $g$ , then  $g'(x)$  is equal to

A.  $\frac{1}{1 + [g(x) - x]^2}$

B.  $\frac{1}{2 - [g(x) - x]^2}$

C.  $\frac{1}{2 + [g(x) - x]^2}$

D. none of these

**Answer: C**



**Watch Video Solution**

**30.** If  $f(x) = x^3 + 3x + 4$  and  $g$  is the inverse function of  $f(x)$ , then the value of  $\frac{d}{dx} \left( \frac{g(x)}{g(g(x))} \right)$  at  $x = 4$  equals

A.  $-\frac{1}{6}$

B. 6

C.  $-\frac{1}{3}$

D. non-existent

**Answer:**



[Watch Video Solution](#)

**31.** If  $y = \frac{\sin^{-1}x}{\sqrt{1-x^2}}$ , then  $\frac{(1-x^2)dy}{dx}$  is equal to x + y (b) 1 + xy 1 - xy (d) xy - 2

A. x+y

B. 1+xy

C. 1-xy

D.  $xy - 2$

**Answer:**



**Watch Video Solution**

32. If  $y = \sqrt{\log x + \sqrt{\log x + \sqrt{\log x + \dots^\infty}}}$ , then  $\frac{dy}{dx}$  is

A.  $\frac{x}{2y - 1}$

B.  $\frac{x}{2y + 1}$

C.  $\frac{1}{x(2y - 1)}$

D.  $\frac{1}{x(1 - 2y)}$

**Answer:**



**Watch Video Solution**

33.  $\frac{d}{dx} \left[ \tan^{-1} \left( \frac{\sqrt{x}(3-x)}{1-3x} \right) \right] =$  (a)  $\frac{1}{2(1+x)\sqrt{x}}$  (b)  $\frac{3}{(1+x)\sqrt{x}}$  (c)  $\frac{2}{(1+x)\sqrt{x}}$  (d)

$$\frac{3}{2(1+x)\sqrt{x}}$$

A.  $\frac{1}{2(1+x)\sqrt{x}}$

B.  $\frac{3}{(1+x)\sqrt{x}}$

C.  $\frac{2}{(1+x)\sqrt{x}}$

D.  $\frac{3}{2(1+x)\sqrt{x}}$

**Answer:**



**Watch Video Solution**

34. Suppose the function  $f(x)-f(2x)$  has the derivative 5 at  $x=1$  and derivative 7 at  $x=2$ . The derivative of the function  $f(x)-f(4x)$  at  $x=1$  has the value equal to

A. 19

B. 9

C. 17

D. 14

**Answer:**



**Watch Video Solution**

35. If  $y\sqrt{x^2 + 1} = \log\left(\sqrt{x^2 + 1} - x\right)$ , show that  $x^2 + 1 \frac{dy}{dx} + xy + 1 = 0$

A. 0

B. 1

C. 2

D. none of these

**Answer:**



**Watch Video Solution**

**36.**

Let  $e^y = \frac{\sqrt{1+\alpha} + \sqrt{1-\alpha}}{\sqrt{1+\alpha} - \sqrt{1-\alpha}}$  and  $\tan \frac{x}{2} = \sqrt{\frac{1-\alpha}{1+\alpha}}$ ,  $\alpha \in [-1, 0] \cup (0, 1]$ . Then  $\left(\frac{dy}{dx}\right)$

A. 1/2

B. 1

C. 2

D. 1/3

**Answer:**



**Watch Video Solution**

**37.** The derivative of  $\tan^{-1} \left( \frac{\sqrt{1+x^2} - 1}{x} \right)$  with respect to  $\tan^{-1} \left( \frac{2x\sqrt{1-x^2}}{1-2x^2} \right)$  at  $x = 0$  is (a)  $\frac{1}{8}$  (b)  $\frac{1}{4}$  (c)  $\frac{1}{2}$  (d) 1

A. 1/8

B.  $1/4$

C.  $1/2$

D. 1

**Answer:**



**Watch Video Solution**

**38.** If  $\ln((e - 1)e^{xy} + x^2) = x^2 + y^2$  then  $\left(\frac{dy}{dx}\right)_{1,0}$  is equal to

A. 0

B. 1

C. 2

D. 3

**Answer:**



**Watch Video Solution**

39. If  $y = x^{x^x}$  then  $\frac{dy}{dx}$  is

A.  $y \left[ x^x (\ln x) \ln x + x^x \right]$

B.  $y \left[ x^x (\ln x) \ln x + x \right]$

C.  $y \left[ x^x (\ln x) \ln x + x^{-1} \right]$

D.  $y \left[ x^x (\ln e^x) \ln x + x^{-1} \right]$

**Answer:**



**Watch Video Solution**

40. The first derivative of the function  $\left[ \cos^{-1} \left( \sin \sqrt{\frac{1+x}{2}} \right) + x^x \right]$  with respect to  $x$  at  $x = 1$  is

A.  $3/4$

B.  $0$

C.  $1/2$

**Answer:**



**Watch Video Solution**

41. Let  $f(x) = x^3, x \in (0, \infty)$  and let  $g(x)$  be inverse of  $f(x)$ , then  $g'(x)$  must be

A.  $x(1 + \log x)$

B.  $x(1 + \log(x))$

C.  $\frac{1}{x(1 + \log g(x))}$

D. non-existent

**Answer:**



**Watch Video Solution**

**42.** If  $y = ax^{n+1} + bx^{-n}$ , then  $x^2 \frac{d^2y}{dx^2}$  is equal to

A.  $n(n-1)y$

B.  $n(n+1)y$

C.  $ny$

D.  $n^2y$

**Answer:**



Watch Video Solution

**43.** If  $y = ax^{n+1} + bx^{-n}$ , then  $x^2 \frac{d^2y}{dx^2}$  is equal to

A.  $m^2(ae^{mx} - be^{-mx})$

B. 1

C. 0

D. none of these

**Answer:**



**Watch Video Solution**

44. Suppose  $f(x) = e^{ax} + e^{bx}$ , where  $a \neq b$ , and that  $f'(x) - 2f(x) - 15f(x) = 0$  for all  $x$ . Then the value of  $ab$  is equal to:

A. 25

B. 9

C. -15

D. -9

**Answer:**



**Watch Video Solution**

45.  $\frac{d^{20}y}{dx^{20}}(2\cos x \cos 3x)$  is equal  $\rightarrow 2^{20}(\cos 2x - 2^{20} \cos 3x)$   
 $2^{20}(\cos 2x + 2^{20} \cos 4x)$   $2^{20}(\sin 2x + 2^{20} \sin 4x)$   $2^{20}(\sin 2x - 2^{20} \sin 4x)$

A.  $2^{20}(\cos 2x - 2^{20} \cos 3x)$

B.  $2^{20}(\cos 2x + 2^{20} \cos 4x)$

C.  $2^{20}(\sin 2x + 2^{20} \sin 4x)$

D.  $2^{20}(\sin 2x - 2^{20} \sin 4x)$

**Answer:**



**Watch Video Solution**

46.  $\frac{d^n}{dx^n}(\log x) = ?$  (a)  $\frac{(n-1)!}{x^n}$  (b)  $\frac{n!}{x^n}$  (c)  $\frac{(n-2)!}{x^n}$  (d)  $(-1)^{n-1} \frac{(n-1)!}{x^n}$

A.  $\frac{(n-1)!}{x^n}$

B.  $\frac{n!}{x^n}$

C.  $\frac{(n-2)!}{x^n}$

D.  $(-1)^{n-1} \frac{(n-1)!}{x^n}$

**Answer:**



**Watch Video Solution**

47. The  $n$ th derivative of the function  $f(x) = \frac{1}{1-x^2}$  [where  $x \in (-1, 1)$ ] at the point  $x = 0$  where  $n$  is even is (b)  $n!$  (c)  $n^n C_2$  (d)  $2^n C_2$

A. 0

B.  $n!$

C.  $b^n C_2$

D.  $2^n C_2$

**Answer:**



**Watch Video Solution**

48. If  $y = x \log\left\{\frac{x}{(a+bx)}\right\}$ , then show that  $x^3 \frac{d^2y}{dx^2} = \left(x \frac{dy}{dx} - y\right)^2$

A.  $x \frac{dy}{dx} - y$

B.  $\left(x \frac{dy}{dx} - y\right)^2$

C.  $y \frac{dy}{dx} - x$

D.  $\left( y \frac{dy}{dx} - x \right)^2$

**Answer:**



**Watch Video Solution**

49. If  $ax^2 + 2hxy + by^2 = 1$ , then  $\frac{d^2y}{dx^2}$  is  $\frac{h^2 - ab}{(hx + by)^2}$  (b)  $\frac{ab - h^2}{(hx + by)^2}$   $\frac{h^2 + ab}{(hx + by)^2}$

(d) none of these

A.  $\frac{h^2 - ab}{(hx + by)^3}$

B.  $\frac{ab - h^2}{(hx + by)^2}$

C.  $\frac{h^2 + ab}{(hx + by)^2}$

D. none of these

**Answer:**



**Watch Video Solution**

**50.** If  $y^{1/m} = \left(x + \sqrt{1+x^2}\right)$ , then  $(1+x^2)y_2 + xy_1$  is (where  $y_r$  represents the  $r$ th derivative of  $y$  w.r.t.  $x$ )

- A.  $m^2y$
- B.  $my^2$
- C.  $m^2y^2$
- D. none of these

**Answer:**



[Watch Video Solution](#)

**51.** If  $(\sin x)(\cos y) = \frac{1}{2}$ , then  $\frac{d^2y}{dx^2}$  at  $\left(\frac{\pi}{4}, \frac{\pi}{4}\right)$  is -4 (b) -2 (c) -6 (d) 0

- A. -4
- B. -2
- C. -6

D. 0

**Answer:**



**Watch Video Solution**

52. A function  $f$  satisfies the condition

$f(x) = f'(x) + f''(x) + \dots + f^{(n)}(x)$ , where  $f(x)$  is a differentiable function indefinitely and dash denotes the order of derivative. If

$f(0) = 1$ , then  $f(x)$  is (a)  $e^{\frac{x}{2}}$  (b)  $e^x$  (c)  $e^{2x}$  (d)  $e^{4x}$

A.  $e^{x/2}$

B.  $e^x$

C.  $e^{2x}$

D.  $e^{4x}$

**Answer:**



**Watch Video Solution**

53. Let  $f(x)$  be a polynomial of degree 3 such that  $f(3) = 1$ ,  $f'(3) = -1$ ,  $f''(3) = 0$ , and  $f'''(3) = 12$ . Then the value of  $f'(1)$  is (a) 12 (b) 23 (c) -13 (d) none of these

A. 12

B. 23

C. -13

D. none of these

**Answer:**



**Watch Video Solution**

54. If  $y^2 = ax^2 + bx + c$ , then  $y^3 \frac{d^2y}{dx^2}$  is (a) a constant (b) a function of  $x$  only (c) a function of  $y$  only (d) a function of  $x$  and  $y$

A. a constant

B. a function of  $x$  only

C. a function of y only

D. a function of x and y

**Answer:**



Watch Video Solution

55. If  $y = x + e^x$ , then  $\frac{d^2y}{dx^2}$  is (a)  $e^x$  (b)  $-\frac{e^x}{(1+e^x)^3}$  (c)  $\frac{e^x}{(1+e^x)^3}$  (d)  $-\frac{1}{(1+e^x)^3}$

A.  $(-\sin x + e^x)^{-1}$

B.  $\frac{\sin x - e^x}{(\cos x + e^x)^2}$

C.  $\frac{\sin x - e^x}{(\cos x + e^x)^3}$

D.  $\frac{\sin x + e^x}{(\cos x + e^x)^3}$

**Answer:**



Watch Video Solution

56. if  $y = \sin mx$ , then the value of the determinant

$$\begin{vmatrix} y & y_1 & y_2 \\ y_3 & y_4 & y_5 \\ y_6 & y_7 & y_8 \end{vmatrix}$$

Where  $y_n = \frac{d^n y}{dx^n}$  is

- A. 1
- B. 0
- C. -1
- D. none of these

**Answer:**



**Watch Video Solution**

57. If  $f''(x) = -f(x)$  and  $g(x) = f'(x)$  and  $F(x) = \left(f\left(\frac{x}{2}\right)\right)^2 + \left(g\left(\frac{x}{2}\right)\right)^2$  and given

that  $F(5) = 5$ , then  $F(10)$  is 5 (b) 10 (c) 0 (d) 15

A. 5

B. 10

C. 0

D. 15

**Answer:**



**Watch Video Solution**

**58.** Let  $y = \ln(1 + \cos x)^2$ . Then the value of  $\frac{d^2y}{dx^2} + \frac{2}{e^{\frac{y}{2}}}$  equal (a) 0 (b)

$$\frac{2}{1 + \cos x} \quad (\text{c}) \frac{4}{1 + \cos x} \quad (\text{d}) \frac{-4}{(1 + \cos x)^2}$$

A. 0

B.  $\frac{2}{1 + \cos x}$

C.  $\frac{4}{1 + \cos x}$

D.  $\frac{-4}{(1 + \cos x)^2}$

**Answer:**



**Watch Video Solution**

**59.**  $x = t \cos t, y = t + \sin t$ . Then  $(d^2x)/(dy^2)$  at "t=(pi)/(2)" is

A.  $\frac{\pi + 4}{2}$

B.  $-\frac{\pi + 4}{2}$

C. -2

D. none of these

**Answer:**



**Watch Video Solution**

**60.** If  $f(x) = (x - 1)^4(x - 2)^3(x - 3)^2(x - 4)$ , then the value of  $f'''(1) + f'(2) + f(3) + f(4)$  equals

A. 0

B. 50

C. 324

D. 648

**Answer:** 648



**Watch Video Solution**

**61.** let  $y = t^{10} + 1$ , and  $x = t^8 + 1$ , then  $\frac{d^2y}{dx^2}$  is

A.  $\frac{5}{2}t$

B.  $20t^8$

C.  $\frac{5}{16t^6}$

D. none of these

**Answer:**



**Watch Video Solution**

62. If  $x = \log p$  and  $y = \frac{1}{p}$ , then

A.  $\frac{d^2y}{dx^2} - 2p = 0$

B.  $\frac{d^2y}{dx^2} + y = 0$

C.  $\frac{d^2y}{dx^2} + \frac{dy}{dx} = 0$

D.  $\frac{d^2y}{dx^2} - \frac{dy}{dx} = 0$

**Answer:**



Watch Video Solution

63. If  $x = \phi(t)$ ,  $y = \psi(t)$ , then  $\frac{d^2y}{dx^2}$  is

A.  $\frac{\phi' \psi'' - \psi' \phi''}{(\phi')^2}$

B.  $\frac{\phi' \psi'' - \psi' \phi''}{(\phi')^3}$

C.  $\frac{\phi''}{\psi''}$

D.  $\frac{\psi'}{\phi'}$

**Answer:**



**Watch Video Solution**

**64.** If  $f(x) = x^4 \tan(x^3) - x \ln(1 + x^2)$ , then the value of  $\frac{d^4(f(x))}{dx^4}$  at  $x = 0$  is

A. 0

B. 6

C. 12

D. 24

**Answer:**



**Watch Video Solution**

**65.** If graph of  $y = f(x)$  is symmetrical about the  $y$ -axis and that of  $y = g(x)$

is symmetrical about the origin and if  $h(x) = f(x)g(x)$ , then  $\frac{d^3h(x)}{dx^3} \text{ at } x = 0$  is

cannot be determined (b)  $f(0)g(0)$  0 (d) none of these

A. cannot be determined

B.  $f(0) \cdot g(0)$

C. 0

D. none of these

**Answer:**



[Watch Video Solution](#)

**66.** Let  $g(x)$  be the inverse of an invertible function  $f(x)$ , which is

differentiable for all real  $x$ . Then  $g''(f(x))$  equals.

A.  $-\frac{f'(x)}{(f'(x))^3}$

- B.  $\frac{f(x)f'(x) - (f(x))^3}{f(x)}$
- C.  $\frac{f(x)f'(x) - (f'(x))^2}{(f'(x))^2}$
- D. none of these

**Answer:**



**Watch Video Solution**

67.  $f(x) = e^x - e^{-x} - 2\sin x - \frac{2}{3}x^3$ . Then the least value of n for which  $\frac{d^n}{dx^n}f(x) \mid_{x=0}$  is nonzero is

- A. 5
- B. 6
- C. 7
- D. 8

**Answer:**



**Watch Video Solution**

68. Let  $y = f(x)$  and  $x = \frac{1}{z}$ . If  $\frac{d^2y}{dx^2} = \lambda(z^3) \frac{dy}{dz} + z^4 \frac{d^2y}{dz^2}$ , then the value of  $\lambda$  is

A. 1

B. 2

C.  $\frac{1}{2}$

D.  $\frac{1}{4}$

**Answer:**



**Watch Video Solution**

69. Let  $x=f(t)$  and  $y=g(t)$ , where  $x$  and  $y$  are twice differentiable function. If

$f'(0)=g'(0)=f''(0)=2$ .  $g''(0)=6$ , then the value of  $\left(\frac{d^2y}{dx^2}\right)_{t=0}$  is equal to

A. 0

B. 1

C. 2

D. 3

**Answer:**



**Watch Video Solution**

70. If  $f(x)$  satisfies the relation  $f\left(\frac{5x - 3y}{2}\right) = \frac{5f(x) - 3f(y)}{2}$   $\forall x, y \in R$ , and  $f(0) = 3$  and  $f'(0) = 2$ , then the period of  $\sin(f(x))$  is 2 $\pi$  (b)  $\pi$  (c) 3 $\pi$  (d) 4 $\pi$

A. 2 $\pi$

B.  $\pi$

C. 3 $\pi$

D. 4 $\pi$

**Answer:**



**Watch Video Solution**

71. A function  $f: \vec{R} \rightarrow \vec{R}$  satisfies

$$\sin x \cos y \left( f(2x + 2y) - f(2x - 2y) \right) = \cos x \sin y \left( f(2x + 2y) + f(2x - 2y) \right) \quad \text{If}$$

$$f'(0) = \frac{1}{2}, \text{ then } a \left( f''(x) = f(x) = 0 \right) b \left( 4f''(x) + f(x) = 0 \right) c \left( f''(x) + f(x) = 0 \right) d \left( 4f''(x) - f(x) = 0 \right)$$

A.  $f'(x) = f(x) = 0$

B.  $4f'(x) + f(x) = 0$

C.  $f'(x) + f(x) = 0$

D.  $4f'(x) - f(x) = 0$

**Answer:**



[Watch Video Solution](#)

72. If  $y = a\sin x + b\cos x$ , then  $y^2 + \left(\frac{dy}{dx}\right)^2$  is a function of  $x$  (b) function of  $y$  function of  $x$  and  $y$  (d) constant

- A. function of  $x$
- B. function of  $y$
- C. function of  $x$  and  $y$
- D. constant

**Answer:**



**Watch Video Solution**

73.  $\int \frac{dx}{\sqrt{(1-\sin 2x)/(1+\sin 2x)}}$  is equal to, (0

- A.  $\sec^2 x$
- B.  $-\sec^2\left(\frac{\pi}{4} - x\right)$
- C.  $\sec^2\left(\frac{\pi}{4} + x\right)$

D.  $\sec^2\left(\frac{\pi}{4} - x\right)$

**Answer:**



**Watch Video Solution**

74. If  $f(x) = |\cos x| + |\sin x|$ , then  $\frac{dy}{dx}$  at  $x = \frac{2\pi}{3}$  is equal to

A.  $\frac{1 - \sqrt{3}}{2}$

B. 0

C.  $\frac{1}{2}(\sqrt{3} - 1)$

D. none of these

**Answer:**



**Watch Video Solution**

75. If  $f(x) = \left|\log_e|x|\right|$ , then  $f'(x)$  equals

A.  $\frac{1}{|x|}$ , where  $x \neq 0$

B.  $\frac{1}{x}$  for  $|x| > 1$  and  $-\frac{1}{x}$  for  $|x| < 1$

C.  $-\frac{1}{x}$  for  $|x| > 1$  and  $\frac{1}{x}$  for  $|x| < 1$

D.  $\frac{1}{x}$  for  $x > 0$  and  $-\frac{1}{x}$  for  $x < 0$

**Answer:**



**Watch Video Solution**

76. If  $f(x) = \sqrt{1 - \sin 2x}$ , then  $f'(x)$  is equal to

(a)  $-(\cos x + \sin x)$ , for  $x \in \left(\frac{\pi}{4}, \frac{\pi}{2}\right)$

(b)  $\cos x + \sin x$ , for  $x \in \left(0, \frac{\pi}{4}\right)$

(c)  $-(\cos x + \sin x)$ , for  $x \in \left(0, \frac{\pi}{4}\right)$

(d)  $\cos x - \sin x$ , for  $x \in \left(\frac{\pi}{4}, \frac{\pi}{2}\right)$

A.  $-(\cos x + \sin x)$ , for  $x \in (\pi/4, \pi/2)$

B.  $\cos x + \sin x$  for  $x \in (0, \pi/4)$

C.  $-(\cos x + \sin x)$ , for  $x \in (0, \pi/4)$

D.  $\cos x - \sin x$ , for  $x \in (\pi/4, \pi/2)$

**Answer:**



**Watch Video Solution**

77. Instead of the usual definition of derivative  $Df(x)$ , if we define a new

kind of derivative  $D^*F(x)$  by the formula

$$D^*(x) = (\lim_{h \rightarrow 0}) \frac{f^2(x+h) - f^2(x)}{h}, \text{ where } f^2(x)$$

$[f(x)]^2$  and  $f(x) = x \log x$ , then  $D^*f(x)(|)_{x=e}$  has the value

(B) 2e (C) 4e (d)  
none of these

A. e

B. 2e

C. 4e

D. none of these

**Answer:**



**Watch Video Solution**

78. if  $y = \frac{(a - x)\sqrt{a - x} - (b - x)\sqrt{x - b}}{\sqrt{a - x} + \sqrt{x - b}}$  then  $\frac{dy}{dx}$  wherever it is defined is equal to :

A.  $\frac{x + (a + b)}{\sqrt{(a - x)(x - b)}}$

B.  $\frac{2x - a - b}{2\sqrt{a - x}\sqrt{x - b}}$

C.  $-\frac{(a + b)}{2\sqrt{(a - x)(x - b)}}$

D.  $\frac{2x + (a + b)}{2\sqrt{(a - x)(x - b)}}$

**Answer:**



**Watch Video Solution**

79. The derivative of  $y = (1-x)(2-x)\dots(n-x)$  at  $x=1$  is

A. 0

B.  $(-1)(n - 1)!$

C.  $n! - 1$

D.  $(-1)^{n-1}(n - 1)!$

**Answer:**



**Watch Video Solution**

80. If  $y = \sqrt{\frac{1-x}{1+x}}$ , then  $\left(1-x^2\right)\frac{dy}{dx}$  is equal to

A.  $y^2$

B.  $1/y$

C.  $-y$

D.  $-y/x$

**Answer:**



**Watch Video Solution**

81. If  $y = \frac{\sqrt{a+x} - \sqrt{a-x}}{\sqrt{a+x} + \sqrt{a-x}}$ , then  $\frac{dy}{dx}$  is equal to (a)  $\frac{ay}{x\sqrt{a^2 - x^2}}$  (b)  $\frac{ay}{\sqrt{a^2 - x^2}}$   
 (c)  $\frac{ay}{x\sqrt{a^2 - x^2}}$  (d) none of these

A.  $\frac{ay}{x\sqrt{a^2 - x^2}}$

B.  $\frac{ay}{\sqrt{a^2 - x^2}}$

C.  $\frac{ay}{x\sqrt{x^2 - a^2}}$

D. none of these

**Answer:**



Watch Video Solution

82. Let  $u(x)$  and  $v(x)$  be differentiable functions such that  
 $\frac{u(x)}{v(x)} = 7$ .  $\frac{u'(x)}{v'(x)} = p$  and  $\left(\frac{u(x)}{v(x)}\right)' = q$ , then  $\frac{p+q}{p-q}$  has the value of to 1 (b) 0

(c) 7 (d) -7

A. 1

B. 0

C. 7

D. -7

**Answer:**



**Watch Video Solution**

83. If  $\sin^{-1} \left( \frac{x^2 - y^2}{x^2 + y^2} \right) = \log a$ , then  $\frac{dy}{dx}$  is equal to (a)  $\frac{x}{y}$  (b)  $\frac{y}{x^2}$  (c)  $\frac{x^2 - y^2}{x^2 + y^2}$  (d)  $\frac{y}{x}$

A.  $\frac{x}{y}$

B.  $\frac{y}{x^2}$

C.  $\frac{x^2 - y^2}{x^2 + y^2}$

D.  $\frac{y}{x}$

**Answer:**



**Watch Video Solution**

**84.** Let  $h(x)$  be differentiable for all  $x$  and let  $f(x) = (kx + e^x)h(x)$ , where  $k$  is some constant. If  $h(0) = 5$ ,  $h'(0) = -2$ , and  $f'(0) = 18$ , then the value of  $k$  is

A. 5

B. 4

C. 3

D. 2.2

**Answer:**



**Watch Video Solution**

**85.** If  $\lim_{t \rightarrow x} \frac{e^{tf(x)} - e^{xt}f(t)}{(t-x)(f(x))^2} = 2$  and  $f(0) = \frac{1}{2}$ , then find the value of  $f'(0)$ .

A. 4

B. 2

C. 0

D. 1

**Answer:**



**Watch Video Solution**

**86.** If  $f(0) = 0$ ,  $f'(0) = 2$ , then the derivative of  $y = f(f(f(x)))$  at  $x = 0$  is 2 (b)

8 (c) 16 (d) 4

A. 2

B. 8

C. 16

D. 4

**Answer:**



Watch Video Solution

87. If  $f(x) = \sqrt{1 + \cos^2(x^2)}$ , then  $f' \left( \frac{\sqrt{\pi}}{2} \right)$  is  $\frac{\sqrt{\pi}}{6}$  (b)  $-\sqrt{\pi/6}$  (c)  $1/\sqrt{6}$  (d)  $\pi/\sqrt{6}$

A.  $\sqrt{\pi/6}$

B.  $-\sqrt{\pi/6}$

C.  $1/\sqrt{6}$

D.  $\pi/\sqrt{6}$

Answer:



Watch Video Solution

88.  $\frac{d}{dx} \cos^{-1} \sqrt{\cos x}$ ,  $0 < x < \frac{\pi}{2}$  is equal to

A.  $\frac{1}{2} \sqrt{1 + \sec x}$

B.  $\sqrt{1 + \sec x}$

C.  $-\frac{1}{2}\sqrt{1 + \sec x}$

D.  $-\sqrt{1 + \sec x}$

**Answer:**



**Watch Video Solution**

89. If  $\tan y = \frac{2^x}{1 + 2^{2x+1}}$ , then  $\frac{dy}{dx} \text{ at } x = 0$  is

A. 1

B. 2

C.  $\ln 2$

D.  $-\frac{1}{10} \ln 2$

**Answer:**



**Watch Video Solution**

90. If  $y = \left(x + \sqrt{x^2 + a^2}\right)^n$ , then  $\frac{dy}{dx}$  is  $\frac{ny}{\sqrt{x^2 + a^2}}$  (b) -  $\frac{ny}{\sqrt{x^2 + a^2}}$  -  $\frac{nx}{\sqrt{x^2 + a^2}}$  (d)  
-  $\frac{nx}{\sqrt{x^2 + a^2}}$

A.  $\frac{ny}{\sqrt{x^2 + a^2}}$

B. -  $\frac{ny}{\sqrt{x^2 + a^2}}$

C.  $\frac{nx}{\sqrt{x^2 + a^2}}$

D. -  $\frac{nx}{\sqrt{x^2 + a^2}}$

**Answer:**



[Watch Video Solution](#)

91. if  $y = \log_{\sin x} \tan x$  then  $\left(\frac{dy}{dx}\right)_{\frac{\pi}{4}}$  is

A.  $\frac{4}{\log 2}$

B. -4log2

C.  $\frac{-4}{\log 2}$

D. none of these

**Answer:**



**Watch Video Solution**

92.  $\frac{d}{dx} \left[ \sin^2 \cot^{-1} \sqrt{\frac{1-x}{1+x}} \right]$  is

A. -1

B.  $\frac{1}{2}$

C.  $-\frac{1}{2}$

D. 1

**Answer:**



**Watch Video Solution**

**93.** The differential coefficient of  $f(\log_e x)$  w. r. t.  $x$ , where  $f(x) = \log_e x$ , is

A.  $\frac{x}{\log_e x}$

B.  $\frac{1}{x} \log_e x$

C.  $\frac{1}{x \log_e x}$

D. none of these

**Answer:**



**Watch Video Solution**

**94.** if  $f(x) = \sqrt{2x^2 - 1}$  and  $y = f(x^2)$  then  $\frac{dy}{dx}$  at  $x = 1$  is:

A. 2

B. 1

C. -2

D. none of these

**Answer:**



**Watch Video Solution**

**95.** If  $u = f(x^3)$ ,  $v = g(x^2)$ ,  $f'(x) = \cos x$ , and  $g'(x) = \sin x$ , then  $\frac{du}{dv}$  is

A.  $\frac{3}{2}x \cos x^3 \operatorname{cosec} x^2$

B.  $\frac{3}{2} \sin x^3 \sec x^2$

C.  $\tan x$

D. none of these

**Answer:**



**Watch Video Solution**

**96.** A function  $f$ , defined for all positive real numbers, satisfies the equation  $f(x^2) = x^3 = x^3$  for every  $x > 0$ . Then the value of  $f'(4)$  is

A. 12

B. 3

C. 3/2

D. cannot be determined

**Answer:**



**Watch Video Solution**

**97.** Let  $f: (-5, 5) \rightarrow \mathbb{R}$  be a differentiable function of with  $f(4) = 1, f'(4) = 1, f(0) = -1$  and  $f'(0) = I$ . If  $g(x) = \left(f\left(2f^2(x) + 2\right)\right)^2$ , then  $g'(0)$  equals

A. 4

B. -4

C. 8

D. -8

**Answer:**



**Watch Video Solution**

**98.** The function  $f(x) = e^x + x$ , being differentiable and one- to -one, has a differentiable inverse  $f^{-1}(x)$ . The value of  $\frac{x}{dx} \left( f^{-1}atf(\log 2) \right)$  is

A.  $\frac{1}{In2}$

B.  $\frac{1}{3}$

C.  $\frac{1}{4}$

D. none of these

**Answer:**



**Watch Video Solution**

**99.** If  $f(x) = x + \tan x$  and  $f$  is the inverse of  $g$ , then  $g'(x)$  is equal to

A.  $\frac{1}{1 + [g(x) - x]^2}$

B.  $\frac{1}{2 - [g(x) - x]^2}$

C.  $\frac{1}{2 + [g(x) - x]^2}$

D. none of these

**Answer: C**



**Watch Video Solution**

**100.** If  $f(x) = x^3 + 3x + 4$  and  $g$  is the inverse function of  $f(x)$ , then the

value of  $\frac{d}{dx} \left( \frac{g(x)}{g(g(x))} \right)$  at  $x = 4$  equals

A.  $\frac{-1}{6}$

B. 6

C.  $\frac{-1}{3}$

D. non-existent

**Answer:**



Watch Video Solution

101. If  $y = \frac{\sin^{-1}x}{\sqrt{1-x^2}}$ , then  $\frac{(1-x^2)dy}{dx}$  is equal to  $x+y$  (b)  $1+xy$  (c)  $1-xy$  (d)

$xy - 2$

A.  $x+y$

B.  $1+xy$

C.  $1-xy$

D.  $xy-2$

**Answer:**



Watch Video Solution

102. If  $y = \sqrt{\log x + \sqrt{\log x + \sqrt{\log x + \dots^\infty}}}$ , then  $\frac{dy}{dx}$  is

A.  $\frac{x}{2y-1}$

- B.  $\frac{x}{2y + 1}$
- C.  $\frac{1}{x(2y - 1)}$
- D.  $\frac{1}{x(1 - 2y)}$

**Answer:**



**Watch Video Solution**

103.  $\frac{d}{dx} \left[ \tan^{-1} \left( \frac{\sqrt{x}(3-x)}{1-3x} \right) \right] =$

- A.  $\frac{1}{2(1+x)\sqrt{x}}$
- B.  $\frac{3}{(1+x)\sqrt{x}}$
- C.  $\frac{2}{(1+x)\sqrt{x}}$
- D.  $\frac{3}{2(1+x)\sqrt{x}}$

**Answer:**



**Watch Video Solution**

**104.** Suppose the function  $f(x) - f(2x)$  has the derivative 5 at  $x = 1$  and derivative 7 at  $x = 2$ . The derivative of the function  $f(x) - f(4x)$  at  $x = 1$  has the value equal to 19 (b) 9 (c) 17 (d) 14

A. 19

B. 9

C. 17

D. 14

**Answer:**



**Watch Video Solution**

**105.** If  $y\sqrt{x^2 + 1} = \log\left(\sqrt{x^2 + 1} - x\right)$ , show that  $(x^2 + 1)\frac{dy}{dx} + xy + 1 = 0$



**Watch Video Solution**

**106.**

Let  $e^y = \frac{\sqrt{1+\alpha} + \sqrt{1-\alpha}}{\sqrt{1+\alpha} - \sqrt{1-\alpha}}$  and  $\tan \frac{x}{2} = \sqrt{\frac{1-\alpha}{1+\alpha}}$ ,  $\alpha \in [-1, 0] \cup (0, 1]$ . Then  $\left(\frac{dy}{dx}\right)$

A. 1/2

B. 1

C. 2

D. 1/3

**Answer:**



**Watch Video Solution**

**107.** The derivative of  $\tan^{-1} \left( \frac{\sqrt{1+x^2} - 1}{x} \right)$  with respect to  $x$  at  $x = 0$  is (a)  $\frac{1}{8}$  (b)  $\frac{1}{4}$  (c)  $\frac{1}{2}$  (d) 1

A. 1/8

B.  $1/4$

C.  $1/2$

D. 1

**Answer:**



**Watch Video Solution**

**108.** If  $\ln((e - 1)e^{xy} + x^2) = x^2 + y^2$  then  $\left(\frac{dy}{dx}\right)_{1,0}$  is equal to

A. 0

B. 1

C. 2

D. 3

**Answer:**



**Watch Video Solution**

**109.** If  $y = (x^x)^x$  then  $\frac{dy}{dx}$  is

A.  $y \left[ x^x (\log ex) \log x + x^x \right]$

B.  $y \left[ x^x (\log ex) \log x + x \right]$

C.  $y \left[ x^x (\log ex) \log x + x^{-1} \right]$

D.  $y \left[ x^x \left( \log_e x \right) \log x + x^{-1} \right]$

**Answer:**



**Watch Video Solution**

**110.** The first derivative of the function  $\left[ \cos^{-1} \left( \sin \sqrt{\frac{1+x}{2}} \right) + x^x \right]$  with respect to  $x$  at  $x = 1$  is

A.  $3/4$

B.  $0$

C.  $1/2$

D. -1/2

**Answer:**



**Watch Video Solution**

**111.**  $f(x) = x^x$ ,  $x \in (0, \infty)$  and let  $g(x)$  be inverse of  $f(x)$  , then  $g(x)'$  must be

A.  $x(1 + \log x)$

B.  $x(1 + \log(x))$

C.  $\frac{1}{x(1 + \log g(x))}$

D. non-existent

**Answer:**



**Watch Video Solution**

**112.** If  $y = ax^{n+1} + bx^{-n}$ , then  $x^2 \frac{d^2y}{dx^2}$  is equal to (a)  $n(n - 1)y$  (b)  $n(n + 1)y$  (c)  $ny$  (d)

$$n^2y$$

A.  $n(n-1)y$

B.  $n(n+1)y$

C.  $ny$

D.  $n^2y$

**Answer:**



**Watch Video Solution**

**113.** If  $y = ax^{n+1} + bx^{-n}$ , then  $x^2 \frac{d^2y}{dx^2}$  is equal to (a)  $n(n - 1)y$  (b)  $n(n + 1)y$  (c)  $ny$  (d)

$$n^2y$$

A.  $m^2(ae^{mx} - be^{-mx})$

B. 1

C. 0

D. none of these

**Answer:**



**Watch Video Solution**

114. Suppose  $f(x) = e^{ax} + e^{bx}$ , where  $a \neq b$ , and that  $f''(x) - 2f'(x) - 15f(x) = 0$

for all  $x$ . Then the product  $ab$  is

A. 25

B. 9

C. -15

D. -9

**Answer:**



**Watch Video Solution**

$$115. \frac{d^{20}y}{dx^{20}}(2\cos x \cos 3x) \text{ is equal } \rightarrow a) 2^{20}(\cos 2x - 2^{20}\cos 3x)$$

$$b) 2^{20}(\cos 2x + 2^{20}\cos 4x) c) 2^{20}(\sin 2x + 2^{20}\sin 4x) d) 2^{20}(\sin 2x - 2^{20}\sin 4x)$$

A.  $2^{20}(\cos 2x - 2^{20}\cos 3x)$

B.  $2^{20}(\cos 2x + 2^{20}\cos 4x)$

C.  $2^{20}(\sin 2x + 2^{20}\sin 4x)$

D.  $2^{20}(\sin 2x - 2^{20}\sin 4x)$

**Answer:**



**Watch Video Solution**

$$116. \frac{d^n}{dx^n}(\log x) = \frac{(n-1)!}{x^n}$$
 (b)  $\frac{n!}{x^n} \frac{(n-2)!}{x^n}$  (d)  $(-1)^{n-1} \frac{(n-1)!}{x^n}$

A.  $\frac{(n-1)!}{x^n}$

B.  $\frac{n!}{x^n}$

C.  $\frac{(n-2)!}{x^n}$

$$D. (-1)^{n-1} \frac{(n-1)!}{x^n}$$

**Answer:**



**Watch Video Solution**

117. If  $y = x \log\left(\frac{x}{a+bx}\right)$ , then  $x^3 \frac{d^2y}{dx^2} =$

A.  $x \frac{dy}{dx} - y$

B.  $\left(x \frac{dy}{dx} - y\right)^2$

C.  $y \frac{dy}{dx} - x$

D.  $\left(y \frac{dy}{dx} - x\right)^2$

**Answer:**



**Watch Video Solution**

**118.** If  $ax^2 + 2hxy + by^2 = 1$ , then  $\frac{d^2y}{dx^2}$  is  $\frac{h^2 - ab}{(hx + by)^2}$  (b)  $\frac{ab - h^2}{(hx + by)^2}$   $\frac{h^2 + ab}{(hx + by)^2}$

(d) none of these

A.  $\frac{h^2 - ab}{(hx + by)^3}$

B.  $\frac{ab - h^2}{(hx + by)^2}$

C.  $\frac{h^2 + ab}{(hx + by)^2}$

D. none of these

**Answer:**



Watch Video Solution

**119.** If  $y^{1/m} = \left(x + \sqrt{1 + x^2}\right)$ , then  $(1 + x^2)y_2 + xy_1$  is (where  $y_r$

represents the  $r$ th derivative of  $y$  w.r.t.  $x$ )

A.  $m^2y$

B.  $my^2$

C.  $m^2y^2$

D. none of these

**Answer:**



**Watch Video Solution**

**120.** If  $(\sin x)(\cos y) = 1/2$ , then  $d^2y/dx^2$  at  $(\pi/4, \pi/4)$  is

A. -4

B. -2

C. -6

D. 0

**Answer:**



**Watch Video Solution**

**121.** A function  $f$  satisfies the condition  $f(x) = f(x) + f'(x) + f''(x) + \dots$ , where  $f(x)$  is a differentiable function indefinitely and dash denotes the order the derivative. If  $f(0) = 1$ , then  $f(x)$  is

A.  $e^{x/2}$

B.  $e^x$

C.  $e^{2x}$

D.  $e^{4x}$

**Answer:**



**Watch Video Solution**

**122.** Let  $f(x)$  be a polynomial of degree 3 such that  $f(3) = 1$ ,  $f'(3) = -1$ ,  $f''(3) = 0$ , and  $f'''(3) = 12$ . Then the value of  $f'(1)$  is

A. 12

B. 23

C. -13

D. none of these

**Answer:**



**Watch Video Solution**

123. If  $y^2 = ax^2 + bx + c$ , then  $y^3 \frac{d^2y}{dx^2}$  is (a) a constant (b) a function of  $x$  only (c) a function of  $y$  only (d) a function of  $x$  and  $y$

A. a constant

B. a function of  $x$  only

C. a function of  $y$  only

D. a function of  $x$  and  $y$

**Answer:**



**Watch Video Solution**

**124.** If  $y = \sin x + e^x$ , then  $\frac{d^2y}{dx^2} =$

A.  $(-\sin x + e^x)^{-1}$

B.  $\frac{\sin x - e^x}{(\cos x + e^x)^2}$

C.  $\frac{\sin x - e^x}{(\cos x + e^x)^3}$

D.  $\frac{\sin x + e^x}{(\cos x + e^x)^3}$

**Answer:**



**Watch Video Solution**

**125.** If  $f''(x) = -f(x)$  and  $g(x) = f'(x)$  and  $F(x) = \left(f\left(\frac{x}{2}\right)\right)^2 + \left(g\left(\frac{x}{2}\right)\right)^2$

and given that  $F(5) = 5$ , then  $F(10)$  is

A. 5

B. 10

C. 0

D. 15

**Answer:**



**Watch Video Solution**

126. Let  $y = \ln(1 + \cos x)^2$ . The value of  $\frac{d^2y}{dx^2} + \frac{2}{e^{y/2}}$  equals

A. 0

B.  $\frac{2}{1 + \cos x}$

C.  $\frac{4}{1 + \cos x}$

D.  $\frac{-4}{(1 + \cos x)^2}$

**Answer:**



**Watch Video Solution**

127.  $x = t \cos t$ ,  $y = t + \sin t$ . Then  $\frac{d^2x}{dy^2}$  is (a)  $\frac{\pi + 4}{2}$  (b)  $-\frac{\pi + 4}{2}$  (c) -2 (d) none of these

A.  $\frac{\pi + 4}{2}$

B.  $-\frac{\pi + 4}{2}$

C. -2

D. none of these

**Answer:**



[Watch Video Solution](#)

128. Let  $y = t^{10} + 1$  and  $x = t^8 + 1$ . Then  $\frac{d^2y}{dx^2}$  is

A.  $\frac{5}{2}t$

B.  $20t^8$

C.  $\frac{5}{16t^6}$

D. none of these

**Answer:**



**Watch Video Solution**

129. If  $x = \log p$  and  $y = \frac{1}{p}$ , then

A.  $\frac{d^2y}{dx^2} - 2p = 0$

B.  $\frac{d^2y}{dx^2} + y = 0$

C.  $\frac{d^2y}{dx^2} + \frac{dy}{dx} = 0$

D.  $\frac{d^2y}{dx^2} - \frac{dy}{dx} = 0$

**Answer:**



**Watch Video Solution**

130. If  $x = \phi(t)$ ,  $y = \psi(t)$ , then  $\frac{d^2y}{dx^2}$  is  $\frac{\phi' \psi - \psi' \phi'}{(\phi')^2}$  (b)  $\frac{\phi' \psi - \psi' \phi'}{(\phi')^3}$  (c)  $\frac{\phi'}{\psi'}$  (d)  $\frac{\psi'}{\phi'}$

A.  $\frac{\phi' \psi' - \psi' \phi'}{(\phi')^2}$

B.  $\frac{\phi' \psi' - \psi' \phi'}{(\phi')^3}$

C.  $\frac{\phi'}{\psi'}$

D.  $\frac{\psi'}{\phi'}$

**Answer:**



**Watch Video Solution**

131. If  $f(x) = x^4 \tan(x^3) - x \ln(1 + x^2)$ , then the value of  $\frac{d^4(f(x))}{dx^4}$  at  $x = 0$  is

0 (b) 6 (c) 12 (d) 24

A. 0

B. 6

C. 12

**Answer:****Watch Video Solution**

132. Let  $g(x)$  be the inverse of an invertible function  $f(x)$ , which is differentiable for all real  $x$ . Then  $g''(f(x))$  equals

A.  $-\frac{f'(x)}{(f(x))^3}$

B.  $\frac{f(x)f'(x) - (f(x))^3}{f(x)}$

C.  $\frac{f(x)f'(x) - (f(x))^2}{(f(x))^2}$

D. none of these

**Answer:****Watch Video Solution**

**133.**  $f(x) = e^x - e^{-x} - 2\sin x - \frac{2}{3}x^3$ . Then the least value of n for which  $\frac{d^n}{dx^n}f(x) \mid_{x=0}$  is nonzero is

A. 5

B. 6

C. 7

D. 8

**Answer:**



**Watch Video Solution**

**134.** Let  $y = f(x)$  and  $x = \frac{1}{z}$ . If  $\frac{d^2y}{dx^2} = \lambda z^3 \frac{dy}{dz} + z^4 \frac{d^2y}{dz^2}$ , then the value of  $\lambda$  is

A. 1

B. 2

C.  $\frac{1}{2}$

D.  $\frac{1}{4}$

**Answer:**



**Watch Video Solution**

135. Let  $x=f(t)$  and  $y=g(t)$ , where  $x$  and  $y$  are twice differentiable function. If

$f'(0) = g'(0) = f''(0) = 2$ .  $g''(0) = 6$ , then the value of  $\left(\frac{d^2y}{dx^2}\right)_{t=0}$  is equal to

A. 0

B. 1

C. 2

D. 3

**Answer:**



**Watch Video Solution**

136. If  $f(x)$  satisfies the relation  $f\left(\frac{5x - 3y}{2}\right) = \frac{5f(x) - 3f(y)}{2}$   $\forall x, y \in R$ , and  $f(0) = 3$  and  $f'(0) = 2$ , then the period of  $\sin(f(x))$  is 2 $\pi$  (b)  $\pi$  (c) 3 $\pi$  (d) 4 $\pi$

A.  $2\pi$

B.  $\pi$

C.  $3\pi$

D.  $4\pi$

**Answer:**



**Watch Video Solution**

137. A function

$f: R \rightarrow R$  satisfies  $\sin x \cos y (f(2x + 2y) - f(2x - 2y)) = \cos x \sin y (f(2x + 2y) + f(2x - 2y))$

" If " $f'(0) = 1/2$ ", then

A.  $f'(x) = f(x) = 0$

B.  $4f'(x) + f(x) = 0$

C.  $f'(x) + f(x) = 0$

D.  $4f'(x) - f(x) = 0$

**Answer:**



**Watch Video Solution**

### Multiple Correct Answers Type

1. If  $y = e^{\sqrt{x}} + e^{-\sqrt{x}}$ , then  $\frac{dy}{dx}$  is equal to (a)  $\frac{e^{\sqrt{x}}}{2\sqrt{x}}$  (b)  $\frac{e^{\sqrt{x}} - e^{-\sqrt{x}}}{2\sqrt{x}}$  (c)  $\frac{1}{2\sqrt{x}}\sqrt{y^2 - 4}$

(d)  $\frac{1}{2\sqrt{x}}\sqrt{y^2 + 4}$

A.  $\frac{e^{\sqrt{x}} - e^{-\sqrt{x}}}{2\sqrt{x}}$

B.  $\frac{e^{\sqrt{x}} - e^{-\sqrt{x}}}{2x}$

C.  $\frac{1}{2\sqrt{x}}\sqrt{y^2 - 4}$

D.  $\frac{1}{2\sqrt{x}}\sqrt{y^2 + 4}$

**Answer:**



**Watch Video Solution**

2. Let  $y = \sqrt{x + \sqrt{x + \sqrt{x + \infty}}}$ ,  $\frac{dy}{dx}$  is equal to (a)  $\frac{1}{2y - 1}$  (b)  $\frac{x}{x + 2y}$  (c)  $\frac{1}{\sqrt{1 + 4x}}$  (d)

$$\frac{y}{2x + y}$$

A.  $\frac{1}{2y - 1}$

B.  $\frac{x}{x + 2y}$

C.  $\frac{1}{\sqrt{1 + 4x}}$

D.  $\frac{y}{2x + y}$

**Answer:**



**Watch Video Solution**

3. If  $f(\theta) = \tan\left(\sin^{-1}\sqrt{\frac{2}{3 + \cos 2\theta}}\right)$ , then find  $f'(\theta)$  and

A.  $f\left(\frac{\pi}{4}\right) = 1$

B.  $f\left(\frac{\pi}{4}\right) = \sqrt{2}$

C.  $\frac{d(f(\theta))}{d(\cos\theta)}$  at  $\theta = \frac{\pi}{4}$  is - 2

D.  $f\left(\frac{\pi}{4}\right) = \sqrt{2}$

**Answer:**



**Watch Video Solution**

4.  $f(x) = |x^2 - 3x|x| + 2|$ . Then which of the following is/are true ?

A.  $f'(x) = 2x - 3$  for  $x \in (0, 1) \cup (2, \infty)$

B.  $f'(x) = 2x + 3$  or  $x \in (-\infty, -2) \cup (-1, 0)$

C.  $f'(x) = -2x - 3$  for  $x \in (-2, -1)$

D. None of these

**Answer:**



5. Let  $f(x) = \frac{\sqrt{x} - 2\sqrt{x-1}}{\sqrt{x-1} - 1} \cdot x$  then

A.  $f(10) = 1$

B.  $f'(3/2) = -1$

C. domain of  $f(x)$  is  $x \geq 1$

D. range of  $f(x)$  is  $(-2, -1] \cup (2, 00)$

**Answer:**



Watch Video Solution

6. If  $x^3 - 2x^2y^2 + 5x + y - 5 = 0$  and  $y(1) = 1$ , then  $y'(1) = \frac{4}{3}$  (b)  $y^1 = -\frac{4}{3}$   
 $y^1 = -8\frac{22}{27}$  (d)  $y'(1) = \frac{2}{3}$

A.  $y'(1) = 4/3$

B.  $y''(1) = -4/3$

C.  $y''(1) = -8\frac{22}{27}$

D.  $y'(1) = 2/3$

**Answer:**



**Watch Video Solution**

7. If  $y = x^{(\log x)^{\log(\log x)}}$  then  $\frac{dy}{dx} =$

A.  $\frac{y}{x} \left( \left( \ln x^{x-1} \right) + 2 \ln x \ln(\ln x) \right)$

B.  $\frac{y}{x} (\log x)^{\log(\log x)} (2\log(\log x) + 1)$

C.  $\frac{y}{x \ln x} \left[ (\ln x)^2 + 2\ln(\ln x) \right]$

D.  $\frac{y \log y}{x \log x} [2\log(\log x) + 1]$

**Answer: B**



**Watch Video Solution**

**8.** If  $f(x - y)$ ,  $f(x)f(y)$ , and  $f(x + y)$  are in A.P. for all  $x, y$ , and  $f(0) \neq 0$ , then

A.  $f(4) = f(-4)$

B.  $f(2) + f(-2) = 0$

C.  $f'(4) + f'(-4) = 0$

D.  $f'(2) = f'(-2)$

**Answer:**



**Watch Video Solution**

**9.** If  $y = \cos^{-1}\left(\frac{2x}{1+x^2}\right)$ , then  $\frac{dy}{dx}$  is (a)  $\frac{-2}{1+x^2}$  for all  $x$  (b)  $\frac{-2}{1+x^2}$  for all  $|x| < 1$   
 $\frac{2}{1+x^2}$  for  $|x| > 1$  (d) none of these

A.  $\frac{-2}{1+x^2}$  for all  $x$

B.  $\frac{-2}{1+x^2}$  for all  $|x| < 1$

C.  $\frac{2}{1+x^2}$  for  $|x| > 1$

D. none of these

**Answer:**



**Watch Video Solution**

**10.**  $f_n(x) = e^{f_{n-1}(x)}$  for all  $n \in N$  and  $f_0(x) = x$ , then  $\frac{d}{dx} \{f_n(x)\}$  is

A.  $f_n(x) \frac{d}{dx} \{f_{n-1}(x)\}$

B.  $f_n(x)f_{n-1}(x)$

C.  $f_n(x)f_{n-1}(x) \dots f_2(x). f_1(x)$

D. None of these

**Answer:**



**Watch Video Solution**

11. Suppose  $f$  and  $g$  are functions having second derivatives  $f'$  and  $g'$  every where, if  $f(x) \cdot g(x) = 1$  for all  $x$  and  $f', g''$  are never zero then

$$\frac{f'(x)}{f(x)} - \frac{g''(x)}{g'(x)}$$
 equals

A.  $\frac{-2f'(x)}{f(x)}$

B.  $\frac{-2g'(x)}{g(x)}$

C.  $\frac{-f'(x)}{f(x)}$

D.  $\frac{2f'(x)}{f(x)}$

**Answer:**



Watch Video Solution

12. If  $y = e^{-x} \cos x$  and  $y_n + k_n y = 0$  where  $y_n = \frac{d^n y}{dx^n}$  and  $k_n$  are constant  
 $n \in N$  then

A.  $k_4 = 4$

B.  $k_8 = -16$

C.  $k_{12} = 20$

D.  $k_{16} = -24$

**Answer: B**



**Watch Video Solution**

13. If a function is represented parametrically by the equations

$$x = \frac{1 + (\log)_e t}{t^2}; y = \frac{3 + 2(\log)_e t}{t}, \text{ then which of the following statements are true?}$$
$$y^{x-2xy'} = y \quad yy' = 2x(y')^2 + 1 \quad xy' = 2y(y')^2 + 2 \quad y^{y-4xy'} = (y')^2$$

A.  $y''(x - 2xy') = y$

B.  $yy' = 2x(y')^2 + 1$

C.  $xy' = 2y(y')^2 + 2$

D.  $y''(y - 4xy') = (y')^2$

**Answer:**



**Watch Video Solution**

14. If  $y = \frac{x^2}{2} + \frac{1}{2}x\sqrt{x^2 + 1} + \frac{1}{2}(\log)_e\sqrt{x + \sqrt{x^2 + 1}}$ , prove that

$2y = xy' + (\log)_e y'$ , where  $y'$  denotes the derivative w.r.t  $x$ .

A.  $y' = x + \sqrt{x^2 + 1}$

B.  $y' = \frac{1}{x + \sqrt{x^2 + 1}}$

C.  $2y = xy' + \log_e y'$

D.  $2y = xy' - \log_e y'$

**Answer:**



**Watch Video Solution**

15. A curve given parametrically by

$x = t + t^3$  and  $y = t^2$ , where  $t \in R$ . For what value(s) of  $t$  is  $\frac{dy}{dx} = \frac{1}{2}$ ?

A.  $1/3$

B. 2

C. 3

D. 1

**Answer:**



**Watch Video Solution**

**16.** If  $e^{\sin(x^2+y^2)} = \tan\frac{y^2}{4} + \sin^{-1}x$ , then  $y'(0)$  can be

A.  $\frac{1}{3\sqrt{\pi}}$

B.  $-\frac{1}{3\sqrt{\pi}}$

C.  $-\frac{1}{5\sqrt{\pi}}$

D.  $-\frac{1}{3\sqrt{5\pi}}$

**Answer:**



**Watch Video Solution**

17. If  $g$  is the inverse of a function  $f$  and  $f'(x) = \frac{1}{1+x^5}$  then  $g'(x)$  is equal to  
(1)  $1+x^5$  (2)  $5x^4$  (3)  $\frac{1}{1+\{g(x)\}^5}$  (4)  $1+\{g(x)\}^5$

- A. a unique point in the interval  $\left(n, n + \frac{1}{2}\right)$
- B. a unique point in the interval  $\left(n + \frac{1}{2}, n + 1\right)$
- C. a unique point in the interval  $(n, n + 1)$
- D. two points in the interval  $(n, n + 1)$

**Answer:**



**Watch Video Solution**

18. If for  $x \in \left(0, \frac{1}{4}\right)$ , the derivative of  $\tan^{-1}\left(\frac{6x\sqrt{x}}{1-9x^3}\right)$  is  $\sqrt{x}g(x)$ , then  $g(x)$  equals:  
(1)  $\frac{3x}{1-9x^3}$  (2)  $\frac{3}{1+9x^3}$  (3)  $\frac{9}{1+9x^3}$  (4)  $\frac{3x\sqrt{x}}{1-9x^3}$

A.  $\lim_{x \rightarrow 0^+} f\left(\frac{1}{x}\right) = 1$

B.  $\lim_{x \rightarrow 0^+} xf\left(\frac{1}{x}\right) = 2$

C.  $\lim_{x \rightarrow 0^+} x^2 f'(x) = 0$

D.  $|f(x)| \leq 2$  for all  $x \in (0, 2)$

**Answer:**



**Watch Video Solution**

19. Let  $f(x) = x \sin \pi x$ ,  $x > 0$ . Then for all natural numbers  $n$ ,  $f'(x)$  vanishes

at a unique point in the interval  $\left(n, n + \frac{1}{2}\right)$  a unique point in the interval

$\left(n + \frac{1}{2}, n + 1\right)$  a unique point in the interval  $(n, n + 1)$  two points in the

interval  $(n, n + 1)$

A. a unique point in the interval  $\left(n, n + \frac{1}{2}\right)$

B. a unique point in the interval  $\left(n + \frac{1}{2}, n + 1\right)$

C. a unique point in the interval  $(n, n + 1)$

D. two points in the interval  $(n, n + 1)$

**Answer: A and C**



**Watch Video Solution**

20. Let  $f: (0, \infty) \rightarrow R$  be a differentiable function such that  $f'(x) = 2 - \frac{f(x)}{x}$

for all  $x \in (0, \infty)$  and  $f(1) = 1$ , then

A.  $\lim_{x \rightarrow 0^+} f\left(\frac{1}{x}\right) = 1$

B.  $\lim_{x \rightarrow 0^+} xf\left(\frac{1}{x}\right) = 2$

C.  $\lim_{x \rightarrow 0^+} x^2 f'(x) = 0$

D.  $|f(x)| \leq 2$  for all  $x \in (0, 2)$

**Answer:**



**Watch Video Solution**

21. Let  $f: R \rightarrow R$  and  $h: R \rightarrow R$  be differentiable functions such that  $f(x) = x^3 + 3x + 2$ ,  $g(f(x)) = x$  and  $h(g(g(x))) = x$  for all  $x \in R$ . Then,  $h'(1)$  equals.



Watch Video Solution

22. If  $y = e^{\sqrt{x}} + e^{-\sqrt{x}}$ , then  $\frac{dy}{dx}$  is equal to

A.  $\frac{e^{\sqrt{x}} - e^{-\sqrt{x}}}{2\sqrt{x}}$

B.  $\frac{e^{\sqrt{x}} - e^{-\sqrt{x}}}{2x}$

C.  $\frac{1}{2\sqrt{x}}\sqrt{y^2 - 4}$

D.  $\frac{1}{2\sqrt{x}}\sqrt{y^2 + 4}$

**Answer:**



Watch Video Solution

**23.** Let  $y = \sqrt{x + \sqrt{x + \sqrt{x + \dots}}}$ , then  $\frac{dy}{dx}$  is equal to

A.  $\frac{1}{2y - 1}$

B.  $\frac{x}{x + 2y}$

C.  $\frac{1}{\sqrt{1 + 4x}}$

D.  $\frac{y}{2x + y}$

**Answer:**



**Watch Video Solution**

**24.** If  $f(x - y)$ ,  $f(x)f(y)$ , and  $f(x + y)$  are in A.P. for all  $x, y$ , and  $f(0) \neq 0$ , then

A.  $f(4) = f(-4)$

B.  $f(2) + f(-2) = 0$

C.  $f'(4) + f'(-4) = 0$

D.  $f'(2) = f'(-2)$

**Answer:**



**Watch Video Solution**

**25.** If a function is represented parametrically by the equations

$x = \frac{1 + \log_e t}{t^2}, y = \frac{3 + 2\log_e t}{t}$ , then which of the following statements are true ?

A.  $y''(x - 2xy') = y$

B.  $yy' = 2x(y')^2 + 1$

C.  $xy' = 2y(y')^2 + 2$

D.  $y''(y - 4xy') = (y')^2$

**Answer:**



**Watch Video Solution**

**26.** If  $y = \frac{x^2}{2} + \frac{1}{2}x\sqrt{x^2 + 1} + \log_e \sqrt{x + \sqrt{x^2 + 1}}$ , then

A.  $y' = x + \sqrt{x^2 + 1}$

B.  $y' = \frac{1}{x + \sqrt{x^2 + 1}}$

C.  $2y = xy' + \log_e y'$

D.  $2y = xy' - \log_e y'$

**Answer:**



**Watch Video Solution**

27. A curve given by

$x = t + t^3$  and  $y = t^2$ , where  $t \in R$ . For what value(s) of  $t$  is  $\frac{dy}{dx} = \frac{1}{2}$ ?

A.  $1/3$

B. 2

C. 3

D. 1

**Answer:**



Watch Video Solution

28. Let  $f(x) = x \sin \pi x$ ,  $x > 0$ . Then for all natural numbers  $n$ ,  $f'(x)$  vanishes

at a unique point in the interval  $\left(n, n + \frac{1}{2}\right)$  a unique point in the interval

$\left(n + \frac{1}{2}, n + 1\right)$  a unique point in the interval  $(n, n + 1)$  two points in the

interval  $(n, n + 1)$

A. a unique point in the interval  $\left(n, n + \frac{1}{2}\right)$

B. a unique point in the interval  $\left(n + \frac{1}{2}, n + 1\right)$

C. a unique point in the interval  $(n, n + 1)$

D. two points in the interval  $(n, n + 1)$

**Answer:**



Watch Video Solution

29. Let  $f: (0, \infty) \rightarrow R$  be a differentiable function such that  $f'(x) = 2 - \frac{f(x)}{x}$

for all  $x \in (0, \infty)$  and  $f(1) = 1$ , then  $f(x)$  is

A.  $\lim_{x \rightarrow 0^+} f\left(\frac{1}{x}\right) = 1$

B.  $\lim_{x \rightarrow 0^+} xf\left(\frac{1}{x}\right) = 2$

C.  $\lim_{x \rightarrow 0^+} x^2 f'(x) = 0$

D.  $|f(x)| \leq 2$  for all  $x \in (0, 2)$

**Answer:**



**Watch Video Solution**

### Linked Comprehension Type

1.  $f(x)$  is a polynomial function,  $f: R \rightarrow R$ , such that  $f(2x) = f(x)f'(x)$ .

$f(x)$  is

A. 4

B. 12

C. 15

D. none of these

**Answer: B**



**Watch Video Solution**

2.  $f(x)$  is a polynomial function,  $f: R \rightarrow R$ , such that  $f(2x) = f(x)f'(x)$ .  $f(x)$  is (A) one-one and onto (B) one-one and into (C) many-one and onto (D) many-one and into

A. one-one and onto

B. one-one and into

C. many-one and onto

D. many-one and into

**Answer:**



3.  $f(x)$  is a polynomial function,  $f: R \rightarrow R$ , such that  $f(2x) = f(x)f'(x)$ .

Equation  $f(x) = x$  has (A) three real and positive roots (B) three real and negative roots (C) one real root (D) three real roots such that sum of roots is zero

A. three real and positive roots

B. three real and negative roots

C. one real root

D. three real roots such that sum of roots is zero

**Answer:**



4.  $f: R \rightarrow R$ ,  $f(x) = x^3 + x^2f'(1) + xf'(2) + f''(3)$  for all  $x \in R$ .

The value of  $f(1)$  is

A. 2

B. 3

C. -1

D. 4

**Answer:**



**Watch Video Solution**

5.  $f: R \rightarrow R$ ,  $f(x) = x^3 + x^2f'(1) + xf''(2) + f'''(3)$  for all  $x \in R$ .

$f(x)$  is

A. one-one and onto

B. one-one and into

C. many-one and onto

D. many-one and into

**Answer:**



Watch Video Solution

6.  $f: R \rightarrow R$ ,  $f(x) = x^3 + x^2f(1) + xf'(2) + f''(3)$  for all  $x \in R$ .

The value of  $f(1) + f'(2) + f''(3)$  is

A. 0

B. -1

C. 2

D. 3

**Answer:**



Watch Video Solution

7. Repeated roots : If equation  $f(x) = 0$ , where  $f(x)$  is a polynomial function, has roots  $\alpha, \alpha, \beta, \dots$  or  $\alpha$  root is repreated root, then  $f(x) = 0$  is equivalent to  $(x - \alpha)^2(x - \beta)\dots = 0$ , from which we can conclude that  $f(x) = 0$  or  $2(x - \alpha)[(x - \beta)\dots] + (x - \alpha)^2[(x - \beta)\dots]' = 0$  or  $(x - \alpha)[2\{(x - \beta)\dots\} + (x - \alpha)(x - \beta)\dots] = 0$ .

has root  $\alpha$ . Thus, if  $\alpha$  root occurs twice in the equation, then it is common in equations  $f(x) = 0$  and  $f'(x) = 0$ . Similarly, if  $\alpha$  root occurs thrice in equation, then it is common in the equations  $f(x)=0$ ,  $f'(x)=0$ , and  $f''(x)=0$ .

If  $x-c$  is a factor of order  $m$  of the polynomial  $f(x)$  of degree  $n$  ( $1 < m < n$ ), then  $x=c$  is a root of the polynomial [where  $f^r(x)$  represent rth derivative of  $f(x)$  w.r.t.  $x$ ]

A.  $f^m(x)$

B.  $f^{m-1}(x)$

C.  $f^n(x)$

D. none of these

**Answer:**



**Watch Video Solution**

**8. Repeated roots :** If equation  $f(x) = 0$ , where  $f(x)$  is a polynomial function, has roots  $\alpha, \alpha, \beta, \dots$  or  $\alpha$  root is repreated root, then  $f(x) = 0$  is equivalent to  $(x - \alpha)^2(x - \beta)\dots = 0$ , from which we can conclude that

$$f(x) = 0 \text{ or } 2(x - \alpha)[(x - \beta)\dots] + (x - \alpha)^2[(x - \beta)\dots]' = 0 \text{ or } (x - \alpha)[2\{(x - \beta)\dots\} + (x - \beta)\dots]' = 0$$

has root  $\alpha$ . Thus, if  $\alpha$  root occurs twice in the equation, then it is common in equations  $f(x) = 0$  and  $f'(x) = 0$ . Similarly, if  $\alpha$  root occurs thrice in equation, then it is common in the equations  $f(x)=0$ ,  $f'(x)=0$ , and  $f'''(x)=0$ .

If  $a_1x^3 + b_1x^2 + c_1x + d_1 = 0$  and  $a_2x^3 + b_2x^2 + c_2x + d_2 = 0$  have a pair of repeated roots common, then

$$\begin{vmatrix} 3a_1 & 2b_1 & c_1 \\ 3a_2 & 2b_2 & c_2 \\ a_2b_1 - a_1b_2 & c_1a_2 - c_2a_1 & d_1a_2 - d_2a_1 \end{vmatrix} =$$

A. 0

B. 1

C. -1

D. 2

**Answer:**



[View Text Solution](#)

**9. Repeated roots :** If equation  $f(x) = 0$ , where  $f(x)$  is a polynomial function, has roots  $\alpha, \alpha, \beta, \dots$  or  $\alpha$  root is repeated root, then  $f(x) = 0$  is equivalent to  $(x - \alpha)^2(x - \beta)\dots = 0$ , from which we can conclude that  $f(x) = 0$  or  $2(x - \alpha)[(x - \beta)\dots] + (x - \alpha)^2[(x - \beta)\dots]' = 0$  or  $(x - \alpha)[2\{(x - \beta)\dots\} + (x - \alpha)] = 0$ . Thus, if  $\alpha$  root occurs twice in the equation, then it is common in equations  $f(x) = 0$  and  $f'(x) = 0$ . Similarly, if  $\alpha$  root occurs thrice in equation, then it is common in the equations  $f(x)=0$ ,  $f'(x)=0$ , and  $f''(x)=0$ . If  $x=c$  is a factor of order  $m$  of the polynomial  $f(x)$  of degree  $n$  ( $1 < m < n$ ), then  $x=c$  is a root of the polynomial [where  $f^r(x)$  represent  $r$ th derivative of  $f(x)$  w.r.t.  $x$ ]

A. If  $p < q < n$ , then  $\alpha$  and  $\beta$  are two of the roots of the equation

$$f^{p-1}(x) = 0.$$

B. If  $q < p < n$ , then  $\alpha$  and  $\beta$  are two of the roots of the equation

$$f^{q-1}(x) = 0.$$

C. If  $p < q < n$ , then equations  $f(x) = 0$  and  $f^p(x) = 0$  have exactly one root common

D. If  $q < p < n$ , then equations  $f^q(x) = 0$  and  $f^p(x) = 0$  have exactly two roots common.

**Answer:**



**Watch Video Solution**

**10.** Equation  $x^n - 1 = 0, n > 1, n \in N$ , has roots  $1, a_1, a_2, \dots, a_{n-1}$ .

The value of  $(1 - a_1)(1 - a_2) \dots (1 - a_{n-1})$  is

A.  $n^2/2$

B. n

C.  $(-1)^n n$

D. none of these

**Answer:**



**Watch Video Solution**

**11.** Equation  $x^n - 1 = 0$ ,  $n > 1$ ,  $n \in N$ , has roots  $1, a_1, a_2, \dots, a_n$ . The value

of  $\sum_{r=2}^n \frac{1}{2 - a_r}$ , is

A.  $\frac{2^{n-1}(n-2) + 1}{2^n - 1}$

B.  $\frac{2^n(n-2) + 1}{2^n - 1}$

C.  $\frac{2^{n-1}(n-1) - 1}{2^n - 1}$

D. none of these

**Answer:**



**Watch Video Solution**

**12.** Equation  $x^n - 1 = 0$ ,  $n > 1$ ,  $n \in N$ , has roots  $1, a_1, a_2, \dots, a_{n-1}$ .

The value of  $\sum_{r=1}^{n-1} \frac{1}{2 - a_r}$  is

A.  $\frac{n}{4}$

B.  $\frac{n(n-1)}{2}$

C.  $\frac{n - 1}{2}$

D. none of these

**Answer:**



**Watch Video Solution**

13.  $f(x) = x^2 + xg'(1) + g''(2)$  and  $g(x) = f(1)x^2 + xf'(x) + f(x)$ .

The value of  $f(3)$  is

A. 1

B. 0

C. -1

D. -2

**Answer:**



**Watch Video Solution**

**14.**  $f(x) = x^2 + xg'(1) + g''(2)$  and  $g(x) = f(1)x^2 + xf'(x) + f(x)$ .

The value of  $g(0)$  is

A. 0

B. -3

C. 2

D. none of these

**Answer:** C



[Watch Video Solution](#)

**15.**  $f(x) = x^2 + xg'(1) + g''(2)$  and  $g(x) = f(1)x^2 + xf'(x) + f(x)$ .

The domain of the function  $\sqrt{\frac{f(x)}{g(x)}}$  is

A.  $(-\infty, 1] \cup (2, 3]$

B.  $(-2, 0] \cup (1, \infty)$

C.  $(-\infty, 0] \cup (2/3, 3]$

D. none of these

**Answer:**



**Watch Video Solution**

**16.**  $g(x + y) = g(x) + g(y) + 3xy(x + y) \quad \forall x, y \in R$  and  $g'(0) = -4$ .

Number of real roots of the equation  $g(x) = 0$  is

A. 2

B. 0

C. 1

D. 3

**Answer:**



**Watch Video Solution**

**17.**  $g(x + y) = g(x) + g(y) + 3xy(x + y) \forall x, y \in R$  and  $g'(0) = -4$ .

For which of the following values of x is  $\sqrt{g(x)}$  not defined ?

A.  $[-2, 0]$

B.  $[2, \infty)$

C.  $[-1, 1]$

D. none of these

**Answer:** [-1,1]



**Watch Video Solution**

**18.**  $g(x + y) = g(x) + g(y) + 3xy(x + y) \forall x, y \in R$  and  $g'(0) = -4$ .

The value of  $g'(1)$  is

A. 0

B. 1

C. -1

D. none of these

**Answer:**



**Watch Video Solution**

19. A curve is represented parametrically by the equations

$$x = f(t) = a^{\ln(b^t)} \text{ and } y = g(t) = b^{-\ln(a^t)} \quad a, b > 0 \text{ and } a \neq 1, b \neq 1 \text{ Where } t \in$$

The value of  $\frac{d^2y}{dx^2}$  at the point where  $f(t)=g(t)$  is

A.  $\frac{1}{f(t)^2}$

B.  $-(g(t))^2$

C.  $\frac{-g(t)}{f(t)}$

D.  $\frac{-f(t)}{g(t)}$

**Answer:**



**Watch Video Solution**

20. A curve is represented parametrically by the equations

$$x = f(t) = a^{\ln(b^t)} \text{ and } y = g(t) = b^{-\ln(a^t)} \quad a, b > 0 \text{ and } a \neq 1, b \neq 1$$

Where  $t \in \mathbb{R}$ . The value of  $\frac{d^2y}{dx^2}$  at the point where  $f(t)=g(t)$  is

A. 0

B.  $\frac{1}{2}$

C. 1

D. 2

**Answer:**



[Watch Video Solution](#)

21. A curve is represented parametrically by the equations

$$x = f(t) = a^{\ln(b^t)} \text{ and } y = g(t) = b^{-\ln(a^t)} \quad a, b > 0 \text{ and } a \neq 1, b \neq 1$$

Where  $t \in \mathbb{R}$ .

The value of  $\frac{f(t)}{f'(t)} \cdot \frac{f'(-t)}{f'(-t)} + \frac{f(-t)}{f'(-t)} \cdot \frac{f'(-t)}{f'(t)}$   $\forall t \in \mathbb{R}$  is

A. -2

B. 2

C. -4

D. 4

**Answer:**



**Watch Video Solution**

**22.** Let  $f: R \rightarrow R$  be a differentiable function satisfying

$f(x + y) = f(x) + f(y) + x^2y + xy^2$  for all real numbers  $x$  and  $y$ . If

$$\lim_{x \rightarrow 0} \frac{f(x)}{x} = 1, \text{ then}$$

The value of  $f(9)$  is

A. 8

B. 10

C. 12

D. 18

**Answer:**



**Watch Video Solution**

23. Let  $f: R \rightarrow R$  be a differentiable function satisfying  $f(x + y) = f(x) + f(y) + x^2y + xy^2$  for all real numbers  $x$  and  $y$ . If  $\lim_{x \rightarrow 0} \frac{f(x)}{x} = 1$ , then

The value of  $f(9)$  is

A. 240

B. 356

C. 252

D. 730

**Answer: C**



**Watch Video Solution**

24. If roots of an equation  $x^n - 1 = 0$  are  $1, a_{1,2}, a_{n-1}$ , then the value of

$(1 - a_1)(1 - a_2)(1 - a_3)\dots(1 - a_{n-1})$  will be

A.  $n^2/2$

B. n

C.  $(-1)^n n$

D. none of these

**Answer:**



Watch Video Solution

**Matrix Match Type**

**1. Match the following lists:**

List I	List II
<p>a. If <math>f(x)</math> is an integrable function for  <math>x \in \left[\frac{\pi}{6}, \frac{\pi}{3}\right]</math> and</p> $I_1 = \int_{\pi/6}^{\pi/3} \sec^2 \theta f(2\sin 2\theta) d\theta, \text{ and}$ $I_2 = \int_{\pi/6}^{\pi/3} \operatorname{cosec}^2 \theta f(2\sin 2\theta) d\theta, \text{ then } I_1/I_2 =$	p. 3
<p>b. If <math>f(x+1) = f(3+x) \forall x</math>, and the value of  <math>\int_a^{a+b} f(x) dx</math> is independent of <math>a</math>, then the  value of <math>b</math> can be</p>	q. 1
<p>c. The value of</p> $2 \int_1^4 \frac{\tan^{-1}[x^2]}{\tan^{-1}[x^2] + \tan^{-1}[25+x^2-10x]} dx$ <p>(where <math>[.]</math> denotes the greatest integer function) is</p>	r. 2
<p>d. If <math>I = \int_0^2 \sqrt{x + \sqrt{x + \sqrt{x + \dots}}} dx</math></p> <p>(where <math>x &gt; 0</math>), then <math>[I]</math> is equal to (where <math>[.]</math> denotes the greatest integer function)</p>	s. 4



**Watch Video Solution**

**2. Match the following lists:**

List I	List II
<p>a. If <math>f(x)</math> is an integrable function for  <math>x \in \left[\frac{\pi}{6}, \frac{\pi}{3}\right]</math> and</p> $I_1 = \int_{\pi/6}^{\pi/3} \sec^2 \theta f(2\sin 2\theta) d\theta, \text{ and}$ $I_2 = \int_{\pi/6}^{\pi/3} \operatorname{cosec}^2 \theta f(2\sin 2\theta) d\theta, \text{ then } I_1/I_2 =$	p. 3
<p>b. If <math>f(x+1) = f(3+x) \forall x</math>, and the value of  <math>\int_a^{a+b} f(x) dx</math> is independent of <math>a</math>, then the  value of <math>b</math> can be</p>	q. 1
<p>c. The value of</p> $2 \int_1^4 \frac{\tan^{-1}[x^2]}{\tan^{-1}[x^2] + \tan^{-1}[25+x^2-10x]} dx$ <p>(where <math>[.]</math> denotes the greatest integer function) is</p>	r. 2
<p>d. If <math>I = \int_0^2 \sqrt{x + \sqrt{x + \sqrt{x + \dots}}} dx</math></p> <p>(where <math>x &gt; 0</math>), then <math>[I]</math> is equal to (where <math>[.]</math> denotes the greatest integer function)</p>	s. 4



**Watch Video Solution**

### 3. Match the following lists :



**View Text Solution**

### 4. Match Column-I to II

**Column-I**

- (A) Tetragonal and Hexagonal
- (B) Cubic and Rhombohedral
- (C) Monoclinic and Triclinic
- (D) Cubic and Orthorhombic

**Column-II**

- (P) are two crystal systems
- (Q)  $\alpha = \beta = \gamma$
- (R)  $a \neq b \neq c$
- (S)  $a = b = c$

**Watch Video Solution**

### 5. Match List I with List II and choose the correct answer

List I	List II
A Hypothalamus	1. Sperm lysins
B Acrosome	2. Estrogen
C Graafian follicle	3. Relaxin
D Leydig cells	4. GnRH
E Parturition	5. Testosterone

- A.  $a \ b \ c \ d$   
 $q \ p \ s \ r$
- B.  $a \ b \ c \ d$   
 $s \ p \ q \ r$
- C.  $a \ b \ c \ d$   
 $r \ q \ s \ p$
- D.  $a \ b \ c \ d$   
 $q \ p \ r \ s$

**Answer:**



**Watch Video Solution**

- 6.** Match the terms given in Column I with the compound given in Column II.

<b>Column I</b>	<b>Column II</b>
A. Acid rain	1. $\text{CHCl}_2\text{-CHF}_2$
B. Photochemical smog	2. CO
C. Combination with haemoglobin	3. $\text{CO}_2$
D. Depletion of ozone layer	4. $\text{SO}_2$
	5. Unsaturated hydrocarbons

- A.  $a \ b \ c \ d$   
 $s \ r \ q \ p$
- B.  $a \ b \ c \ d$   
 $q \ s \ r \ p$
- C.  $a \ b \ c \ d$   
 $s \ r \ p \ q$

<i>a</i>	<i>b</i>	<i>c</i>	<i>d</i>
<i>q</i>	<i>s</i>	<i>p</i>	<i>r</i>

**Answer:**



**Watch Video Solution**

### Numerical Value Type

1.  $f'(x) = \varphi'(x) = f(x)$  for all  $x$ . Also,  $f(3) = 5$  and  $f'(3) = 4$ . Then the value of  $[f(10)]^2 - [\varphi(10)]^2$  is \_\_\_\_\_



**Watch Video Solution**

2. If  $y=f(x)$  is an odd differentiable function defined on  $(-\infty, \infty)$  such that  $f'(3) = -2$ , then  $f'(-3)$  equal to-



**Watch Video Solution**

3. If  $x^3 + 3x^2 - 9x + c$  is of the form  $(x - \alpha)^2(x - \beta)$  then  $c$  is equal to \_\_\_\_\_



[Watch Video Solution](#)

4. If graph of  $y = f(x)$  is symmetrical about the point  $(5, 0)$  and  $f'(7) = 3$ ,  
then the value of  $f'(3)$  is \_\_\_\_\_



[Watch Video Solution](#)

5. Let  $g(x) = f(x)\sin x$ , where  $f(x)$  is a twice differentiable function on  $(-\infty, \infty)$  such that  $f'(-\pi) = 1$ . The value of  $g''(-\pi)$  equals \_\_\_\_\_



[Watch Video Solution](#)

6. Let  $f(x) = (x - 1)(x - 2)(x - 3)(x - n)$ ,  $n \in N$ , and  $f(n) = 5040$ . Then the value of  $n$  is \_\_\_\_\_



[Watch Video Solution](#)

7.  $y = f(x)$ , where  $f$  satisfies the relation  $f(x + y) = 2f(x) + xy(y) + y\sqrt{f(x)}$   $\forall x, y \in R$  and  $f'(0) = 0$ . Then  $f(6)$  is equal of  $f(-3)$  is \_\_\_\_\_



**Watch Video Solution**

8. If function  $f$  satisfies the relation  $f(x) \cdot f'(-x) = f(-x) \cdot f'(x)$  for all  $x$ , and  $f(0) = 3$ , and if  $f(3) = 3$ , then the value of  $f(-3)$  is \_\_\_\_\_



**Watch Video Solution**

9. If  $y = \frac{a + bx^{\frac{3}{2}}}{x^{\frac{5}{4}}}$  and  $y' = 0$  at  $x = 5$ , then the value of  $\frac{a^2}{b^2}$  is \_\_\_\_\_



**Watch Video Solution**

10. Prove that  $\frac{2^{\log_2^{1/4}x} - 3^{\log_{27}(x^2+1)^3} - 2x}{7^{4\log_{49}x} - x - 1} > 0, \forall x \in (0, \infty).$



Watch Video Solution

11.  $\lim_{h \rightarrow 0} \frac{(e+h)^{\ln(e+h)} - e}{h}$  is-



Watch Video Solution

12. If the function  $f(x) = -4e^{\frac{1-x}{2}} + 1 + x + \frac{x^2}{2} + \frac{x^3}{3}$  and  $g(x) = f^{-1}(x)$ , then

the reciprocal of  $g' \left( \frac{-7}{6} \right)$  is \_\_\_\_\_



Watch Video Solution

13. Suppose that  $f(0) = 0$  and  $f'(0) = 2$ , and  $g(x) = f(-x + f(f(x)))$ . The value of  $g'(0)$  is equal to -



Watch Video Solution

14. Let  $f(x)$  be a polynomial with real coefficients such that  $f(x) = f'(x) \times f''(x)$ . If  $f(x)=0$  is satisfied  $x=1,2,3$  only, then the value of  $f'(1)f'(2)f'(3)$  is

 Watch Video Solution

15. A nonzero polynomial with real coefficients has the property that  $f(x) = f'(x) \cdot f''(x)$ . If  $a$  is the leading coefficient of  $f(x)$ , then the value of  $1/a$  is -

 Watch Video Solution

16. A function is represented parametrically by the equations

$$x = \frac{1+t}{t^3}; y = \frac{3}{2t^2} + \frac{2}{t} \text{ Then the value of } \frac{f(dy)}{dx} - x \left( \frac{dy}{dx} \right)^3 \text{ is } \underline{\hspace{2cm}}$$

 Watch Video Solution

17. Let  $z = (\cos x)^5$  and  $y = \sin x$ . Then the value of  $\frac{d^2 z}{dy^2} \text{ at } x = \frac{2\pi}{9}$  is \_\_\_\_\_.



Watch Video Solution

18. Let  $g(x) = \begin{cases} \frac{x^2 + x \tan x - x \tan 2x}{ax + \tan x - \tan 3x}, & x \neq 0, x = 0 \text{ if } g'(0) \text{ exists and is equal} \\ b & \text{to nonzero value } b, \text{ then } 52 \frac{b}{a} \text{ is equal to } \end{cases}$



Watch Video Solution

19. Let  $f(x) = x + \cfrac{1}{2x + \cfrac{1}{2x + \cfrac{1}{2x + \dots \dots \infty}}}$ .

Then the value of  $f(50) \cdot f'(50)$  is -



Watch Video Solution

20. Let  $F(x) = f(x)g(x)h(x)$  for all real  $x$ , where  $f(x)$ ,  $g(x)$ , and  $h(x)$  are differentiable functions. At some point  $x_0$ ,  $F'(x_0) = 21F(x_0)$ ,  $f'(x_0) = 4f(x_0)$ ,  $g'(x_0) = -7g(x_0)$ , and  $h'(x_0) = kh(x_0)$ , then the value of  $k$  is



[Watch Video Solution](#)

21. If  $y = \frac{\sqrt[3]{1+3x}\sqrt[4]{1+4x}\sqrt[5]{1+5x}}{\sqrt[7]{1+7x}\sqrt[8]{1+8x}}$ , then  $y'(0)$  is equal to -



[Watch Video Solution](#)

22. Let  $f(\theta) = \sin\left(\tan^{-1}\left(\frac{\sin\theta}{\sqrt{\cos 2\theta}}\right)\right)$ , where  $-\frac{\pi}{4} < \theta < \frac{\pi}{4}$  then the value of  $\frac{d}{d(\tan\theta)}f(\theta)$  is



[Watch Video Solution](#)

**23.** The slope of the tangent to the curve  $(y - x^5)^2 = x(1 + x^2)^2$  at the point  $(1, 3)$  is.



**Watch Video Solution**

**24.** Let  $f: R \rightarrow R$  be a differentiable function with  $f(0) = 1$  and satisfying the equation  $f(x + y) = f(x)f'(y) + f'(x)f(y)$  for all  $x, y \in R$ . Then, the value of  $(\log)_e(f(4))$  is \_\_\_\_\_



**Watch Video Solution**

**25.**  $f'(x) = \varphi'(x) = f(x)$  for all  $x$ . Also,  $f(3) = 5$  and  $f'(3) = 4$ . Then the value of  $[f(10)]^2 - [\varphi(10)]^2$  is \_\_\_\_\_



**Watch Video Solution**

26. If  $y=f(x)$  is an odd differentiable function defined on  $(-\infty, \infty)$  such that  $f'(3) = -2$  then  $f'(-3)$  is equal to-

 Watch Video Solution

27. If  $x^3 + 3x^2 - 9x + \lambda$  is of the form  $(x - \alpha)^2(x - \beta)$  then  $\lambda$  is equal to

 Watch Video Solution

28. If graph of  $y = f(x)$  is symmetrical about the point  $(5, 0)$  and  $f'(7) = 3$ , then the value of  $f'(3)$  is \_\_\_\_\_

 Watch Video Solution

29. Let  $f(x) = (x - 1)(x - 2)(x - 3)(x - n)$ ,  $n \in N$ , and  $f(n) = 5040$ . Then the value of  $n$  is \_\_\_\_\_

 Watch Video Solution

30.  $y = f(x)$ , where  $f$  satisfies the relation  $f(x + y) = 2f(x) + xy(y) + y\sqrt{f(x)}$   $\forall x, y \in R$  and  $f'(0) = 0$ . Then  $f(6)$  is equal of  $f(-3)$  is \_\_\_\_\_



Watch Video Solution

31. If function  $f$  satisfies the relation  $f(x)xf'(-x) = f(-x)xf'(x)f$  or all  $x$ , and  $f(0) = 3$ , and  $f(3) = 3$ , then the value of  $f(-3)$  is \_\_\_\_\_



Watch Video Solution

32. If  $y = \frac{a + bx^{\frac{3}{2}}}{x^{\frac{5}{4}}}$  and  $y' = 0$  at  $x = 5$ , then the value of  $\frac{a^2}{b^2}$  is \_\_\_\_\_



Watch Video Solution

33.  $\lim_{h \rightarrow 0} \frac{(e+h)^{\ln(e+h)} - e}{h}$  is-



Watch Video Solution

34. If the function  $f(x) = -4e^{\frac{1-x}{2}} + 1 + x + \frac{x^2}{2} + \frac{x^3}{3}$  and  $g(x) = f^{-1}(x)$ , then  
the reciprocal of  $g' \left( \frac{-7}{6} \right)$  is \_\_\_\_\_



Watch Video Solution

35. Suppose that  $f(0) = 0$  and  $f'(0) = 2$ , and let  $g(x) = f(-x + f(f(x)))$ . The value of  $g'(0)$  is equal to \_\_\_\_\_



Watch Video Solution

36. A nonzero polynomial with real coefficients has the property that  $f(x) = f(x) \cdot f'(x)$ . If  $a$  is the leading coefficient of  $f(x)$ , then the value of

$1/a$  is -



Watch Video Solution

37. A function is represented parametrically by the equations

$$x = \frac{1+t}{t^3}; y = \frac{3}{2t^2} + \frac{2}{t} \text{ Then the value of } \left| \frac{dy}{dx} - x \left( \frac{dy}{dx} \right)^3 \right| \text{ is } \underline{\hspace{2cm}}$$



Watch Video Solution

38. Let  $z = (\cos x)^5$  and  $y = \sin x$ . Then the value of  $2 \frac{d^2 z}{dy^2} \text{ at } x = \frac{2\pi}{9}$  is \_\_\_\_\_.



Watch Video Solution

39. Let  $g(x) = \begin{cases} x^2 + x \tan x - x \tan 2x \\ \frac{ax + \tan x - \tan 3x}{ax + \tan x - \tan 3x}, x \neq 0, x = 0 \text{ if } g'(0) \text{ exists and is equal} \end{cases}$   
to nonzero value  $b$ , then  $52 \frac{b}{a}$  is equal to \_\_\_\_\_



Watch Video Solution

40. Let  $f(x) = x + \frac{1}{2x + \frac{1}{2x + \frac{1}{2x + \infty}}}$  Compute the value of  $f(50)f'(50)$



Watch Video Solution

41. Let  $F(x) = f(x)g(x)h(x)$  for all real  $x$ , where  $f(x)$ ,  $g(x)$ , and  $h(x)$  are differentiable functions. At some point  $x_0$ ,  $F'(x_0) = 21F(x_0)$ ,  $f'(x_0)4f(x_0)$ ,  $g'(x_0) = -7g(x_0)$ , then the value of  $g'(1)$  is \_\_\_\_\_



Watch Video Solution

42. If  $y = \frac{\sqrt[3]{1+3x}\sqrt[4]{1+4x}\sqrt[5]{1+5x}}{\sqrt[7]{1+7x}\sqrt[8]{1+8x}}$ , then  $y'(0)$  is equal to -



Watch Video Solution

43. If  $f(\theta) = \sin\left(\tan^{-1}\left(\frac{\sin\theta}{\sqrt{\cos 2\theta}}\right)\right)$ , where  $-\frac{\pi}{4} < \theta < \frac{\pi}{4}$ , then the value of  $\frac{d}{d(\tan\theta)} f(\theta)$  is

 Watch Video Solution

44. The slope of the tangent to the curve  $(y - x^5)^2 = x(1 + x^2)^2$  at the point  $(1, 3)$  is.

 Watch Video Solution

45. Let  $f: R \rightarrow R$  be a differentiable function with  $f(0) = 1$  and satisfying the equation  $f(x+y) = f(x)f'(y) + f'(x)f(y)$  for all  $x, y \in R$ . Then, the value of  $(\log)_e(f(4))$  is \_\_\_\_\_

 Watch Video Solution

1. Let  $y$  be an implicit function of  $x$  defined by  $x^{2x} - 2x^x \cot y - 1 = 0$ . Then  $y'(1)$  equals: 1 b. log2 c. -log2 d. -1

A. -1

B. 1

C. log 2

D. -log 2

**Answer: A**



**Watch Video Solution**

2. Let  $f: (1, 1) \rightarrow \mathbb{R}$  be a differentiable function with  $f(0) = -1$  and  $f'(0) = 1$ . Let  $g(x) = [f(2f(x) + 2)]^2$ . Then  $g'(0) =$  (1) 4 (2) 0 (3) 2 (4) 4

A. -2

B. 4

C. -4

**Answer:****Watch Video Solution**

3.  $\frac{d^2x}{dy^2}$  equals:

(1.)  $\left(\frac{d^2y}{dx^2}\right)^{-1}$

(2) -  $\left(\frac{d^2y}{dx^2}\right)\left(\frac{dy}{dx}\right)^{-3}$

(3) -  $\left(\frac{d^2y}{dx^2}\right)^{-1}\left(\frac{dy}{dx}\right)^{-2}$

(4) -  $\left(\frac{d^2y}{dx^2}\right)^{-1}\left(\frac{dy}{dx}\right)^3$

A. -  $\left(\frac{d^2y}{dx^2}\right)\left(\frac{dy}{dx}\right)^{-3}$

B.  $\left(\frac{d^2y}{dx^2}\right)^{-1}$

C.  $-\left(\frac{d^2y}{dx^2}\right)^{-1} \left(\frac{dy}{dx}\right)^{-3}$

D.  $\left(\frac{d^2y}{dx^2}\right) \left(\frac{dy}{dx}\right)^{-2}$

**Answer:**



**Watch Video Solution**

4. If  $y = \sec(\tan^{-1}x)$ , then  $\frac{dy}{dx}$  at  $x = 1$  is equal to:  $\frac{1}{\sqrt{2}}$  (b)  $\frac{1}{2}$  (c) 1 (d)  $\sqrt{2}$

A.  $1/2$

B. 1

C.  $\sqrt{2}$

D.  $1\sqrt{2}$

**Answer:**



**Watch Video Solution**

5. If  $g$  is the inverse of a function  $f$  and  $f'(x) = \frac{1}{1+x^5}$  then  $g'(x)$  is equal to

- (1)  $1+x^5$  (2)  $5x^4$  (3)  $\frac{1}{1+\{g(x)\}^5}$  (4)  $1+\{g(x)\}^5$

A.  $1+x^5$

B.  $5x^4$

C.  $\frac{1}{1+\{g(x)\}^5}$

D.  $1+\{g(x)\}^5$

**Answer:**



**Watch Video Solution**

6. If for  $x \in \left(0, \frac{1}{4}\right)$ , the derivative of  $\tan^{-1}\left(\frac{6x\sqrt{x}}{1-9x^3}\right)$  is  $\sqrt{x}g(x)$ , then  $g(x)$

equals: (1)  $\frac{3x}{1-9x^3}$  (2)  $\frac{3}{1+9x^3}$  (3)  $\frac{9}{1+9x^3}$  (4)  $\frac{3x\sqrt{x}}{1-9x^3}$

A.  $\frac{3}{1+9x^3}$

B.  $\frac{9}{1+9x^3}$

C.  $\frac{3x\sqrt{x}}{1 - 9x^3}$

D.  $\frac{3x}{1 - 9x^3}$

**Answer:**



**Watch Video Solution**

7. Let  $y$  be an implicit function of  $x$  defined by  $x^{2x} - 2x^x \cot y - 1 = 0$ . Then  $y'$

(1) equals

A. -1

B. 1

C.  $\log 2$

D.  $-\log 2$

**Answer:**



**Watch Video Solution**

8. Let  $f: (1, 1) \rightarrow \mathbb{R}$  be a differentiable function with  $f(0) = -1$  and  $f'(0) = 1$ . Let

$g(x) = [f(2f(x) + 2)]^2$ . Then  $g'(0) =$  (1) 4 (2) 0 (3) 2 (4) -4

A. -2

B. 4

C. -4

D. 0

**Answer:**



**Watch Video Solution**

9.  $\frac{d^2x}{dy^2}$  equals: (b)  $\left(\frac{d^2y}{dx^2}\right)^{-1}$  (c)  $-\left(\frac{d^2y}{dx^2}\right)^{-1} \left(\frac{dy}{dx}\right)^{-3} \left(\frac{d^2y}{dx^2}\right) \left(\frac{dy}{dx}\right)^{-2}$  (d)

$-\left(\frac{d^2y}{dx^2}\right) \left(\frac{dy}{dx}\right)^{-3}$

A. -1

B. 1

C.  $\log 2$

D.  $-\log 2$

**Answer: A**



**Watch Video Solution**

10. If  $y = \sec(\tan^{-1}x)$ , then  $\frac{dy}{dx}$  at  $x = 1$  is equal to: (a)  $\frac{1}{\sqrt{2}}$  (b)  $\frac{1}{2}$  (c) 1 (d)  $\sqrt{2}$

A. -2

B. 4

C. -4

D. 0

**Answer:**



**Watch Video Solution**

11. If  $g$  is the inverse of a function  $f$  and  $f'(x) = \frac{1}{1+x^5}$  then  $g(x)$  is equal to

- (1)  $1+x^5$  (2)  $5x^4$  (3)  $\frac{1}{1+\{g(x)\}^5}$  (4)  $1+\{g(x)\}^5$

A.  $-\left(\frac{d^2y}{dx^2}\right)\left(\frac{dy}{dx}\right)^{-3}$

B.  $\left(\frac{d^2y}{dx^2}\right)^{-1}$

C.  $-\left(\frac{d^2y}{dx^2}\right)^{-1}\left(\frac{dy}{dx}\right)^{-3}$

D.  $\left(\frac{d^2y}{dx^2}\right)\left(\frac{dy}{dx}\right)^{-2}$

**Answer:**



**Watch Video Solution**

12. If for  $x \in \left(0, \frac{1}{4}\right)$ , the derivative of  $\tan^{-1}\left(\frac{6x\sqrt{x}}{1-9x^3}\right)$  is  $\sqrt{x}g(x)$ , then  $g(x)$

- equals: (1)  $\frac{3x}{1-9x^3}$  (2)  $\frac{3}{1+9x^3}$  (3)  $\frac{9}{1+9x^3}$  (4)  $\frac{3x\sqrt{x}}{1-9x^3}$

A.  $1/2$

B. 1

C.  $\sqrt{2}$

D.  $1\sqrt{2}$

**Answer:**



**Watch Video Solution**

13. Let  $f(x) = x \sin \pi x$ ,  $x > 0$ . Then for all natural numbers  $n$ ,  $f'(x)$  vanishes

at a unique point in the interval  $\left(n, n + \frac{1}{2}\right)$  a unique point in the interval

$\left(n + \frac{1}{2}, n + 1\right)$  a unique point in the interval  $(n, n + 1)$  two points in the

interval  $(n, n + 1)$

A.  $1 + x^5$

B.  $5x^4$

C.  $\frac{1}{1 + \{g(x)\}^5}$

D.  $1 + \{g(x)\}^5$

**Answer:**



**Watch Video Solution**

14. Let  $f: (0, \infty) \rightarrow R$  be a differentiable function such that  $f'(x) = 2 - \frac{f(x)}{x}$

for all  $x \in (0, \infty)$  and  $f(1) = 1$ , then  $f(x)$  is

A.  $\frac{3}{1 + 9x^3}$

B.  $\frac{9}{1 + 9x^3}$

C.  $\frac{3x\sqrt{x}}{1 - 9x^3}$

D.  $\frac{3x}{1 - 9x^3}$

**Answer:**



**Watch Video Solution**