

MATHS

BOOKS - SAI MATHS (TELUGU ENGLISH)

MEASURES OF DISPERSION AND PROBABILITY

Problems

1. The probability of a coin showing head is p and then 100 such coins are tossed. If the probability of 50 coins showing head is same as the probability of 51 coins showing head then p equals

A. $\frac{1}{2}$

B. $\frac{49}{100}$

C. $\frac{51}{101}$

D. $\frac{50}{101}$

Answer: C



Watch Video Solution

2. X is a binomial variate with parameters $n = 6$ and p . If

$4P(X = 4) = P(X = 2)$, then p is

A. $\frac{1}{2}$

B. $\frac{1}{3}$

C. $\frac{1}{4}$

D. $\frac{1}{6}$

Answer: B



Watch Video Solution

3. In a certain college, 4% of men and 1% of women are taller than 1.8m. Also, 60% of students are women. If a student selected at random is found to be taller than 1.8 m, then the probability that the student being a woman is

A. $\frac{3}{11}$

B. $\frac{5}{11}$

C. $\frac{6}{11}$

D. $\frac{8}{11}$

Answer: A



Watch Video Solution

4. If A and B are two events such that $P(A/B) = 0.6$, $P(B/A) = 0.3$, $P(A) = 0.1$ then, $P(\bar{A} \cap \bar{B}) =$

A. 0.88

B. 0.12

C. 0.6

D. 0.4

Answer: A



[View Text Solution](#)

5. If A and B are events such that $P(A \cup B) = \frac{5}{6}$, $P(\bar{A}) = \frac{1}{4}$ and $P(B) = \frac{1}{3}$, then A and B

are

- A. Mutually exclusive
- B. independent
- C. Exhaustive events
- D. Exhaustive and independent

Answer: B



[Watch Video Solution](#)

6. Two teams A and B have the same mean and their coefficients of variance are 4, 2 respectively. If σ_A, σ_B are the standard deviations of teams A, B respectively then the relation between them is

A. $\sigma_A = \sigma_B$

B. $\sigma_B = 2\sigma_A$

C. $\sigma_A = 2\sigma_B$

D. $\sigma_B = 4\sigma_A$

Answer: C



Watch Video Solution

7. In a data the number i is repeated i times for $i = 1, 2, \dots, n$.

Then the mean of the data is

A. $\frac{2n + 1}{6}$

B. $\frac{2n + 1}{4}$

C. $\frac{2n + 1}{3}$

D. $\frac{2n + 1}{2}$

Answer: C



View Text Solution

8. The arithmetic mean of the observations 10, 8, 5, a, b is 6 and their variance is 6.8. Then $ab =$

A. 6

B. 4

C. 3

D. 12

Answer: D



Watch Video Solution

9. If the median of the data 6, 7, $x-2$, x , 18, 21 written in ascending order is 16, then the variance of that data is

A. $30\frac{1}{5}$

B. $31\frac{1}{3}$

C. $32\frac{1}{2}$

D. $33\frac{1}{3}$

Answer: B



[Watch Video Solution](#)

10. Two persons A and B are rolling die on the condition that the person who gets 3 will win the game. If A starts the game,

then find the probabilities of A and B respectively to win the game.

A. $\frac{6}{11}, \frac{5}{11}$

B. $\frac{5}{11}, \frac{6}{11}$

C. $\frac{8}{11}, \frac{3}{11}$

D. $\frac{3}{11}, \frac{8}{11}$

Answer: A



[Watch Video Solution](#)

11. The letters of the word 'QUESTION' are arranged in a row at random. The probability that there are exactly two letters between Q and S is

A. $\frac{1}{14}$

B. $\frac{5}{7}$

C. $\frac{1}{7}$

D. $\frac{5}{28}$

Answer: D



Watch Video Solution

12. If $\frac{1 + 3P}{3}$, $\frac{1 - 2P}{2}$ are probabilities of two mutually exclusive events, then P lies in the interval.

A. $\left[-\frac{1}{3}, \frac{1}{2} \right]$

B. $\left[-\frac{1}{2}, \frac{1}{2} \right]$

C. $\left[-\frac{1}{3}, \frac{2}{3} \right]$

D. $\left[-\frac{1}{3}, \frac{2}{3}\right]$

Answer: A



Watch Video Solution

13. The probability that an event does not happen in one trial is 0.8 The probability that the event happens atmost once in three trials is

A. 0.896

B. 0.791

C. 0.642

D. 0.592

Answer: A



[Watch Video Solution](#)

14. If the mean and variance of a binomial variate X are 8 and 4 respectively, then $P(X < 3)$ equals to

A. $\frac{265}{2^{15}}$

B. $\frac{137}{2^{14}}$

C. $\frac{137}{2^{16}}$

D. $\frac{265}{2^{16}}$

Answer: C



[View Text Solution](#)

15. A candidate take three tests in succession and the probability of passing the first test is p . The probability of passing each succeeding test p or $\frac{p}{2}$ according as he passes or fails in the preceding one. The candidate is selected if he passes at least two tests. The probability that the candidate is selected is

A. $p^2(2 - p)$

B. $p(2-p)$

C. $p + p^2 + p^3$

D. $p^2(1 - p)$

Answer: C



Watch Video Solution

16. A six-faced unbiased die is thrown twice and the sum of the numbers appearing on the upper face is observed to be 7 . The probability that the number 3 has appeared at least once , is

A. $\frac{43835}{100000}$

B. $\frac{1}{2}$

C. $\frac{1}{3}$

D. $\frac{1}{4}$

Answer: C



[Watch Video Solution](#)

17. If A,B,C are mutually exclusive and exhaustive events of a random experiment such that $P(B) =$

$\frac{3}{2}P(A)$ and $P(C) = \frac{1}{2}P(B)$ then $P(A \cup C) =$

A. $\frac{10}{13}$

B. $\frac{3}{13}$

C. $\frac{6}{13}$

D. $\frac{7}{13}$

Answer: D



[Watch Video Solution](#)

18. If x_1, x_2, \dots, x_n are n observations such that $\sum x_i^2 = 400$ and $\sum x_i = 80$ then the least value of n is

A. 18

B. 12

C. 15

D. 16

Answer: D



Watch Video Solution

19. The mean of four observations is 3. If the sum of the squares of these observations is 48 then their standard deviation is

A. $\sqrt{7}$

B. $\sqrt{2}$

C. $\sqrt{3}$

D. $\sqrt{5}$

Answer: C



Watch Video Solution

20. Two numbers are chosen at random from (1,2,3,4,5,6,7,8) at a time. The probability that smaller of the two numbers is less than 4 is

A. $\frac{7}{14}$

B. $\frac{8}{14}$

C. $\frac{9}{14}$

D. $\frac{10}{14}$

Answer: C



Watch Video Solution

21. Two fair dice are rolled. The probability of the sum of digits on their faces to be greater than or equal to 10 is

A. $\frac{1}{5}$

B. $\frac{1}{4}$

C. $\frac{1}{8}$

D. $\frac{1}{6}$

Answer: D



[Watch Video Solution](#)

22. A bag contains $2n + 1$ coins . It is known that n of these coins have a head on both sides, whereas the remaining $n + 1$ coins are fair . A coin is picked up at random from the beg and

tossed . If the probability that the toss results in a head is $\frac{31}{42}$, then n is equal to

A. 10

B. 11

C. 12

D. 13

Answer: A



[View Text Solution](#)

23. The random variable takes the values 1,2,3,.....m. If

$P(X = n) = \frac{1}{m}$ to each n, then the variance of X is

A. $\frac{(m + 1)(2m + 1)}{6}$

B. $\frac{m^2 - 1}{12}$

C. $\frac{m + 1}{2}$

D. $\frac{m^2 + 1}{12}$

Answer: B



Watch Video Solution

24. If X is a Poisson variate and $P(X=1)=2P(X=2)$ then $P(X=3)=$

A. $\frac{e^1}{6}$

B. $\frac{e^2}{2}$

C. $\frac{e^{-1}}{2}$

D. $\frac{e^{-1}}{3}$

Answer: A



[Watch Video Solution](#)

25. If X is a random Poisson variate such that $\alpha = P(X = 1) = P(X = 2)$ then $P(X=4)=$

A. 2α

B. $\frac{\alpha}{3}$

C. α^{-2}

D. αe^2

Answer: B



[Watch Video Solution](#)

26. X follows a binomial distribution with parameters n and p

where $0 < p < 1$.

If $\frac{P(X = r)}{P(X = n - r)}$ is independent of n and r then $p =$

A. $\frac{1}{2}$

B. $\frac{1}{3}$

C. $\frac{1}{4}$

D. $\frac{1}{8}$

Answer: A



[Watch Video Solution](#)

27. In an entrance test there are multiple choice questions ,
then the probability that he was guessing is

A. $\frac{37}{40}$

B. $\frac{1}{37}$

C. $\frac{36}{37}$

D. $\frac{1}{9}$

Answer: B



[View Text Solution](#)

28. There are four machines and it is known that exactly two of them are faulty . They are tested one by one , in a random order till both the faulty machines are identified . Then , the probability that only two tests are needed is ,

A. $\frac{1}{3}$

B. $\frac{1}{6}$

C. $\frac{1}{2}$

D. $\frac{1}{4}$

Answer: A



[View Text Solution](#)

29. A fair coin is tossed 100 times . The probability of getting tails an odd number of times is

A. $\frac{1}{2}$

B. $\frac{1}{4}$

C. $\frac{1}{8}$

D. $\frac{3}{8}$

Answer: A



[View Text Solution](#)

30. A class has fifteen boys and five girls. Suppose three students are selected at random from the class. The probability that there are two boys and one girl is

A. $\frac{35}{76}$

B. $\frac{35}{38}$

C. $\frac{7}{76}$

D. $\frac{35}{72}$

Answer: A



[Watch Video Solution](#)

31. Seven white balls and three black balls are randomly arranged in a row. The probability that no two black balls are placed adjacently is

A. $\frac{1}{2}$

B. $\frac{7}{15}$

C. $\frac{2}{15}$

D. $\frac{1}{3}$

Answer: B



[Watch Video Solution](#)

32. Let A and B be events in a sample space S such that $P(A)=0.5$, $P(B)=0.4$ and $P(A \cup B) = 0.6$. Observe the

following lists :

List I

- i) $P(A \cap B)$
- ii) $P(A \cap \bar{B})$
- iii) $P(\bar{A} \cap B)$
- iv) $P(\bar{A} \cap \bar{B})$

List II

- a) 0.4
- b) 0.2
- c) 0.3
- d) 0.1

- A.

(i)	(ii)	(iii)	(iv)
(1)	(2)	(3)	(4)
- B.

(i)	(ii)	(iii)	(iv)
(3)	(2)	(4)	(1)
- C.

(i)	(ii)	(iii)	(iv)
(3)	(2)	(1)	(4)
- D.

(i)	(ii)	(iii)	(iv)
(3)	(1)	(2)	(4)

Answer: B



Watch Video Solution

33. The probability that an individual suffers a bad reaction from an injection is 0.001. The probability that out of 2000

individulas exactly three will suffer bad reaction is

A. $\frac{1}{e^2}$

B. $\frac{2}{3e^2}$

C. $\frac{8}{3e^2}$

D. $\frac{4}{3e^2}$

Answer: D



[Watch Video Solution](#)

34. A urn A contains 3 white and 5 black balls. Another urn B contains 6 white and 8 black balls. A ball is picked from A at random and then transferred to B. Then a ball is picked at random from B. The probability that it is a white ball is

A. $\frac{14}{40}$

B. $\frac{15}{40}$

C. $\frac{16}{40}$

D. $\frac{17}{40}$

Answer: D



Watch Video Solution

35. If $A(i=1,2,3,\dots,n)$ are n independent events with $P(A) = \frac{1}{1+i}$

for each i , then the probability that none of A , occur is :

A. $\frac{n-1}{n+1}$

B. $\frac{n}{n+1}$

C. $\frac{n}{n+2}$

D. $\frac{1}{n+1}$

Answer: D

 [Watch Video Solution](#)

36. Suppose that A and B are two independent events such that $P(A \cap B) = \frac{3}{25}$ and $P(A) = \frac{8}{25}$ Then $P(B) =$

A. $\frac{11}{25}$

B. $\frac{3}{11}$

C. $\frac{1}{11}$

D. $\frac{9}{11}$

Answer: A

 [Watch Video Solution](#)

37. Suppose that a random variable X follows Poisson distribution . If $P(X=1)=P(X=2)$ then $P(X=5)=$

A. $\frac{2}{3}e^{-2}$

B. $\frac{3}{4}e^{-2}$

C. $\frac{4}{15}e^{-2}$

D. $\frac{7}{8}e^{-2}$

Answer: C



Watch Video Solution

38. If the mean and variance of a binomial variable X are 2 and 1 respectively, then $P(X \geq 1) =$

A. $\frac{2}{3}$

B. $\frac{15}{16}$

C. $\frac{7}{8}$

D. $\frac{4}{5}$

Answer: B



Watch Video Solution

39. If A and B are events of a random experiment such that

$$P(A \cup B) = \frac{4}{5}, P(\bar{A} \cup \bar{B}) = P(\bar{A} \cap \bar{B}) = \frac{7}{10} \text{ and } P(B) = \frac{2}{5}$$

, then P(A) equal

A. $\frac{9}{10}$

B. $\frac{8}{10}$

C. $\frac{7}{10}$

D. $\frac{3}{5}$

Answer: D



Watch Video Solution

40. The probability of choosing randomly a number c from the set $\{1,2,3,\dots,9\}$ such that the quadratic equation

$x^2 + 4x + c = 0$ has real roots is

A. $\frac{1}{9}$

B. $\frac{2}{9}$

C. $\frac{3}{9}$

D. $\frac{4}{9}$

Answer: D



View Text Solution

41. Suppose that E_1 and E_2 are two events of a random experiment such that

$$P(E_1) = \frac{1}{4}, P(E_2 | E_1) = \frac{1}{2} \text{ and } P(E_1 | E_2) = \frac{1}{4}.$$

Observe the lists given below :

List I

- A) $P(E_2)$
- B) $P(E_1 \cup E_2)$
- C) $P(\overline{E_1} | \overline{E_2})$
- D) $P(E_1 | E_2)$

List II

- i) $1/4$
- ii) $5/8$
- iii) $1/8$
- iv) $1/2$
- v) $3/8$
- vi) $3/4$

The correct matching of the list I from the list II is :

- A. (A) (B) (C) (D)
(a) (ii) (iii) (vi) (i)

- B. (A) (B) (C) (D)
 (a) (iv) (v) (vi) (i)
- C. (A) (B) (C) (D)
 (a) (iv) (ii) (vi) (i)
- D. (A) (B) (C) (D)
 (a) (i) (ii) (iii) (i)

Answer: C



Watch Video Solution

42. X is a binomial variate with parameters $n = 6$ and p . If $4P(X = 4) = P(X = 2)$, then p is

- A. $\frac{1}{3}$
- B. $\frac{1}{2}$
- C. $\frac{2}{3}$
- D. $\frac{3}{4}$

Answer: A



Watch Video Solution

43. If A and B are two independent events of a random experiment such that $P(A \cap B) = \frac{1}{6}$ and $P(\bar{A} \cap \bar{B}) = \frac{1}{3}$, then $P(A) =$

A. $\frac{1}{4}$

B. $\frac{1}{3}$

C. $\frac{1}{2}$

D. $\frac{2}{3}$

Answer: A



Watch Video Solution

44. Let S be the sample space of the random experiment of throwing simultaneously two unbiased dice with six faces (numbered 1 to 6) and let $E_k = \{(a,b) \in S : ab = k\}$ for $k \geq 1$

A. $p_1 < p_{30} < p_4 < p_6$

B. $p_{36} < p_6 < p_2 < p_4$

C. $p_1 < p_{11} < p_4 < p_6$

D. $p_{36} < p_{11} < p_6 < p_4$

Answer: A:C



Watch Video Solution

45. For $L=1,2,3$ the box B_k contains k red balls and $(k+1)$ white balls. Let $P(B_1) = \frac{1}{2}$, $P(B_2) = \frac{1}{3}$, $P(B_3) = \frac{1}{6}$. A box is

selected at random and a ball is drawn from it . If a red ball is drawn, then the probability that it has come from box B_2 is

A. $\frac{35}{78}$

B. $\frac{14}{39}$

C. $\frac{10}{13}$


D. $\frac{12}{13}$

Answer: A



[Watch Video Solution](#)

46. The distribution of a random variable X is given below



The value of k is

A. $\frac{1}{10}$

B. $\frac{2}{10}$

C. $\frac{3}{10}$

D. $\frac{7}{10}$

Answer: B



[View Text Solution](#)

47. If X is a poisson distribution such that

$P(X=1)=P(X=2)$ then, $P(X=4)=$

A. $\frac{1}{2e^2}$

B. $\frac{1}{3e^2}$

C. $\frac{2}{3e^2}$

D. $\frac{1}{e^2}$

Answer: C



Watch Video Solution

48. Four numbers are chosen at random from $(1,2,3,\dots,40)$. The probability that they are not consecutive is

A. $\frac{1}{2470}$

B. $\frac{4}{7069}$

C. $\frac{2469}{2470}$

D. $\frac{7965}{7969}$

Answer: C



Watch Video Solution

49. If A and B are mutually exclusive events with $P(B) \neq 1$ then $P(A | \bar{B})$ is equal to {Here \bar{B} is the complement of the event B)

A. $\frac{1}{P(B)}$

B. $\frac{1}{1 - P(B)}$

C. $\frac{P(A)}{P(B)}$

D. $\frac{P(A)}{1 - P(B)}$

Answer: D



Watch Video Solution

50. A bag contains 6 white and 4 black balls. Two balls are drawn at random. The probability that they are of the same colour is

A. $\frac{1}{15}$

B. $\frac{2}{5}$

C. $\frac{4}{15}$

D. $\frac{7}{15}$

Answer: D



[Watch Video Solution](#)

51. The mean and standard deviation of a binomial variate X are 4 and $\sqrt{3}$ respectively. Then $P(X = \geq 1) =$

A. $1 - \left(\frac{1}{4}\right)^{16}$

B. $1 - \left(\frac{3}{4}\right)^{16}$

C. $1 - \left(\frac{2}{3}\right)^{16}$

D. $1 - \left(\frac{1}{3}\right)^{16}$

Answer: B



Watch Video Solution

52. In the random experiment of tossing two unbiased dice, let E be the event of getting the sum 8 and F be the event of getting even numbers on both the dice. Then ,

Statement I Statement II

$$P(E) = \frac{7}{36} \quad P(F) = \frac{1}{3}$$

Which of the following is a correct statement ?

A. Both I and II are true

B. Neither I nor II is true

C. I is true , II is false

D. I is false , II is true

Answer: B



[Watch Video Solution](#)

53. Seven balls are drawn simultaneously from a bag containing 5 white and 6 green balls. The probability of drawing 3 white and 4 green balls is

A. $\frac{7}{11c_7}$

B. $\frac{5c_3 + 6c_4}{11c_7}$

C. $\frac{{}^5C_2 {}^6C_2}{{}^{11}C_7}$

D. $\frac{{}^6C_3 {}^5C_4}{{}^{11}C_7}$

Answer: C



Watch Video Solution

54. In a book of 500 pages , it is found that there are 250 typing errors . Assume that Poisson law holds for the number of errors per page . Then, the probability that a random sample of 2 pages will contains no error is

A. $e^{-0.3}$

B. $e^{-0.5}$

C. e^{-1}

D. e^{-2}

Answer: C



Watch Video Solution

55. If A and B are two independent events such that

$P(B) = \frac{2}{7}$, $P(A \cup B^c) = 0.8$, then $P(A)$ is equal to

A. 0.1

B. 0.2

C. 0.3

D. 0.4

Answer: C



Watch Video Solution

56. A number n is chosen at random from $(1,2,3,4,\dots,1000)$. The probability that n is a number that leaves remainder 1 when divided by 7 is

A. $\frac{71}{500}$

B. $\frac{143}{1000}$

C. $\frac{72}{500}$

D. $\frac{71}{1000}$

Answer: B



Watch Video Solution

57. A coin and six faced die, both unbiased are thrown simultaneously . The probability of getting a head on the coin and an odd number on the die is

A. $\frac{1}{2}$

B. $\frac{3}{4}$

C. $\frac{1}{4}$

D. $\frac{2}{3}$

Answer: C



[Watch Video Solution](#)

58. A number n is chosen at random from

$S = \{1,2,3,\dots, 50\}$ Let

$$A = \left\{ n \in S : n + \frac{50}{n} > 27 \right\}$$

$B = \{n \in S : n \text{ is a prime}\}$ and

$C = \{n \in S : n \text{ is a square}\}$. Then , correct order of their probabilities is

A. $P(A) < P(B) < P(C)$

B. $P(A) > P(B) > P(C)$

C. $P(B) < P(A) < P(C)$

D. $P(A) > P(C) > P(B)$

Answer: C



[View Text Solution](#)

59. Box A contains 2 black and 3 red balls , while box B contains 3 black and 4 red balls . Out of these two boxes one

is selected at random and the probability of choosing box A is double that of box B . If a red ball is drawn from the selected box , then the probability that it has come from box B, is

A. $\frac{21}{41}$

B. $\frac{10}{31}$

C. $\frac{12}{31}$

D. $\frac{13}{41}$

Answer: B



[View Text Solution](#)

60. If the range of random variable X is {0,1,2,3,4....} with

$$P(X = k) = \frac{(k + 1)a}{3^k} \text{ for } k \geq 0, \text{ then } a \text{ is equal to}$$

A. $\frac{2}{3}$

B. $\frac{4}{9}$

C. $\frac{8}{27}$

D. $\frac{16}{81}$

Answer: B



[View Text Solution](#)

61. For a Binomial variate X with $n = 6$, if $P(X = 2) = 9 P(X = 4)$, then its variance is ,

A. $\frac{8}{9}$

B. $\frac{1}{4}$

C. $\frac{9}{8}$

D. 4

Answer: C



[View Text Solution](#)

62. Suppose E and F are two events of a random experiment
If the probability of occurrence of E is $\frac{1}{5}$ and the
probability of occurrence of F given E is $\frac{1}{10}$, then the
probability of non-occurrence of at least one of the events E
and F, is

A. $\frac{1}{18}$

B. $\frac{1}{2}$

C. $\frac{49}{50}$

D. $\frac{1}{50}$

Answer: C



[View Text Solution](#)

63. Six faces of an unbiased die are numbered with 2,3,5,7,11 and 13 . If two such dice are thrown , then the probability that the sum on the uppermost faces of the dice is an odd number , is

A. $\frac{5}{18}$

B. $\frac{5}{36}$

C. $\frac{13}{18}$

D. $\frac{25}{36}$

Answer: A



[View Text Solution](#)

64. If X is a Poisson variate with $P(X = 0) = 0.8$, then the variance of X is .

A. $\log_e 20$

B. $\log_{10} 20$

C. $\log_e 5/4$

D. 0

Answer: C



[View Text Solution](#)

65. An unbiased coin is tossed to get 2 points for turning up a head and one point for the tail . If three unbiased conis are

tossed simultaneously , then the probability of getting a total of odd number of points is ,

A. $\frac{1}{2}$

B. $\frac{1}{4}$

C. $\frac{1}{8}$

D. $\frac{3}{8}$

Answer: A



[View Text Solution](#)

66. If $P(A \cup B) = 0.8$, $P(A \cap B) = 0.3$ then

$$P(\bar{A}) + P(\bar{B}) =$$

A. 0.3

B. 0.5

C. 0.8

D. 0.9

Answer: D



Watch Video Solution

67. An unbiased coin is tossed n times. If the probability of getting atleast one head is greater than 0.8, then the least value of n is

A. 2

B. 3

C. 4

D. 5

Answer: B



Watch Video Solution

68. A bag X contains 2 white and 3 black balls and another bag Y contains 4 white 2 black balls . One bag is selected at random and a ball is drawn from it . Then , the probability for the ball chosen to be white is ,

A. $\frac{2}{15}$

B. $\frac{7}{15}$

C. $\frac{8}{15}$

D. $\frac{14}{15}$

Answer: C



[View Text Solution](#)

69. For a Poisson variate X , if $P(X = 2) = 3 P(X = 3)$ then the mean of X is

A. 1

B. $\frac{1}{2}$

C. $\frac{1}{3}$

D. $\frac{1}{4}$

Answer: A



[View Text Solution](#)

70. A random variate X takes the values 0,1,2,3 and its mean is 1.3 . If $P(X = 3) = 2P(X = 1)$ and $P(X = 2) = 0.3$, then $P(X = 0)$ is equal to

A. 0.1

B. 0.2

C. 0.3

D. 0.4

Answer: D



[View Text Solution](#)

71. A bag contains 5 black balls 4 white balls and 3 red balls . If a ball is selected at random , the probability that it is a black or a red ball, is

A. $\frac{1}{3}$

B. $\frac{1}{4}$

C. $\frac{5}{12}$

D. $\frac{2}{3}$

Answer: D



View Text Solution

72. The probability of getting qualified in IIT JEE EAMCET by student are respectively $\frac{1}{5}$ and $\frac{3}{5}$. The probability that the student gets qualified for at least one of these tests, is

A. $\frac{3}{25}$

B. $\frac{8}{25}$

C. $\frac{17}{25}$

D. $\frac{22}{25}$

Answer: C



[View Text Solution](#)

73. One die and a coin (both unbiased) are tossed simultaneously . The probability of getting 5 on the top of the die and tail on the coin , is

A. $\frac{1}{2}$

B. $\frac{1}{12}$

C. $\frac{1}{6}$

D. $\frac{1}{8}$

Answer: B



[View Text Solution](#)

74. In a Binomial distribution, the probability of getting success is $\frac{1}{4}$ and the standard deviation is 3 . Then its mean is

A. 6

B. 8

C. 10

D. 12

Answer: D



[View Text Solution](#)

75. If the mean of a Poisson distribution is $\frac{1}{2}$, then the ratio of $P(X = 3)$ to $P(X = 2)$ is

A. 1 : 2

B. 1 : 4

C. 1 : 6

D. 1 : 8

Answer: C



View Text Solution

76. A random variable X takes the values 0,1 and 2. If $P(X = 1) = P(X=2)$ and $P(X = 0) = 0.4$, then the mean of random variable X is

A. 0.2

B. 0.7

C. 0.5

D. 0.9

Answer: D



[View Text Solution](#)

77. In a competition A,B,C are participating . The probability that A wins is twice that of B , the probability that B wins is twice that of C, then probability that A losses is

A. $\frac{1}{7}$

B. $\frac{2}{7}$

C. $\frac{4}{7}$

D. $\frac{3}{7}$

Answer: D



View Text Solution

78. The probability that a number selected at random from the set of numbers $\{1, 2, 3, \dots, 100\}$ is a cube, is

A. $\frac{1}{25}$

B. $\frac{2}{25}$

C. $\frac{3}{25}$

D. $\frac{4}{25}$

Answer: A



[View Text Solution](#)

79. The events A and B have probabilities 0.25 and respectively . The probability that both A and B simultaneously is 0.14 , then the probability that n A nor B occurs, is

A. 0,39

B. 0,29

C. 0.11

D. 0.25

Answer: A



[View Text Solution](#)

80. Two dice are rolled simultaneously . The probability that the sum of the two numbers on the dice is a prime number , is

A. $\frac{5}{12}$

B. $\frac{7}{12}$

C. $\frac{9}{14}$

D. None of these

Answer: A



[View Text Solution](#)

81. Find the Binomial probability distribution whose mean is 3
red variance is 2 .

A. $\left(\frac{2}{3} + \frac{1}{3}\right)^9$

B. $\left(\frac{5}{3} + \frac{2}{3}\right)^9$

C. $\left(\frac{3}{2} + \frac{1}{2}\right)^9$

D. None of these

Answer: A



[View Text Solution](#)

82. For a Binomial variate X , if $n = 4$ and $P(X = 4) = 6 P(X = 2)$,

then the value of p is

A. $\frac{3}{7}$

B. $\frac{4}{7}$

C. $\frac{6}{7}$

D. $\frac{5}{7}$

Answer: C



View Text Solution