



MATHS

BOOKS - SAI MATHS (TELUGU ENGLISH)

MOCK TEST 1

Mathematics

1. If $f: R$ to R is defined by $f(x) = x^2 - 6x + 4$ then , $f(3x + 4) =$

A. $3x^2 + 2x + 2$

B. $9x^2 + 6x - 4$

C. $2x + 2$

D. $x^2 + 6x + 9$

Answer: B



Watch Video Solution

2. The domain of $\frac{\sqrt{7+x} + \sqrt{7-x}}{x}$ is

A. $[-7, 0) \cup (0, 7]$

B. $\mathbb{R} - \{0\}$

C. $(-7, 7)$

D. $[-7, 7]$

Answer: A



Watch Video Solution

3. $4^n - 3n + k$ is divisible by 9 for $n \in \mathbb{N}$. Then the numerically least -ve integral value of k is ,

A. -5

B. -1

C. -3

D. 0

Answer: B

 [View Text Solution](#)

4. If $A = \begin{bmatrix} 1 & 4 \\ 2 & 8 \end{bmatrix}$, $B = \begin{bmatrix} 3 & 2 \\ 1 & 3 \end{bmatrix}$ then $AB =$

A. $\begin{bmatrix} 9 & 16 \\ 13 & 20 \end{bmatrix}$

B. $\begin{bmatrix} 7 & 14 \\ 14 & 28 \end{bmatrix}$

C. $\begin{bmatrix} 10 & 14 \\ 14 & 24 \end{bmatrix}$

D. $\begin{bmatrix} 13 & 14 \\ 15 & 16 \end{bmatrix}$

Answer: B

 [Watch Video Solution](#)

5. The inverse of the matrix $\begin{bmatrix} 1 & 1 & 1 \\ 0 & 2 & 1 \\ 1 & 0 & 1 \end{bmatrix}$ is,

A. $\begin{bmatrix} -2 & -1 & 1 \\ 1 & 0 & 1 \\ 2 & -1 & -2 \end{bmatrix}$

B. $\begin{bmatrix} -2 & 1 & 1 \\ -1 & 0 & 1 \\ 2 & -1 & -2 \end{bmatrix}$

C. $\begin{bmatrix} 2 & 1 & 1 \\ -1 & 0 & 1 \\ 2 & -1 & -2 \end{bmatrix}$

D. $\begin{bmatrix} 2 & -1 & -1 \\ 1 & 0 & -1 \\ -2 & 1 & 2 \end{bmatrix}$

Answer: D



[Watch Video Solution](#)

6. The equations $x + y + z = 0$, $x + 2y - 4z = 0$, $2x - y - z = 0$ have

A. no solution

B. infinity many solution

C. unique solution

D. none

Answer: C



Watch Video Solution

7. If a, b, c are the position vectors of the vertices of a triangle ABC , then the position vector of its centroid is

A. $\frac{a + b + c}{3}$

B. $\frac{a - b + c}{3}$

C. $-\frac{a + b + c}{3}$

D. $\frac{a - b - c}{3}$

Answer: A



View Text Solution

8. If the position vector of P,Q are $i + j + k$ and $i + 4j + 7k$ respectively then the position vector of the point which divides \overline{PQ} in the ratio 2 : 1 is

A. $2i + j + k$

B. $3i + 2j + 4k$

C. $\frac{i + j + k}{3}$

D. $i + 3j + 5k$

Answer: D



Watch Video Solution

9. The angle between a diagonal of a cube and the diagonal of a face of the cube is

A. $\sin^{-1}\left(\frac{1}{3}\right)$

B. $\cos^{-1}\left(\frac{1}{3}\right)$

C. $\cos^{-1}\left(\sqrt{\frac{2}{3}}\right)$

D. $\cos^{-1}\left(\frac{2}{3}\right)$

Answer: C



Watch Video Solution

10. $(a \times b)^2 + (a \cdot b)^2 =$

A. $(a + b)^2$

B. a^2b^2

C. $(a - b)^2$

D. ab

Answer: B



Watch Video Solution

11. If a and b are vectors, satisfying $|a| = |b| = 2$ and $(a, b) = 38$, then the area of triangle constructed with the vector $a - b$ and $a + b$ is

- A. 2 sq. units
- B. 3 sq. units
- C. 1 sq. units
- D. 4 sq. units

Answer: A



[View Text Solution](#)

12. The volume of the parallelepiped whose edges are given by $i + 2j + 3k, 2i + 3j + 2k, 2i + 3j + k$ is

- A. 8
- B. 6
- C. 7

D. 12

Answer: B



Watch Video Solution

13. The value of $\cos 22\frac{1^\circ}{2}$ is

A. $\sqrt{2} + 1$

B. $\frac{\sqrt{3} + 1}{2\sqrt{2}}$

C. $\sqrt{\frac{\sqrt{2} + 1}{2\sqrt{2}}}$

D. $\frac{\sqrt{\sqrt{2} - 1}}{2\sqrt{2}}$

Answer: C



Watch Video Solution

14. The value of $\frac{\tan 70^\circ - \tan 20^\circ}{\tan 50^\circ} =$

A. 1

B. -1

C. -2

D. 2

Answer: D



Watch Video Solution

15. Solution of $4 \cos^2 x + 2 \sin^2 x = 3$ is

A. $n\pi \pm \frac{\pi}{4}$

B. $n\pi \pm \frac{\pi}{3}$

C. $n\pi \pm \frac{\pi}{6}$

D. $n\pi \pm \frac{\pi}{2}$

Answer: A



View Text Solution

16. If $a_1, a_2, a_3, \dots, a_n$ is an A.P. with common difference d , then

$$\tan \left[\tan^{-1} \left(\frac{d}{1 + a_1 a_2} \right) + \tan^{-1} \left(\frac{d}{1 + a_2 a_3} \right) + \dots \tan^{-1} \left(\frac{d}{1 + a_{n-1} a_n} \right) \right] =$$

A. $\frac{nd}{1 + x_1 x_n}$

B. $\frac{(n-1)d}{x_1 + x_n}$

C. $\frac{x_n - x_1}{x_n + x_1}$

D. $\frac{(n-1)d}{1 + x_1 x_n}$

Answer: D



Watch Video Solution

17. If $\sin h^{-1}(4) + \sin h^{-1}(5) = \beta$, then $\sin h(\beta) =$

A. $5\sqrt{10} + 4\sqrt{12}$

B. $4\sqrt{26} + 5\sqrt{17}$

C. $4\sqrt{20} + 5\sqrt{15}$

D. $4\sqrt{21} + 5\sqrt{11}$

Answer: B



[View Text Solution](#)

18. In a right angled triangle ABC , $r : R : r_1 =$

A. $1 : 2 : 3$

B. $5 : 12 : 10$

C. $2 : 5 : 12$

D. $7 : 6 : 8$

Answer: C



[View Text Solution](#)

19. In Δabc , if $b = \sqrt{2} + 1$, $A = 60^\circ$, $C = 30^\circ$, then $c =$

A. $\frac{\sqrt{2} - 1}{2}$

B. $\sqrt{2} + 1$

C. $-\frac{(\sqrt{2} + 1)}{2}$

D. $\frac{\sqrt{2} + 1}{2}$

Answer: D



Watch Video Solution

20. A flag staff on top of the tower 30mt of height subtends an angle of $\tan^{-1}\left(\frac{1}{3}\right)$ at a point on the ground 50mt from the foot of the tower.

The height of the flag staff is



Watch Video Solution

21. If $K \neq 1$ then, $\left| \frac{z - z_1}{z - z_2} \right| = K$ is α

A. Straight line

B. Ellipse

C. Circle

D. Parabola

Answer: C



[View Text Solution](#)

22. If, $\left(\frac{2+2i}{2-2i}\right)^3 + \left(\frac{2-2i}{2+2i}\right)^3 = \alpha + ib$, then α and b are

A. 0, 0

B. 0, 1

C. 1, 1

D. ,1,2`

Answer: A



[Watch Video Solution](#)

23. If $\cos \alpha + \cos \beta + \cos \gamma = 0 = \sin \alpha + \sin \beta + \sin \gamma$ then
 $\cos(2\alpha - \beta - \gamma) + \cos(2\beta - \gamma - \alpha) + \cos(2\gamma - \alpha - \beta) =$

A. 3

B. 2

C. 1

D. 0

Answer: A



[Watch Video Solution](#)

24. If ω is a complex cube root of unity, then $\cos \left[(\omega^7 + \omega^{11})\pi + \frac{\pi}{3} \right]$

A. 3

B. 2

C. 1

D. 0

Answer: B



Watch Video Solution

25. If $x^8 + 1 = 0$, then $X =$

A. $\text{cis} \left[(2k + 1) \frac{\pi}{8} \right], K = 0, 1, 2, 3, 4, 5, 6, 7$

B. $\text{cis} \left[(2k - 1) \frac{\pi}{8} \right], K = 0, 1, 2, 3, 4, 5, 6, 7$

C. $\text{cis} \left[(3k + 1) \frac{\pi}{8} \right], K = 0, 1, 2, 3, 4, 5, 6, 7$

D. $\text{cis} \left[(3k - 1) \frac{\pi}{8} \right], k = 0, 1, 2, 3, 4, 5, 6, 7$

Answer: A



Watch Video Solution

26. $\sqrt{a + \sqrt{a + \sqrt{a + \dots \infty}}} = \dots \dots$

A. $\frac{1 - \sqrt{4a - 1}}{2}$

B. $\frac{1 + \sqrt{4a - 1}}{2}$

C. $\frac{1 + \sqrt{4a + 1}}{2}$

D. $\frac{-1 - \sqrt{4a + 1}}{2}$

Answer: C



Watch Video Solution

27. $y = \frac{x^2 - 4x + 1}{x^2 - 4x + 3}$ does not lie between

A. $\left[\frac{3}{7}, 1 \right]$

B. $\left(\frac{-3}{7}, -1 \right)$

C. $\left[\frac{-3}{7}, -1 \right]$

D. $\left(\frac{3}{7}, 1 \right)$

Answer: D



View Text Solution

28. The remainder when $x^5 - 2x^4 + 4x^3 + 3x^2 + 5x - 7$ is divided by $x - 1$ is

A. 4

B. 3

C. 2

D. 1

Answer: A



[Watch Video Solution](#)

29. The equation whose roots are reciprocal of $3x^4 + 2x^2 - 2x - 3 = 0$ is

A. $x^4 + 3x^3 - 2x^2 + 3x - 3 = 0$

B. $3x^4 + 2x^3 - 2x^2 - 3 = 0$

C. $3x^4 - 2x^3 + 2x^2 + 3$

D. $3x^4 - 2x^3 - 2x^2 + 3 = 0$

Answer: B



[Watch Video Solution](#)

30. The sum of all 4 digit numbers that can be formed using the digit 0,1,2,3,4 without using zero is

A. 3, 33, 000

B. 5, 55, 500

C. 6, 66, 600

D. 2, 22, 200

Answer: C



[View Text Solution](#)

31. How many straight lines can be drawn by joining 15 distinct points on a circle ?

A. 100

B. 95

C. 90

D. 105

Answer: D



[Watch Video Solution](#)

32. The term independent of 'x' in the expansion of $\left(x^2 + \frac{1}{x}\right)^{12}$ is

A. 505

B. 500

C. 495

Answer: C [Watch Video Solution](#)

33. The sum of $(n + 1)$ terms of the series $\frac{C_0}{2} - \frac{C_1}{3} + \frac{C_2}{4} \dots$ is

A. $\frac{1}{n(n + 1)}$

B. $\frac{2}{(n + 1)(n + 2)}$

C. $\frac{1}{n + 1}$

D. $\frac{1}{n + 2}$

Answer: B [Watch Video Solution](#)

34. $\frac{x^2 + 5x + 7}{(x + 1)^3} =$

A. $\frac{1}{x+1} + \frac{3}{(x+1)^2} + \frac{3}{(x+1)^3}$

B. $\frac{1}{x+1} + \frac{3}{(x+1)^2} + \frac{2}{(x+1)^3}$

C. $\frac{2}{x+1} + \frac{1}{(x+1)^2} + \frac{3}{(x+1)^3}$

D. $\frac{1}{x+1} + \frac{2}{(x+1)^2} + \frac{3}{(x+1)^3}$

Answer: A



Watch Video Solution

35. The range of the series of value 10,12,13,11,7,6 is

A. 7

B. 8

C. 6

D. 9

Answer: A



Watch Video Solution

36. Standard deviation of first 'n' natural numbers is

A. $\sqrt{\frac{n^2 + 1}{12}}$

B. $\sqrt{\frac{n - 1}{nm}}$

C. $\sqrt{\frac{n^2 - 1}{12}}$

D. none

Answer: C



[Watch Video Solution](#)

37. Three persons A,B and C are to speak at a function along with 7 other persons. If the persons speak in random orders the probability that A speaks before B and B speaks before C is

A. $\frac{3}{5}$

B. $\frac{3}{8}$

C. $\frac{1}{6}$

D. none

Answer: C



View Text Solution

38. If $p(A) = \frac{3}{5}$, $p(B) = \frac{2}{5}$, $P(A \cap B) = \frac{1}{5}$ then, $P(\bar{A} \cap \bar{B}) =$

A. $\frac{6}{5}$

B. 1

C. $\frac{1}{2}$

D. $\frac{1}{5}$

Answer: D



Watch Video Solution

39. A,B,C are 3 newspaper from a city. 20% of the population read A, 16% read B, 14% read C, 8% both A and B, 5% both A and C, 4% both B and C, 2% all the three. Find the percentage of the populations who read atleast one newspaper.

A. 0.4

B. 0.35

C. 0.25

D. 0.3

Answer: B



Watch Video Solution

40. A bag contains 6 white and 4 black balls. A fair die is rolled and a number of balls equal to that appearing on the die is chosen from the bag at random. The probability that all the balls selected are white is

A. $\frac{1}{7}$

B. $\frac{1}{5}$

C. $\frac{1}{8}$

D. $\frac{1}{6}$

Answer: B



Watch Video Solution

41. For a binomial variate X if $n = 5$, and $P(X = 1) = 2P(X = 3)$, then

$p =$

A. $\frac{1}{3}$

B. $\frac{1}{10}$

C. $\frac{1}{15}$

D. $\frac{1}{20}$

Answer: A

 [Watch Video Solution](#)

42. The locus of a point P such that area of ΔPAB is 9 sq. units where $A = (2, 3)$ and $B(-4, 5)$ is

A. $3x + y - 2 = 0$

B. $x - 3y - 2 = 0$

C. $x + 3y - 2 = 0$

D. $x - 3y + 2 = 0$

Answer: C

 [Watch Video Solution](#)

43. If the axes are rotated through an angle 60° then the coordinates of a point $(2, -4\sqrt{3})$ in the old system are

A. $(5\sqrt{3}, -7)$

B. $(7, -\sqrt{3})$

C. $(3\sqrt{3}, -5)$

D. $(-1, -5)$

Answer: B



Watch Video Solution

44. Area of the triangle formed by the line passing through the points $(1,2),(-3,4)$ with the coordinate axes is

A. $\frac{25}{4}$ sq. units

B. $\frac{50}{3}$ sq. Units

C. $\frac{25}{2}$ sq. units

D. 25 sq. units

Answer: A



View Text Solution

45. For $a \neq b \neq c$ if the lines $x + 2ay + a = 0$, $x + 3by + b = 0$ and $x + 4cy + c = 0$ are concurrent, then a, b, c are in

A. A.G.P

B. G. P

C. A. P

D. H.P

Answer: D

 [Watch Video Solution](#)

46. The equations of the lines passing through $(4, 5)$ and making equal angles with the lines $3x = 4y + 7$, $5y = 12x + 6$ are

A. $9x - 7y - 1 = 0$, $7x + 9y - 73 = 0$

B. $y + 2 = 0$, $\sqrt{3}x - y - (2 + 3\sqrt{3}) = 0$

C. $3x + y = 4, x - 3y = -2$

D. none of the above

Answer: A



Watch Video Solution

47. If the slope of one of the lines $2x^2 + 3xy + \lambda y^2 = 0$ is 2 then the angle between the lines is

A. $\frac{\pi}{3}$

B. $\frac{\pi}{6}$

C. $\frac{\pi}{2}$

D. $\frac{\pi}{4}$

Answer: C



Watch Video Solution

48. The point of intersection of the pair of lines represented by

$$xy + 2x + 2y + 4 = 0 \text{ is}$$

A. $(-2, 2)$

B. $(-2, -2)$

C. $(2, -2)$

D. $(2, 2)$

Answer: B



[Watch Video Solution](#)

49. The ratio in which the line segment joining the points

$A(-3, -2, -1), B(1, 2, 1)$ is divided by the xz -plane is

A. 3:1

B. 2:1

C. 1:2

D. 1 : 1

Answer: D



[Watch Video Solution](#)

50. If the d.r.'s of two lines are $(1, 0, 0)$ and $(0, 1, 0)$ then the angle between those two lines is,

A. $\frac{\pi}{6}$

B. $\frac{\pi}{2}$

C. $\frac{\pi}{4}$

D. none

Answer: B



[Watch Video Solution](#)

51. The perpendicular distance from the origin to the plane $x + y + z + 3 = 0$ is

A. $\sqrt{3}$

B. $\sqrt{2}$

C. 1

D. 2

Answer: A



[Watch Video Solution](#)

52. $\lim_{x \rightarrow 0} \frac{\sqrt{4+x} - \sqrt[3]{8+3x}}{x} =$

A. -3

B. 0

C. $\frac{-1}{2}$

D. $\frac{1}{2}$

Answer: B

 [Watch Video Solution](#)

53. If $f(x) = x$ for $x < 0$ then $\lim_{x \rightarrow 0} f(x) =$

$= 0$ or $x = x0$

$= x^2$ or $x > 0$

A. 1

B. -1

C. 0

D. 2

Answer: C

 [View Text Solution](#)

54. Let $f(x) = \frac{x + x^2 + \dots + x^n - n}{x - 1}$, $x \neq 1$, the value of $f(1)$

A. $\frac{n + 1}{2}$

B. $\frac{n(n - 1)}{2}$

C. n

D. $\frac{n(n + 1)}{2}$

Answer: D



Watch Video Solution

55. $\frac{d}{dx} \left\{ \frac{a - b \cos x}{a + b \cos x} \right\} =$

A. $\frac{2ab \sin x}{(a - b \cos x)^2}$

B. $\frac{2ab \sin x}{(a + b \cos x)^2}$

C. $\frac{ab \sin x}{(a - b \cos x)^2}$

D. $\frac{ab \sin x}{(a + b \cos x)^2}$

Answer: B



Watch Video Solution

56. $\frac{d}{dx} [\cos^{-1}(4x^3 - 3x)] =$

A. $\frac{4}{\sqrt{1-x^2}}$

B. $\frac{1}{\sqrt{1-x^2}}$

C. $\frac{2}{\sqrt{1-x^2}}$

D. $\frac{3}{\sqrt{1-x^2}}$

Answer: D



Watch Video Solution

57. If $ax^2 + 2hxy + by^2 = 1$ then $(hx + by)^3 y_2 =$

A. $h^2 - ab$

B. 0

C. $ab - h^2$

D. $ab - h^3$

Answer: A



[Watch Video Solution](#)

58. If there is an error of 0.02 cm , while measuring the side of an equilateral triangle as 2 cm , then the percentage error in area is

A. 1

B. 3

C. 2

D. 4

Answer: C



[Watch Video Solution](#)

59. The length of the subtangent of the curve $x^2 + 3xy + 2y^2 = 0$ at (1,2)

is

A. $\frac{4}{11}$

B. $\frac{3}{11}$

C. $\frac{11}{2}$

D. $\frac{11}{4}$

Answer: D



Watch Video Solution

60. A particle moves on a line according to the law $s = at^2 + bt + c$. If the displacement after one second is 16 cm, the velocity after 2 second is 24 cm/sec and the acceleration is 8 cm/sec², then $(a, b, c) =$

A. (8,4,4)

B. (4,8,4)

C. (8,8,4)

D. (4,4,8)

Answer: B



[Watch Video Solution](#)

61. The constant c of Lagrange's mean value theorem for $f(x) = 2 \sin x + \sin 2x$ in $[0, \pi]$ is

A. $\frac{\pi}{3}$

B. $\frac{\pi}{6}$

C. $\frac{\pi}{2}$

D. $\frac{\pi}{4}$

Answer: A



[Watch Video Solution](#)

62. The equation of the diameter of the circle $x^2 + y^2 - 4x - 6y + 3 = 0$ that is perpendicular to $2x + 5y - 6 = 0$ is

A. $5x - 2y = 4$

B. $5x + 2y = 4$

C. $5x - 2y = -4$

D. $5x + 2y = -4$

Answer: A



[Watch Video Solution](#)

63. The equation of the circle touching both axes, lying in the first quadrant and having the radius 3 is

A. $x^2 + y^2 + 6x + 6y + 9 = 0$

B. $x^2 + y^2 - 6x - 6y + 9 = 0$

$$C. x^2 + y^2 - 6x - 6y - 9 = 0$$

$$D. x^2 + y^2 + 6x - 6y + 0$$

Answer: B



Watch Video Solution

64. If the points $(k,1)$ $(2,-3)$ are conjugate w.r.t.

$x^2 + y^2 + 4x - 6y - 12 = 0$ then k

A. $\frac{2}{3}$

B. $\frac{5}{4}$

C. 1

D. -3

Answer: B



Watch Video Solution

65. The radical axis of the circles $x^2 + y^2 + 2x - 3y - 5 = 0$ is

A. $4x - 7y - 20 = 0$

B. $4x + 7y - 20 = 0$

C. $4x + 7y + 20 = 0$

D. $4x - 7y + 20 = 0$

Answer: C



View Text Solution

66. The condition that the circle $x^2 + y^2 + 2gx + 2fy + c = 0$ to bisect the circumference of the circle $x^2 + y^2 + 2g^1x + 2f^1y + c^1 = 0$ is

A. $g(g - g^1) + f(f - f^1) = c - c^1$

B. $g^1(g - g^1) + f^1(f - f^1) = c - c^1$

C. $2g^1(g - g^1) + 2f^1(f - f^1) = c - c^1$

D. $2g(g - g^1) + 2f(f - f^1) = c - c^1$

Answer: C



Watch Video Solution

67. The equation of the parabola having focus (1,2) and directrix

$$x + 2y + 6 = 0 \text{ is}$$

A. $4x^2 + 4xy + y^2 + 22x - 44y - 11 = 0$

B. $4x^2 - 4xy + y^2 - 22x + 44y - 11 = 0$

C. $4x^2 + 4xy + y^2 + 22x + 54y + 11 = 0$

D. $4x^2 - 4xy + y^2 - 22x - 44y - 11 = 0$

Answer: D



Watch Video Solution

68. If the points (2,4) ,(k,6) are conjugate with respect to the parabola

$$y^2 = 4x \text{ then } k =$$

A. -12

B. -2

C. $\frac{7}{2}$

D. 10

Answer: D



Watch Video Solution

69. The condition that the line $lx + my + n = 0$ to be a normal to the

ellipse $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$ is

A. $\frac{a^2}{l^2} - \frac{b^2}{m^2} = \frac{(a^2 - b^2)^2}{n^2}$

B. $\frac{a^2}{l^2} + \frac{b^2}{m^2} = \frac{(a^2 - b^2)^2}{n^2}$

C. $\frac{a^2}{l^2} + \frac{b^2}{m^2} = \frac{(a^2 + b^2)^2}{n^2}$

D. $\frac{a^2}{l^2} - \frac{b^2}{m^2} = \frac{(a^2 + b^2)^2}{n^2}$

Answer: B



Watch Video Solution

70. The length of the latus - rectum of the ellipse

$$9x^2 + 25y^2 - 18x - 100y - 116 = 0 \text{ is}$$

A. $9/2$

B. $8/5$

C. $8/3$

D. $18/5$

Answer: B



Watch Video Solution

71. The foci of the ellipse $\frac{x^2}{16} + \frac{y^2}{b^2} = 1$ and the hyperbola

$$\frac{x^2}{144} - \frac{y^2}{81} = \frac{1}{25}$$

coincide. Then, the value of b^2 is

A. 7

B. 1

C. 9

D. 5

Answer: A

 [Watch Video Solution](#)

72. $\int \frac{1 + \sin^2 x}{1 + \cos 2x} dx =$

A. $\tan x + \frac{x}{2} + c$

B. $2 \tan x - \frac{x}{2} + c$

C. $\tan x - \frac{x}{2} + c$

D. $2 \tan x - \frac{x}{2} + c$

Answer: C

 [Watch Video Solution](#)

$$73. \int \frac{1}{\sqrt{x} + x} dx =$$

A. $2 \log(1 + \sqrt{x}) + c$

B. $\log(x + \sqrt{x}) + c$

C. $2 \log(x + \sqrt{x}) + c$

D. $\log(1 + \sqrt{x}) + c$

Answer: A



Watch Video Solution

$$74. \int \frac{1}{(1 + x^2)\sqrt{1 - x^2}} dx =$$

A. $\frac{-1}{\sqrt{2}} \tan^{-1} \left(\frac{\sqrt{1 + x^2}}{\sqrt{2}x} \right) + c$

B. $\frac{1}{\sqrt{2}} \tan^{-1} \left(\frac{\sqrt{1 + x^2}}{\sqrt{2}x} \right) + c$

C. $\frac{-1}{\sqrt{2}} \tan^{-1} \left(\frac{\sqrt{1 - x^2}}{\sqrt{2}x} \right) + c$

D. none

Answer: C

 [Watch Video Solution](#)

$$75. \int \frac{(x+1)dx}{x(1+xe^x)} =$$

A. $\log(1+xe^x)xe^x + c$

B. $\log\left[\frac{xe^x}{1+xe^x}\right] + c$

C. $\log\left[\frac{1+xe^x}{xe^x}\right] + C$

D. $\log(1+xe^x) + C$

Answer: D

 [Watch Video Solution](#)

$$76. \int_0^1 \frac{dx}{e^x + e^{-x}} =$$

A. $\tan^{-1}(e) - \frac{\pi}{2}$

B. $\tan^{-1}(e) + \frac{\pi}{4}$

C. $\tan^{-1}(e) - \frac{\pi}{4}$

D. none

Answer: C

 [Watch Video Solution](#)

77. $\int_0^{\pi/4} \left[\sqrt{\frac{1 - \sin 2x}{1 + \sin 2x}} \right] dx =$

A. $\log 2$

B. $\frac{1}{2} \log 2$

C. $-\log 2$

D. $\frac{-1}{2} \log 2$

Answer: B

 [Watch Video Solution](#)

78. The area of the region bounded by the curves $y = 2^x$, $y = 2x - x^2$ and the lines $x = 0$, $x = 2$ is

A. $\frac{3}{\log 2} - \frac{4}{3}$

B. $\frac{3}{\log 2} + \frac{4}{3}$

C. $3 \log 2 - \frac{4}{3}$

D. $\frac{1}{\log 2} - \frac{4}{3}$

Answer: A



Watch Video Solution

79. The solution of $e^{x-y} dx + e^{y-x} dy = 0$ is

A. $e^x - e^y = C$

B. $e^{2x} - e^{2y} = C$

C. $e^{2x} + e^{2y} = c$

D. $e^x + e^y = C$

Answer: B



Watch Video Solution

80. Integrating factor of $\sin x \frac{dy}{dx} + y \cos x = \sin 2x$ is

A. $-\sin x$

B. $-\cos x$

C. $\cos x$

D. $\sin x$

Answer: C



Watch Video Solution

1. If a unit vector is represented by $0.5\hat{i} + 0.8\hat{j} + c\hat{k}$, the value of c is

A. $\sqrt{0.89}$

B. 0.2

C. 0.3

D. $\sqrt{0.11}$

Answer: D



[Watch Video Solution](#)

2. A thermos flask contains 250 g coffee at $90^\circ C$. To this 20 g of milk at $5^\circ C$ is added. After equilibrium is established, the temperature of the liquid is

(Assume no heat loss to the thermos bottle . Take specific heat of coffee and milk as $1.00\text{cal} / \text{g}^\circ C$)

A. $3.23^\circ C$

B. $3.17^{\circ}C$

C. $83.7^{\circ}C$

D. $37.8^{\circ}C$

Answer: C



Watch Video Solution

3. A copper rod of length 75 cm and an iron rod of length 125cm are joined together end to end . Both are of circular cross section with diameter 2 cm . The free ends of the copper and iron are maintained at $100^{\circ}C$ and $0^{\circ}C$ respectively . The surface of the bars are insulated thermally . The temperature of the copper -iron junction is [Thermal conductivity of the copper is $386.4W/m - K$ and that of iron is $48.46W/m - K$].

A. $100^{\circ}C$

B. $0^{\circ}C$

C. $93^{\circ}C$

D. $50^{\circ}C$

Answer: C



[Watch Video Solution](#)

4. Total emf produd in a thermocouple does not depend on

A. The metals in the thermocouple

B. Thomson coefficients of the metals in the thermocouple

C. Temperature of the junctions

D. The duration of time for which the current is passed through thermocouple

Answer: D



[Watch Video Solution](#)

5. A ball is projected vertically down with an initial velocity from a height of 20 m on to a horizontal floor. During the impact it loses 50% of the energy and rebounds to the same height, the initial velocity of its projection is

A. $20ms^{-1}$

B. $15ms^{-1}$

C. $10ms^{-1}$

D. $5ms^{-1}$

Answer: A



Watch Video Solution

6. Moment of inertia of a body about an axis is $4kg - m^2$. The body is initially at rest and a torque of 8 N-m starts along the same axis. Work done by the R, about an axis which is a tangent and parallel to its torque in 20 s, in joules, is

A. 40

B. 640

C. 2560

D. 3200

Answer: D



Watch Video Solution

7. An electrically charged particle enters into a uniform magnetic induction field in a direction perpendicular to the field with a velocity v . Then, it travels

A. In a straight line without acceleration

B. With force in the direction of the field

C. In a circular path with a radius directly proportional to v^2

D. In a circular path with radius directly proportional to its velocity

Answer: D



Watch Video Solution

8. At a certain place, the angle of dip is 60° and the horizontal component of the earth's magnetic field (B_H) is 0.8×10^{-4} T. The earth's overall magnetic field is

A. $1.5 \times 10^{-4} T$

B. $1.6 \times 10^{-3} T$

C. $1.5 \times 10^{-3} T$

D. $1.6 \times 10^{-4} T$

Answer: D



Watch Video Solution

9. Wires A and B have resistivities ρ_A and ρ_B . If the diameter of the wire B is twice that of A and the Two wires have same resistance, then $\frac{L_b}{L_A}$ is

A. 2

B. 1

C. $\frac{1}{2}$

D. $\frac{1}{4}$



Answer: A



[Watch Video Solution](#)

10. Use Lenz's law to determine the direction of induced current in the situations described by fig. a, b.

A circular loop being deformed into a narrow straight wire.



A. Towards left

B. In a direction opposite to change of the magnetic flux

C. Towards right

D. In the direction of change of the magnetic flux

Answer: B



Watch Video Solution

11. A motor of power P_0 is used to deliver water at a certain rate through a given horizontal pipe. To increase the rate of flow of water through the

same pipe n times, the power of the motor is increased to P_1 to P_0 is

A. $n : 1$

B. $n^2 : 1$

C. $n^3 : 1$

D. $n^4 : 1$

Answer: A



[Watch Video Solution](#)

12. A body of mass 5 kg makes an elastic collision with another body at rest and continues to move in the original direction after collision with velocity equal to $\frac{1}{10}$ th of its original velocity . Then the mass of the second body is

A. $4.09kg$

B. $0.5kg$

C. 5 kg

D. 5.09kg

Answer: A



Watch Video Solution