



## MATHS

### BOOKS - SAI MATHS (TELUGU ENGLISH)

### MOCK TEST 2

#### Maths

1. If  $f: R \rightarrow R, S: R \rightarrow R$  are defined by  $f(x) = 3x-4, g(x) = 5x-1$  then,

$$(f \circ g^{-1})(2) =$$

A.  $\frac{11}{5}$

B.  $-\frac{11}{5}$

C.  $\frac{1}{5}$

D.  $-\frac{1}{5}$

**Answer: B**

[View Text Solution](#)

2. The range of  $y = 2x^2 + x + \frac{2}{2x^2 + x + 1}$  is

- A.  $\left( 11 - \frac{\sqrt{2}}{7}, 11 + \frac{\sqrt{2}}{7} \right)$   
B.  $\left[ 11 - \frac{\sqrt{2}}{7}, 11 + \frac{\sqrt{2}}{7} \right]$   
C.  $\left( -11 + \frac{\sqrt{2}}{7}, -11 - \frac{\sqrt{2}}{7} \right)$   
D.  $\left[ -11 + \frac{\sqrt{2}}{7}, -11 + \frac{\sqrt{2}}{7} \right]$

**Answer: B**

[View Text Solution](#)

3.  $\sum \left( \frac{1^2 + 2^2 + 3^2 + \dots + n^2}{1 + 2 + 3 + \dots + n} \right)$

A.  $n^2 + \frac{11}{12}n$

B.  $n^2 - 2\frac{n}{6}$

C.  $\frac{n^2 + 2n}{3}$

D. None

**Answer: C**

 [Watch Video Solution](#)

4. If  $A = \begin{bmatrix} 1 & 1 & 0 \\ 0 & 1 & 0 \\ 1 & 0 & 1 \end{bmatrix}$  then  $A^3 =$

A.  $\begin{bmatrix} 1 & 3 & 0 \\ 0 & 1 & 0 \\ 3 & 3 & 1 \end{bmatrix}$

B.  $\begin{bmatrix} 1 & 0 & 3 \\ 3 & 1 & 3 \\ 0 & 0 & 1 \end{bmatrix}$

C.  $\begin{bmatrix} 3 & 3 & 1 \\ 1 & 3 & 0 \\ 0 & 1 & 0 \end{bmatrix}$

D.  $\begin{bmatrix} 3 & 1 & 0 \\ 3 & 3 & 1 \\ 1 & 0 & 0 \end{bmatrix}$

**Answer: A**

 [View Text Solution](#)

5. If  $\frac{1}{a} + \frac{1}{b} + \frac{1}{c} = 0$  then  $\begin{vmatrix} 1+a & 1 & 1 \\ 1 & 1+b & 1 \\ 1 & 1 & 1+c \end{vmatrix} =$

A.  $-abc$

B. 0

C.  $abc$

D. None

**Answer: C**



**Watch Video Solution**

6. If the system of equations  $\lambda x + 3y + z = 0$ ,  $4x + \lambda y + 3z = 0$ ,  $2x + 3y + \lambda z = 0$  has non-trivial solution, then  $\lambda =$

A. 6

B. 3

C. 2

D. None

**Answer: D**



[View Text Solution](#)

7. If  $z_1, z_2, z_3$  are collinear and  $z_3 - \frac{z_1}{z_2 - z_1}$  is purely real, then  $\arg\left(z_3 - \frac{z_1}{z_2 - z_1}\right)$

A. 0

B. 1

C. 2

D. 3

**Answer: A**



[View Text Solution](#)

8. If  $\omega, \omega^2$  are cube root of unity then,  $\frac{\omega}{1} + \omega^2 + \frac{\omega}{1} + \omega =$

A.  $-2\omega^2$

B.  $-2\omega$

C. 2

D. -2

**Answer: D**



[View Text Solution](#)

9. If  $z = 3 + 3i$  then,  $z^2 + z + 15 =$



[Watch Video Solution](#)

10. If the area of the triangle on the complex plane formed by the points  $z, iz$  and  $z+iz$  is 50 sq. units then  $|z|$  is

A. 15

B. 10

C. 5

D. None

**Answer: B**



[View Text Solution](#)

11. If  $A, B$  and  $C$  are the angles of a triangle such that  $\cos A + \cos B + \cos C = 0 = \sin A + \sin B + \sin C$ , then  $\sin 3A + \sin 3B + \sin 3C =$

A. 1

B. 2

C. 0

D. 3

**Answer: C**

 [View Text Solution](#)

12. If  $x^2 + 3x + 2 = 0$ ,  $x^2 + 6x + k = 0$  have a common root then  $p =$

A. 10 (or) 16

B. 11 (or) 15

C. 8 (or) 4

D. 5 (or) 8

**Answer: D**

 [Watch Video Solution](#)

13. If  $x$  is real, then the maximum value of  $\frac{x^2 + 14x + 9}{x^2 + 2x + 3}$  is

A. 6

B. 8

C. 4



D. 2

**Answer: C**



[Watch Video Solution](#)

14. The condition that the roots of  $x^3 + 3px^2 + 3qx + r = 0$  may be in

A.P is

A.  $2q^3 + r^2 = 3pqr$

B.  $2p^3 + r = 3pq$

C.  $p^3r = q^3$

D. None

**Answer: B**



[Watch Video Solution](#)

15. The roots of  $2x^5 + x^4 - 12x^3 - 12x^2 + x + 2 = 0$  are

A.  $-1, -2, -\frac{1}{2}, 3 \pm \frac{\sqrt{5}}{2}$

B.  $2 \pm \sqrt{3}, 3 \pm \sqrt{2}$

C.  $2, \frac{1}{2}, 3, \frac{1}{3}, 1$

D.  $1, 2, \frac{1}{2}, 5 \pm \sqrt{11} \frac{i}{6}$

**Answer: A**



[Watch Video Solution](#)

16. How many numbers can be made with digit 3,4,5,6,7,8 lying between 3000 and 4000 which are divisible by 5 without repetition

A. 120

B. 60

C. 24

D. 12

**Answer: D**



**Watch Video Solution**

17. If  $n_{pr} = 840$ ,  $n_{cr} = 35$ , then  $n$  is equal to

A. 1

B. 7

C. 4

D. 10

**Answer: B**



**View Text Solution**

18. The co-efficient of  $x^{11}$  in the expansion of  $(1 + 3x + 2x^2)^6$  is

A. 216

B. 144

C. 576

D. 288

**Answer: C**



**Watch Video Solution**

19.  $1 + \frac{1}{3}x + \frac{1.4}{3.6}x^2 + \frac{1.4.7}{3.6.9}x^3 + \dots =$

A.  $(1 + x)^{\frac{1}{3}}$

B.  $(1 - x)^{\frac{1}{3}}$

C.  $(1 - x)^{-\frac{1}{3}}$

D.  $x$

**Answer: C**



**Watch Video Solution**

20. The partial fractions of  $\frac{1}{x^3(x+2)} =$

A.  $\frac{1}{8}x + \frac{1}{4}x^2 + \frac{1}{2}x^3 - \frac{1}{8(x+2)}$

B.  $\frac{1}{8}x - \frac{1}{4}x^2 + \frac{1}{2}x^3 - \frac{1}{8(x+2)}$

C.  $\frac{1}{8}x + \frac{1}{4}x^2 + \frac{1}{2}x^3 + \frac{1}{8(x+2)}$

D.  $\frac{1}{8}x - \frac{1}{4}x^2 - \frac{1}{2}x^3 + \frac{1}{8(x+2)}$

**Answer: B**



**Watch Video Solution**

21. If  $70 \sin^2 \theta + 3 \cos^2 \theta = 4$ , then  $\tan \theta$

A.  $\pm \frac{1}{\sqrt{3}}$

B.  $\frac{1}{3}$

C.  $\pm \sqrt{3}$

D.  $\pm 1$

**Answer: A**



**Watch Video Solution**

22. If 
$$\begin{vmatrix} 1 + \sin^2 \theta & \cos^2 \theta & 4 \sin 4\theta \\ \sin^2 \theta & 1 + \cos^2 \theta & 4 \sin 4\theta \\ \sin^2 \theta & \cos^2 \theta & 1 + 4 \sin 4\theta \end{vmatrix} = 0$$
, then the value of  $\theta$  is

A.  $7\frac{\pi}{24}$  or  $11\frac{\pi}{24}$

B.  $\frac{\pi}{24}$

C.  $5\frac{\pi}{24}$

D. None

**Answer: A**



**Watch Video Solution**

23. 
$$\tan \left[ \frac{1}{2} \sin^{-1} \frac{2a}{1+a^2} + \frac{1}{2} \cos^{-1} \frac{1-a^2}{1+a^2} \right] =$$

A.  $1 - \frac{a^2}{1 + a^2}$

B.  $2 \frac{a}{1 + a^2}$

C.  $\frac{1 + a^2}{1 - a^2}$

D.  $2 \frac{a}{1} - a^2$

**Answer: D**



**Watch Video Solution**

**24.** If  $\sin x \cosh y = \cos \theta$ ,  $\cos x \sinh y = \sin \theta$  then  $\sinh^2 y =$

A.  $\cosh^2 y$

B.  $\cos^2 x$

C.  $\sec^2 x$

D.  $\cosh^2 x$

**Answer: B**



**Watch Video Solution**

25. If  $m \cdot \tan(\theta - 30^\circ) = n \cdot \tan(\theta + 120^\circ)$ , then  $\cos 2\theta =$

A.  $\sin 2\theta$

B.  $\cos 2\theta$

C.  $2 \sin 2\theta$

D.  $2 \cos 2\theta$

**Answer: D**



**Watch Video Solution**

26. If  $\tan\left(B - \frac{C}{2}\right) = x \frac{\cot A}{2}$ , then  $x =$

A.  $a - \frac{b}{a+b}$

B.  $b - \frac{c}{b+c}$

C.  $c - \frac{a}{c+a}$



D. None

**Answer: B**



[View Text Solution](#)

27. If  $P_1, P_2, P_3$  are altitudes of  $\triangle ABC$  from the vertices A,B,C and  $\Delta$  is the area of triangle then,  $\frac{1}{P_1^2} + \frac{1}{P_2^2} + \frac{1}{P_3^2} =$

A.  $a^2 + b^2 + \frac{c^2}{4}\Delta^2$

B.  $a^2 - b^2 - \frac{c^2}{\Delta^2}$

C.  $a + b + \frac{c}{\Delta}$

D.  $a^2 + b^2 + \frac{c^2}{\Delta^2}$

**Answer: A**



[View Text Solution](#)

28. From the top of a mast of 60 m height, the angle of depression of an object is  $45^\circ$ . The distance of the object from the ship is

- A. 45 m
- B. 80 m
- C. 60 m
- D. 90 m

**Answer: C**



[Watch Video Solution](#)

29. If the position vector of A,B,C are  $2i + 3j + 4k$ ,  $i + 2j$ ,  $j + 2k$  and

$\vec{AB} = P\vec{AC}$  then P=

- A.  $-\frac{1}{2}$
- B.  $\frac{1}{4}$
- C.  $\frac{1}{2}$

D.  $-\frac{1}{4}$

**Answer: C**



[View Text Solution](#)

**30.** If the position vectors of A, B are  $2a - 3b$ ,  $3a + 2b$  respectively then the position of vector of C in AB produced such that  $AC = 2 AB$  is

A.  $4a+7b$

B.  $5b-2a$

C.  $3a+2b$

D.  $3b-2a$

**Answer: A**



[Watch Video Solution](#)

31. The vector equation of the line passing through the point  $i+j-k$  and parallel to the vector  $2i+3j-k$  is

A.  $r=(i+j-k)+t(2i+3j-k)$

B.  $r=(i-j+k)+t(2i-3j-k)$

C.  $r=(i-j-k)+t(2i-3j-k)$

D.  $r=(-i-j+k)+t(-2i+3j+k)$

**Answer: A**



[View Text Solution](#)

32. The length of projection of  $i+2j+3k$  in the direction of  $3i-4j+5k$  is

A.  $\sqrt{2}$

B.  $\sqrt{3}$

C. 1

D. 2

**Answer: A**

 [Watch Video Solution](#)

**33.** A unit vector perpendicular to each of the vector  $3i+2j+4k$  and  $2i+j-k$  is,

A.  $\pm 6i + 8j + \frac{k}{\sqrt{101}}$

B.  $\pm 6i - 8j + \frac{k}{\sqrt{104}}$

C.  $\pm 6i - 8j - \frac{k}{\sqrt{101}}$

D.  $\pm 6i - 8j - 4\frac{k}{\sqrt{101}}$

**Answer: A**

 [View Text Solution](#)

**34.** The vector  $i+j+k, i+2j+3k, 2i+3j+k$  are

A. Collinear

B. Non-coplanar

C. Coplanar

D. None

**Answer: B**



[View Text Solution](#)

**35.** The variance of 6,5,8,10,3,4,9,11 is

A. 8

B. 9

C. 7.5

D. 10

**Answer: C**



[View Text Solution](#)

36. If 6 cards are drawn at random, from a pack of cards, then the probability to get 3 red and 3 black cards is

A.  $\frac{{}^{28}C_3 \times {}^{28}C_3}{{}^{56}C_6}$

B. None

C.  $\frac{{}^{16}C_3 \times {}^{16}C_3}{{}^{32}C_6}$

D.  $\frac{{}^{26}C_3 \times {}^{26}C_3}{{}^{52}C_6}$

**Answer: D**



**Watch Video Solution**

37. Let  $A, B, C$  are three events such that  $P(A) = 0.2, P(B) = 0.5, P(C) = 0.6, P(A \cap B) = 0.15, P(A \cap C) = 0.2$ , then,

A.  $0.05 \leq P(B \cap C) \leq 0.5$

B.  $0.03 \leq P(B \cap C) \leq 0.38$

C.  $0.06 \leq P(B \cap C) \leq 0.24$

D.  $0.04 \leq P(B \cap C) \leq 0.22$

**Answer: C**



[View Text Solution](#)

**38.** In a class, 40% students study botany, 25% Zoology and 15% bith Botany and Zoology.A student from the class is selected at random. The probability that he studies Botany,if it is known that he studies Zoology is

A.  $\frac{1}{8}$

B.  $\frac{2}{5}$

C.  $\frac{3}{8}$

D.  $\frac{3}{5}$

**Answer: D**



[View Text Solution](#)



39. There are 2 red, 4 green balls in bag A, bag B, there are 5 red and 7 green balls. If one ball is randomly replaced from A into B and a ball is drawn from B then the probability for the ball to be red is

A.  $\frac{17}{40}$

B.  $\frac{14}{40}$

C.  $\frac{19}{45}$

D.  $\frac{16}{39}$

**Answer: D**



[View Text Solution](#)

40. If  $X$  is random variable with distribution given below

$x: 1 \ 2 \ 3 \ 4$

$P(X=x): k \ k \ 2k \ 3k$

The value of  $k$  and its mean are



Watch Video Solution

41. If  $X$  is a poisson distribution such that  $P(X=1)=P(X=2)$  then,  $P(X=4)=$

A.  $\frac{1}{3}e^2$

B.  $\frac{2}{3}e^2$

C.  $\frac{4}{3}e^2$

D. none

Answer: B



View Text Solution

42. A(2, 1) and B(2, 3) are two points. If P is a point such that  $PA + PB = 2$ ,

then the locus of P is

A.  $4x^2 - 12y^2 - 16x + 124y - 69 = 0$

B.  $4x^2 + 12y^2 - 16x - 124y + 69 = 0$

C.  $4x^2 + 12y^2 + 16x - 124y + 69 = 0$

D.  $4x^2 + 12y^2 + 16x + 124y - 69 = 0$

**Answer: A**



[View Text Solution](#)

**43.** The transformed equation of  $xy + 2x - 5y - 11 = 0$  when the origin is shifted to the point  $(2, 3)$  is,

A.  $xy - 5x - 3y + 16 = 0$

B.  $xy + 5x + 3y - 16 = 0$

C.  $xy + 5x - 3y - 16 = 0$

D.  $xy - 5x + 3y + 16 = 0$

**Answer: C**



[View Text Solution](#)

44. The area of the triangle formed by the line  $\frac{x}{5} + \frac{y}{4} = 1$  with the coordinate axes is

- A. 20 sq.units
- B. 15 sq.units
- C. 5 sq.units
- D. 10 sq.units

**Answer: D**

 [View Text Solution](#)

45. The perpendicular distance of the straight line  $3x + 4y - 8 = 0$  from the point of intersection of the lines  $3x + 2y + 4 = 0$ ,  $2x + 5y - 1 = 0$  is

- A.  $\frac{11}{5}$  units
- B.  $\frac{12}{5}$  units
- C.  $\frac{8}{5}$  units

D. 2units

**Answer: D**



[Watch Video Solution](#)

**46.** The diagonal of a square is  $8x - 15y = 0$  and one vertex of the square is  $(1, 2)$ . The equations to the sides of the square passing through this vertex are

A.  $23x + 7y = 9, 7x - 23y = 52$

B.  $23x + 7y = 9, 7x - 23y = 53$

C.  $22x + 8y = 9, 22x - 8y = 52$

D. None

**Answer: B**



[Watch Video Solution](#)

47. The area of the triangle formed by the pair of lines

$3x^2 + 8xy - 3y^2 = 0$  and the line  $3x + 4y - 5 = 0$  is,

A.  $\frac{3}{5}$  sq. units

B.  $\frac{5}{3}$  sq. units

C.  $\frac{4}{5}$  sq. units

D.  $\frac{5}{4}$  sq. units

**Answer: B**



[Watch Video Solution](#)

48. If  $x^2 - 10xy + 4y^2 + 6x + 2y + k = 0$  represents a pair of straight

lines then,  $k =$

A.  $-\frac{1}{3}$

B.  $\frac{2}{3}$

C.  $-\frac{2}{3}$

D. None

**Answer: A**

 [Watch Video Solution](#)

49. The equation of the circle concentric with  $x^2 + y^2 - 2x + 8y - 23 = 0$  and passing through  $(2, 3)$  is

A.  $x^2 + y^2 - 2x - 8y - 33 = 0$

B.  $x^2 + y^2 + 6x - 4y - 12 = 0$

C.  $x^2 + y^2 + x + 8y + 33 = 0$

D.  $x^2 + y^2 - 6x + 4y - 12 = 0$

**Answer: A**

 [Watch Video Solution](#)

50. If the tangent to the circle  $x^2 + y^2 = 5$  at  $(1, -2)$  also touches the circle  $x^2 + y^2 - 8x + 6y + 20 = 0$  then the point of contact is

- A.  $(-1, 0)$
- B.  $(1, 0)$
- C.  $(3, -1)$
- D.  $(5, 2)$ .

**Answer: C**



[Watch Video Solution](#)

51. If the circles  $x^2 + y^2 - 6x - 8y + c = 0$  and  $x^2 + y^2 = 9$  have three common tangents then  $c =$

- A. 17
- B. 19
- C. 21



D. 20

**Answer: C**

 [View Text Solution](#)

52. The equation of the circle which cuts orthogonally the circle  $x^2 + y^2 - 4x + 2y - 7 = 0$  and having centre at  $(2, 3)$  is,

A.  $3x^2 - 3y^2 + 2x - 12y + 2 = 0$

B.  $x^2 + y^2 + 6x + 4y + 19 = 0$

C.  $x^2 + y^2 - 2x + 4y + 2 = 0$

D.  $x^2 + y^2 - 4x - 6y + 9 = 0$

**Answer: D**

 [View Text Solution](#)

53. If  $(2, 1)$  is limiting point of coaxial system of which  $x^2 + y^2 - 6x - 4y - 3 = 0$  is a member, then the other limiting point is

A.  $(-5,-6)$

B.  $(-2,-3)$

C.  $(3,2)$

D.  $(5,6)$

**Answer: A**



[View Text Solution](#)

54. The locus of the point of intersection of tangents to parabola  $y^2 = 4(x + 1)$  and  $y^2 = 8(x + 2)$  which are perpendicular to each other is

A.  $x-3=0$

B.  $x+3=0$

C.  $y-3=0$

D.  $y+3=0$

**Answer: B**



[View Text Solution](#)

55. If the normals at the points  $t_1$  and  $t_2$  on  $y^2 = 4ax$  at the point  $t_3$  on the parabola, the  $t_1 t_2 =$

A. 4

B. 3

C. 2

D.  $2t_3$

**Answer: C**



[View Text Solution](#)

56. The eccentricity of the ellipse  $9x^2 + 16y^2 = 144$  is

A.  $\frac{7}{4}$

B.  $\frac{7}{2}$

C.  $\frac{\sqrt{7}}{2}$

D.  $\frac{\sqrt{7}}{4}$

Answer: D



Watch Video Solution

57. The condition that the line  $x \cos \alpha + y \sin \alpha = P$  may be a normal to

the ellipse  $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$  is

A.  $\frac{a^2}{\cos^2 \alpha} - \frac{b^2}{\sin^2 \alpha} = \left( a^2 - \frac{b^2}{P^2} \right)^2$

B.  $\frac{a^2}{\cos^2 \alpha} - \frac{b^2}{\sin^2 \alpha} = \left( a^2 + \frac{b^2}{P^2} \right)^2$

C.  $\frac{a^2}{\cos^2 \alpha} + \frac{b^2}{\sin^2 \alpha} = \left( a^2 - \frac{b^2}{P^2} \right)^2$

D.  $\frac{a^2}{\cos^2 \alpha} + \frac{b^2}{\sin^2 \alpha} = \left( a^2 + \frac{b^2}{P^2} \right)^2$

**Answer: C**



**View Text Solution**

**58.** The distance between the foci is  $4\sqrt{13}$  and the length of conjugate axis is 8 then, the eccentricity of the hyperbola is

A.  $\frac{\sqrt{15}}{4}$

B.  $\frac{\sqrt{13}}{4}$

C.  $\frac{\sqrt{13}}{3}$

D.  $\frac{\sqrt{13}}{2}$

**Answer: C**



**Watch Video Solution**

**59.** If the d.c.'s  $(l, m, n)$  of two lines are connected by the relations  $l + m + n = 0$  and  $2mn + 3ln - 5lm = 0$  then the angle between

the lines is

A.  $\frac{\pi}{2}$

B.  $\frac{\pi}{6}$

C.  $\frac{\pi}{3}$

D. None

**Answer: A**



[Watch Video Solution](#)

**60.** The plane  $2x + 3y + kz - 7 = 0$  is parallel to the line whose d.r's are  $(2, -3, 1)$  then  $k =$

A. 5

B. 10

C. 15

D. 20

**Answer: A**



**Watch Video Solution**

**61.** The centroid of the triangle formed by the points (1, 2,3), (2, 3, 1), (3, 1, 2) is

A. (1,1,1)

B. (2,2,2)

C. (1,2,2)

D. (3,1,3)

**Answer: B**



**View Text Solution**

**62.** 
$$\lim_{x \rightarrow 0} \left( \frac{1 - 2 \cos x + \cos 2x}{x^2} \right)$$

A. 2

B. 3

C. -1

D. 0

**Answer: C**

 [Watch Video Solution](#)

63.  $\lim_{x \rightarrow 2^-} \{x + (x - [x])^2\} =$

A. 3

B. 1

C. 2

D. None

**Answer: A**

 [Watch Video Solution](#)



64.

If

$$f(x) = \left\{ \left( -2 \sin x, f \text{ or } x \leq -\frac{\pi}{2} \right) \left( a \sin x + b, f \text{ or } -\frac{\pi}{2} < x < \frac{\pi}{2} \right) \right.$$

everywhere then the ordered pair(a,b) is

A. (-1,1)

B. (1,1)

C. (0,0)

D. None

**Answer: A**



[View Text Solution](#)

65. If  $y = x \log \left| x + \sqrt{1+x^2} \right| - \sqrt{1+x^2}$  then  $\frac{dy}{dx} =$

A.  $\cos^{-1} hx$

B.  $\frac{1}{2} \log(x - \sqrt{1+x^2})$

C.  $\cos ec^{-1}hx$

D.  $\sin^{-1}hx$

**Answer: D**



**Watch Video Solution**

66. The derivative of  $e^{\sin^{-1}x}$  w.r.t  $\log x$  is

A.  $\frac{e^{\sin^{-1}x}}{\sqrt{1+x^2}}$

B.  $x \frac{e^{\sin^{-1}x}}{\sqrt{1+x^2}}$

C.  $\frac{e^{\sin^{-1}x}}{\sqrt{1+x^2}}$

D.  $x \frac{e^{\sin^{-1}x}}{\sqrt{1-x^2}}$

**Answer: D**



**Watch Video Solution**

67. If  $y = ae^{nx} + be^{-nx}$  then  $y_2 =$

A.  $-n^2y$

B.  $ny$

C.  $n^2y$

D.  $-ny$

Answer: C



Watch Video Solution

68. Equation of the tangent to the curve  $y^2 = 4ax$  at  $(at^2, 2at)$  is

A.  $xt + y - 2at - at^3 = 0$

B.  $x + yt = at^2$

C.  $x - yt + at^2 = 0$

D.  $xt - y - 2at - at^3 = 0$

**Answer: C**



**Watch Video Solution**

69. the side of a square is equal to the diameter of a circle. If the side and radius change at the same rate then the ratio of the change of their areas is

A.  $2 : \pi$

B.  $1 : 1$

C.  $\pi : 2$

D.  $1 : \pi$

**Answer: A**



**Watch Video Solution**

70. Verify Rolle's theorem for the function  $f(x) = x(x + 3)e^{-\frac{x}{2}}$  in  $[-3, 0]$ .

A. -2

B. 1

C. -1

D. 0

**Answer: A**



[Watch Video Solution](#)

71. Show that when the curved surface of a right circular cylinder inscribed in a sphere of radius  $R$  is maximum, then the height of the cylinder is  $\sqrt{2R}$ .

A.  $\sqrt{3}R$

B.  $\sqrt{2}R$

C.  $2R$

D.  $R$

**Answer: B**



**Watch Video Solution**

$$72. \int \left( \sin 2 \frac{x}{a \cos^2 x} + b \sin^2 x \right) dx =$$

A.  $\frac{1}{b} - a \log|a \cos^2 x - b \sin^2 x| + c$

B.  $\frac{1}{b} - a \log|a \cos^2 x + b \sin^2 x| + c$

C.  $\frac{1}{b} + a \log|a \cos^2 x + b \sin^2 x| + c$

D.  $\frac{1}{b} + a \log|a \cos^2 x - b \sin^2 x| + c$

**Answer: B**



**View Text Solution**

$$73. \int \frac{3x - 4}{\sqrt{2x^2 + 4x + 5}} dx =$$

A.  $\frac{3}{2} \sqrt{2x^2 + 4x + 5} - \frac{7}{\sqrt{2}} \sin^{-1} \left( \sqrt{2} \frac{x - 1}{\sqrt{3}} \right) + c$

$$\text{B. } \frac{3}{2}\sqrt{2x^2 + 4x + 5} + \frac{7}{\sqrt{2}}\sin^{-1}h\left(\sqrt{2}\frac{x+1}{\sqrt{3}}\right) + c$$

$$\text{C. } \frac{3}{2}\sqrt{2x^2 + 4x + 5} - \frac{7}{\sqrt{2}}\sin^{-1}h\left(\sqrt{2}\frac{x+1}{\sqrt{3}}\right) + c$$

$$\text{D. } \frac{3}{2}\sqrt{2x^2 + 4x + 5} + \frac{7}{\sqrt{2}}\sin^{-1}h\left(\sqrt{2}\frac{x-1}{\sqrt{3}}\right) + c$$

**Answer: C**



**Watch Video Solution**

$$74. \int \frac{1}{2\cos x + 3\sin x + 4} dx =$$

$$\text{A. } \frac{2}{\sqrt{3}}\tan^{-1}\left[\frac{1}{\sqrt{3}}\left(2\frac{\tan x}{2} + 3\right)\right] + c$$

$$\text{B. } \frac{2}{\sqrt{3}}\tan^{-1}\left[\frac{1}{\sqrt{3}}\left(2\frac{\tan x}{2} - 3\right)\right] + c$$

$$\text{C. } \frac{1}{\sqrt{3}}\tan^{-1}\left[\frac{1}{\sqrt{3}}\left(2\frac{\tan x}{2} + 3\right)\right] + c$$

D. None

**Answer: A**



**Watch Video Solution**

$$75. \int e^x \left( \frac{2 + \sin 2x}{1 + \cos 2x} \right) dx =$$

A.  $e^x \cos 2x + c$

B.  $e^x \cot x + c$

C.  $2e^x \sec^2 x + c$

D.  $e^x \tan x + c$

**Answer: D**



**Watch Video Solution**

$$76. \int_0^{\pi} \sin^3 x \cos^4 x dx =$$

A.  $\frac{2}{35}$

B.  $\frac{4}{35}$

C.  $\frac{4}{30}$

D.  $\frac{2}{30}$



**Answer: B**

 [Watch Video Solution](#)

77.  $\lim_{n \rightarrow \infty} \left[ \frac{1}{3n+1} + \frac{1}{3n+2} + \dots + \frac{1}{3n+n} \right] =$

A.  $\log\left(\frac{4}{3}\right)$

B.  $\log\left(\frac{1}{3}\right)$

C.  $\log\left(\frac{3}{2}\right)$

D. 0

**Answer: A**

 [Watch Video Solution](#)

78. The area of the region bounded by the curves

$y = |x - 1|$  and  $y = 3 - |x|$  is

A. 1

B. 2

C. 3

D. 4

**Answer: D**



**Watch Video Solution**

79. The degree of the differential equation  $\left[5 + \frac{d^2y}{dx^2}\right]^{\frac{3}{2}} = \frac{dy}{dx}$

A. 1

B. 2

C. 3

D. None

**Answer: C**



**Watch Video Solution**

80. The solution of  $\frac{dy}{dx} = \frac{x - 2y + 3}{2x - y + 5}$  is

A.  $x^2 + 4xy - y^2 - 6x + 10y = c$

B.  $x^2 + 4xy + y^2 - 6x - 10y = c$

C.  $x^2 - 4xy - y^2 - 6x + 10y = c$

D.  $x^2 - 4xy + y^2 + 6x - 10y = c$

**Answer: D**



**Watch Video Solution**

81. If  $f: R \rightarrow R, S: R \rightarrow R$  are defined by  $f(x) = 3x-4, g(x) = 5x-1$  then,

$(f \circ g^{-1})(2) =$

A.  $\frac{11}{5}$

B.  $-\frac{11}{5}$

C.  $\frac{1}{5}$

D.  $-\frac{1}{5}$

**Answer: B**

 [Watch Video Solution](#)

82. The range of  $y = 2x^2 + x + \frac{2}{2x^2 + x + 1}$  is

A.  $\left(11 - \frac{\sqrt{2}}{7}, 11 + \frac{\sqrt{2}}{7}\right)$

B.  $\left[11 - \frac{\sqrt{2}}{7}, 11 + \frac{\sqrt{2}}{7}\right]$

C.  $\left(-11 + \frac{\sqrt{2}}{7}, -11 - \frac{\sqrt{2}}{7}\right)$

D.  $\left[-11 + \frac{\sqrt{2}}{7}, -11 + \frac{\sqrt{2}}{7}\right]$

**Answer: B**

 [View Text Solution](#)

$$83. \sum \left( \frac{1^2 + 2^2 + 3^2 + \dots + n^2}{1 + 2 + 3 + \dots + n} \right)$$

A.  $n^2 + \frac{11}{12}n$

B.  $n^2 - 2\frac{n}{6}$

C.  $n^2 + 2\frac{n}{3}$

D. None

**Answer: C**



**Watch Video Solution**

$$84. \text{ If } A = \begin{bmatrix} 1 & 1 & 0 \\ 0 & 1 & 0 \\ 1 & 0 & 1 \end{bmatrix} \text{ then } A^3 =$$

A.  $\begin{bmatrix} 1 & 3 & 0 \\ 0 & 1 & 0 \\ 3 & 3 & 1 \end{bmatrix}$

B.  $\begin{bmatrix} 1 & 0 & 3 \\ 3 & 1 & 3 \\ 0 & 0 & 1 \end{bmatrix}$

C.  $\begin{bmatrix} 3 & 3 & 1 \\ 1 & 3 & 0 \\ 0 & 1 & 0 \end{bmatrix}$

D.  $\begin{bmatrix} 3 & 1 & 0 \\ 3 & 3 & 1 \\ 1 & 0 & 0 \end{bmatrix}$

**Answer: A**



**Watch Video Solution**

85. If  $\frac{1}{a} + \frac{1}{b} + \frac{1}{c} = 0$  then  $\begin{vmatrix} 1+a & 1 & 1 \\ 1 & 1+b & 1 \\ 1 & 1 & 1+c \end{vmatrix} =$

A.  $-abc$

B. 0

C.  $abc$

D. None

**Answer: C**



**Watch Video Solution**

86. If the system of equations  $\lambda x + 3y + z = 0$ ,  $4x + \lambda y + 3z = 0$ ,  $2x + 3y + \lambda z = 0$  has non-trivial solution, then  $\lambda =$

A. 6

B. 3

C. 2

D. None

**Answer: D**



[Watch Video Solution](#)

87. If  $z_1, z_2, z_3$  are collinear and  $z_3 - \frac{z_1}{z_2 - z_1}$  is purely real, then

$$\arg\left(z_3 - \frac{z_1}{z_2 - z_1}\right)$$

A. 0

B. 1

C. 2

D. 3

**Answer: A**



[View Text Solution](#)

88. If  $\omega, \omega^2$  are cube root of unity then,  $\frac{\omega}{1} + \omega^2 + \frac{\omega}{1} + \omega =$

A.  $-2\omega^2$

B.  $-2\omega$

C. 2

D. -2

**Answer: D**



[View Text Solution](#)



89. If  $z = 3 + 3i$  then,  $z^2 + z + 15 =$

A.  $12+3i$

B.  $12-3i$

C.  $-12 - 3i$

D.  $-12 + 3i$

**Answer: A**



[View Text Solution](#)

90. If the area of the triangle on the complex plane formed by the points  $z$ ,  $iz$  and  $z+iz$  is 50 sq. units then  $|z|$  is

A. 15

B. 10

C. 5

D. None

**Answer: B**



[View Text Solution](#)

91. If  $A, B$  and  $C$  are the angles of a triangle such that  $\cos A + \cos B + \cos C = 0 = \sin A + \sin B + \sin C$ , then  $\sin 3A + \sin 3B + \sin 3C =$

A. 1

B. 2

C. 0

D. 3

**Answer: C**



[View Text Solution](#)

92. If  $x^2 + 3x - 2 = 0$ ,  $x^2 + 6x + k = 0$  have a common root then  $p =$

A. 10 (or) 16

B. 11 (or) 15

C. 8 (or) 4

D. 5 (or) 8

**Answer: D**

 [View Text Solution](#)

93. If  $x$  is real, then the maximum value of  $\frac{x^2 + 14x + 9}{x^2 + 2x + 3}$  is

A. 6

B. 8

C. 4

D. 2

**Answer: C**

 [Watch Video Solution](#)

94. The condition that the roots of  $x^3 + 3px^2 + 3qx + r = 0$  may be in

A.P is

A.  $2q^3 + r^2 = 3pqr$

B.  $2p^3 + r = 3pq$

C.  $p^3r = q^3$

D. None

**Answer: B**



**Watch Video Solution**

95. The roots of  $2x^5 + x^4 - 12x^3 - 12x^2 + x + 2 = 0$  are

A.  $-1, -2, -\frac{1}{2}, 3 \pm \frac{\sqrt{5}}{2}$

B.  $2 \pm \sqrt{3}, 3 \pm \sqrt{2}$

C.  $2, \frac{1}{2}, 3, \frac{1}{3}, 1$

D.  $1, 2, \frac{1}{2}, 5 \pm \sqrt{11} \frac{i}{6}$

**Answer: A**

 [Watch Video Solution](#)

**96.** How many numbers can be made with digit 3,4,5,6,7,8 lying between 3000 and 4000 which are divisible by 5 without repetition

A. 120

B. 60

C. 24

D. 12

**Answer: D**

 [View Text Solution](#)

97. If  ${}^n P_r = 840$ ,  ${}^n C_r = 35$  then  $n =$

A. 1

B. 7

C. 4

D. 10

**Answer: B**



[Watch Video Solution](#)

98. The co-efficient of  $x^{11}$  in the expansion of  $(1 + 3x + 2x^2)^6$  is

A. 216

B. 144

C. 576

D. 288

**Answer: C**



**Watch Video Solution**

99.  $1 + \frac{1}{3}x + \frac{1.4}{3.6}x^2 + \frac{1.4.7}{3.6.9}x^3 + \dots =$

A.  $(1 + x)^{\frac{1}{3}}$

B.  $(1 - x)^{\frac{1}{3}}$

C.  $(1 - x)^{-\frac{1}{3}}$

D.  $x$

**Answer: C**



**Watch Video Solution**

100. The partial fractions of  $\frac{1}{x^3(x + 2)} =$

A.  $\frac{1}{8}x + \frac{1}{4}x^2 + \frac{1}{2}x^3 - \frac{1}{8(x + 2)}$

B.  $\frac{1}{8}x - \frac{1}{4}x^2 + \frac{1}{2}x^3 - \frac{1}{8(x+2)}$

C.  $\frac{1}{8}x + \frac{1}{4}x^2 + \frac{1}{2}x^3 + \frac{1}{8(x+2)}$

D.  $\frac{1}{8}x - \frac{1}{4}x^2 - \frac{1}{2}x^3 + \frac{1}{8(x+2)}$

**Answer: B**



**Watch Video Solution**

**101.** If  $7 \sin^2 \theta + 3 \cos^2 \theta = 4$ , then  $\tan \theta$

A.  $\pm \frac{1}{\sqrt{3}}$

B.  $\frac{1}{3}$

C.  $\pm \sqrt{3}$

D.  $\pm 1$

**Answer: A**



**View Text Solution**



102. If  $\begin{vmatrix} 1 + \sin^2 \theta & \cos^2 \theta & 4 \sin 4\theta \\ \sin^2 \theta & 1 + \cos^2 \theta & 4 \sin 4\theta \\ \sin^2 \theta & \cos^2 \theta & 1 + 4 \sin 4\theta \end{vmatrix} = 0$ , then the value of  $\theta$  is

A.  $7\frac{\pi}{24}$  or  $11\frac{\pi}{24}$

B.  $\frac{\pi}{24}$

C.  $5\frac{\pi}{24}$

D. None

**Answer: A**



**Watch Video Solution**

103.  $\tan \left[ \frac{1}{2} \sin^{-1} \frac{2a}{1+a^2} + \frac{1}{2} \cos^{-1} \frac{1-a^2}{1+a^2} \right] =$

A.  $1 - \frac{a^2}{1+a^2}$

B.  $2\frac{a}{1+a^2}$

C.  $\frac{1+a^2}{1-a^2}$

D.  $2\frac{a}{1} - a^2$

**Answer: D**



**Watch Video Solution**

**104.** If  $\sin x \cosh y = \cos \theta$ ,  $\cos x \sinh y = \sin \theta$  then  $\sinh^2 y =$

A.  $\cosh^2 y$

B.  $\cos^2 x$

C.  $\sec^2 x$

D.  $\cosh^2 x$

**Answer: B**



**Watch Video Solution**

**105.**  $m \cdot (\tan \theta - 30) = n(\tan \theta + 120)$  then,  $m + \frac{n}{m} - n =$

A.  $\sin 2\theta$

B.  $\cos 2\theta$

C.  $2 \sin 2\theta$

D.  $2 \cos 2\theta$

**Answer: D**



[View Text Solution](#)

106. If  $\tan\left(B - \frac{C}{2}\right) = x \frac{\cot A}{2}$ , then  $x =$

A.  $a - \frac{b}{a + b}$

B.  $b - \frac{c}{b + c}$

C.  $c - \frac{a}{c + a}$

D. None

**Answer: B**



[View Text Solution](#)

107. If  $P_1, P_2, P_3$  are altitudes of  $\triangle ABC$  from the vertices A,B,C and  $\Delta$  is the area of triangle then,  $\frac{1}{P_1^2} + \frac{1}{P_2^2} + \frac{1}{P_3^2} =$

A.  $a^2 + b^2 + \frac{c^2}{4}\Delta^2$

B.  $a^2 - b^2 - \frac{c^2}{\Delta^2}$

C.  $a + b + \frac{c}{\Delta}$

D.  $a^2 + b^2 + \frac{c^2}{\Delta^2}$

**Answer: A**



[View Text Solution](#)

108. From the top of a mast of 60 m height, the angle of depression of an object is  $45^\circ$ . The distance of the object from the ship is

A. 45 m

B. 80 m

C. 60 m

D. 90 m

**Answer: C**



[Watch Video Solution](#)

**109.** If the position vector of A,B,C are  $2i + 3j + 4k, i + 2j, j + 2k$  and

$$\vec{AB} = P\vec{AC} \text{ then } P =$$

A.  $-\frac{1}{2}$

B.  $\frac{1}{4}$

C.  $\frac{1}{2}$

D.  $-\frac{1}{4}$

**Answer: C**



[View Text Solution](#)

110. If the position vectors of A, B are  $2a - 3b$ ,  $3a + 2b$  respectively then the position of vector of C in AB produced such that  $AC = 2 AB$  is

A.  $4a+7b$

B.  $5b-2a$

C.  $3a+2b$

D.  $3b-2a$

**Answer: A**



[Watch Video Solution](#)

111. The vector equation of the line passing through the point  $i+j-k$  and parallel to the vector  $2i+3j-k$  is

A.  $r=(i+j-k)+t(2i+3j-k)$

B.  $r=(i-j+k)+t(2i-3j-k)$

C.  $r=(i-j-k)+t(2i-3j-k)$

$$D. r = (-i - j + k) + t(-2i + 3j + k)$$

**Answer: A**



[Watch Video Solution](#)

**112.** The length of projection of  $i + 2j + 3k$  in the direction of  $3i - 4j + 5k$  is

A.  $\sqrt{2}$

B.  $\sqrt{3}$

C. 1

D. 2

**Answer: A**



[Watch Video Solution](#)

113. A unit vector perpendicular to each of the vector  $3i+2j+4k$  and  $2i+j-k$  is,

A.  $\pm \frac{-6i + 11j - k}{\sqrt{158}}$

B.  $\pm 6i - 8j + \frac{k}{\sqrt{104}}$

C.  $\pm 6i - 8j - \frac{k}{\sqrt{101}}$

D.  $\pm 6i - 8j - 4\frac{k}{\sqrt{101}}$

**Answer: A**



[Watch Video Solution](#)

114. The vector  $i+j+k, i+2j+3k, 2i+3j+k$  are

A. Collinear

B. Non-coplanar

C. Coplanar

D. None



**Answer: B**



[View Text Solution](#)

**115.** The variance of 6,5,8,10,3,4,9,11 is

A. 8

B. 9

C. 7.5

D. 10

**Answer: C**



[Watch Video Solution](#)

**116.** If 6 cards are drawn at random, from a pack of cards, then the probability to get 3 red and 3 black cards is

A.  $\frac{{}^{28}C_3 \times {}^{28}C_3}{{}^{56}C_6}$

B. None

C.  $\frac{{}^{16}C_3 \times {}^{16}C_3}{{}^{32}C_6}$

D.  $\frac{{}^{26}C_3 \times {}^{26}C_3}{{}^{52}C_6}$

**Answer: D**



**Watch Video Solution**

117. Let  $A, B, C$  are three events such that  $P(A) = 0.2, P(B) = 0.5, P(C) = 0.6, P(A \cap B) = 0.15, P(A \cap C) = 0.2$ , then,

A.  $0.05 \leq P(B \cap C) \leq 0.5$

B.  $0.03 \leq P(B \cap C) \leq 0.38$

C.  $0.06 \leq P(B \cap C) \leq 0.24$

D.  $0.04 \leq P(B \cap C) \leq 0.22$

**Answer: C**



[View Text Solution](#)

**118.** In a class, 40% students study botany, 25% Zoology and 15% bith Botany and Zoology. A student from the class is selected at random. The probability that he studies Botany, if it is known that he studies Zoology is

A.  $\frac{1}{8}$

B.  $\frac{2}{5}$

C.  $\frac{3}{8}$

D.  $\frac{3}{5}$

**Answer: D**



[View Text Solution](#)

119. There are 2 white , 4 black balls in urn A, In urn B, there are 5 white and 7 black balls. If one ball is randomly replaced from A and B , and a ball is drawn from B then the probability for the ball to be white one is

A.  $\frac{17}{40}$

B.  $\frac{14}{40}$

C.  $\frac{19}{45}$

D.  $\frac{16}{39}$

**Answer: D**



[Watch Video Solution](#)

120. If X is random variable with distribution given below

x: 1 2 3 4

P(X=x): k k 2k 3k

The value of k and its mean are

A.  $\frac{3}{19}, \frac{8}{19}$

B.  $\frac{1}{19}, \frac{21}{19}$

C.  $\frac{4}{19}, \frac{18}{19}$

D.  $\frac{5}{19}, \frac{15}{19}$

**Answer: B**



**View Text Solution**

**121.** If  $X$  is a poisson distribution such that  $P(X=1)=P(X=2)$  then,  $P(X=4)=$

A.  $\frac{1}{3}e^2$

B.  $\frac{2}{3}e^2$

C.  $\frac{4}{3}e^2$

D. none

**Answer: B**



**Watch Video Solution**

122. A(2, 1) and B(2, 3) are two points. If P is a point such that  $PA + PB = 2$ , then the locus of P is

A.  $4x^2 - 12y^2 - 16x + 124y - 69 = 0$

B.  $4x^2 + 12y^2 - 16x - 124y + 69 = 0$

C.  $4x^2 + 12y^2 + 16x - 124y + 69 = 0$

D.  $4x^2 + 12y^2 + 16x + 124y - 69 = 0$

**Answer: A**



[View Text Solution](#)

123. The transformed equation of  $xy + 2x - 5y - 11 = 0$  when the origin is shifted to the point (2, 3) is,

A.  $xy - 5x - 3y + 16 = 0$

B.  $xy + 5x + 3y - 16 = 0$

C.  $xy+5x-3y - 16 = 0$

D.  $xy-5x+3y + 16 = 0$

**Answer: C**



**Watch Video Solution**

**124.** The area of the triangle formed by the line  $\frac{x}{5} + \frac{y}{4} = 1$  with the coordinate axes is

A. 20 sq.units

B. 15 sq.units

C. 5 sq.units

D. 10 sq.units

**Answer: D**



**Watch Video Solution**

**125.** The perpendicular distance of the straight line  $3x + 4y - 8 = 0$  from the point of intersection of the lines  $3x + 2y + 4 = 0$ ,  $2x + 5y - 1 = 0$  is

A.  $\frac{11}{5}$  units

B.  $\frac{12}{5}$  units

C.  $\frac{8}{5}$  units

D.  $\frac{9}{5}$  units

**Answer: D**



[View Text Solution](#)

**126.** The diagonal of a square is  $8x - 15y = 0$  and one vertex of the square is  $(1, 2)$ . The equations to the sides of the square passing through this vertex are

A.  $23x + 7y = 9$ ,  $7x - 23y = 52$

B.  $23x + 7y = 9$ ,  $7x - 23y = 53$



C.  $22x + 8y = 9$ ,  $22x - 8y = 52$

D. None

**Answer: B**



[View Text Solution](#)

**127.** The area of the triangle formed by the pair of lines  $3x^2 + 8xy - 3y^2 = 0$  and the line  $3x + 4y - 5 = 0$  is,

A.  $\frac{3}{5}$  sq. units

B.  $\frac{5}{3}$  sq. units

C.  $\frac{4}{5}$  sq. units

D.  $\frac{5}{4}$  sq. units

**Answer: B**



[Watch Video Solution](#)

128. If  $x^2 - 10xy + 4y^2 + 6x + 2y + k = 0$  represents a pair of straight lines then,  $k =$

A.  $-\frac{1}{3}$

B.  $\frac{2}{3}$

C.  $-\frac{2}{3}$

D. None

**Answer: A**



[View Text Solution](#)

129. The equation of the circle concentric with  $x^2 + y^2 - 2x + 8y - 23 = 0$  and passing through  $(2, 3)$  is

A.  $x^2 + y^2 - 2x - 8y - 33 = 0$

B.  $x^2 + y^2 + 6x - 4y - 12 = 0$

C.  $x^2 + y^2 + x + 8y + 33 = 0$

$$D. x^2 + y^2 - 6x + 4y - 12 = 0$$

**Answer: A**



[Watch Video Solution](#)

**130.** If the tangent to the circle  $x^2 + y^2 = 5$  at  $(1, -2)$  also touches the circle  $x^2 + y^2 - 8x + 6y + 20 = 0$  then the point of contact is

A.  $(-1, 0)$

B.  $(1, 0)$

C.  $(3, -1)$

D.  $(5, 2)$ .

**Answer: C**



[Watch Video Solution](#)

131. If the circles  $x^2 + y^2 - 6x - 8y + c = 0$  and  $x^2 + y^2 = 9$  have three common tangent then  $c =$

A. 17

B. 19

C. 21

D. 20

**Answer: C**



[Watch Video Solution](#)

132. The equation of the circle which cuts orthogonally the circle  $x^2 + y^2 - 4x + 2y - 7 = 0$  and having centre at  $(2, 3)$  is,

A.  $3x^2 - 3y^2 + 2x - 12y + 2 = 0$

B.  $x^2 + y^2 + 6x + 4y + 19 = 0$

C.  $x^2 + y^2 - 2x + 4y + 2 = 0$

$$D. x^2 + y^2 - 4x - 6y + 9 = 0$$

**Answer: D**



[View Text Solution](#)

**133.** If  $(2, 1)$  is limiting point of coaxial system of which  $x^2 + y^2 - 6x - 4y - 3 = 0$  is a member, then the other limiting point is

A.  $(-5,-6)$

B.  $(-2,-3)$

C.  $(3,2)$

D.  $(5,6)$

**Answer: A**



[View Text Solution](#)

134. The locus of the point of intersection of tangents to parabola  $y^2 = 4(x + 1)$  and  $y^2 = 8(x + 2)$  which are perpendicular to each other is

A.  $x-3=0$

B.  $x+3=0$

C.  $y-3=0$

D.  $y+3=0$

**Answer: B**



[View Text Solution](#)

135. If the normals at the points  $t_1$  and  $t_2$  on  $y^2 = 4ax$  at the point  $t_3$  on the parabola, the  $t_1 t_2 =$

A. 4

B. 3

C. 2

D.  $2t_3$

**Answer: C**

 [View Text Solution](#)

**136.** The eccentricity of the ellipse  $9x^2 + 16y^2 = 144$  is

A.  $\frac{7}{4}$

B.  $\frac{7}{2}$

C.  $\frac{\sqrt{7}}{2}$

D.  $\frac{\sqrt{7}}{4}$

**Answer: D**

 [View Text Solution](#)

137. The condition that the line  $x \cos \alpha + y \sin \alpha = P$  may be a normal to the ellipse  $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$  is

A.  $\frac{a^2}{\cos^2 \alpha} - \frac{b^2}{\sin^2 \alpha} = \left(a^2 - \frac{b^2}{P^2}\right)^2$

B.  $\frac{a^2}{\cos^2 \alpha} - \frac{b^2}{\sin^2 \alpha} = \left(a^2 + \frac{b^2}{P^2}\right)^2$

C.  $\frac{a^2}{\cos^2 \alpha} + \frac{b^2}{\sin^2 \alpha} = \left(a^2 - \frac{b^2}{P^2}\right)^2$

D.  $\frac{a^2}{\cos^2 \alpha} + \frac{b^2}{\sin^2 \alpha} = \left(a^2 + \frac{b^2}{P^2}\right)^2$

**Answer: C**



[View Text Solution](#)

138. The distance between the foci is  $4\sqrt{13}$  and the length of conjugate axis is 8 then, the eccentricity of the hyperbola is

A.  $\frac{\sqrt{15}}{4}$

B.  $\frac{\sqrt{13}}{4}$

C.  $\frac{\sqrt{13}}{3}$



D.  $\frac{\sqrt{13}}{2}$

**Answer: C**



[View Text Solution](#)

**139.** If the d.c.'s ( $l, m, n$ ) of two lines are connected by the relations  $l + m + n = 0$  and  $2mn + 3ln - 5lm = 0$  then the angle between the lines is

A.  $\frac{\pi}{2}$

B.  $\frac{\pi}{6}$

C.  $\frac{\pi}{3}$

D. None

**Answer: A**



[Watch Video Solution](#)

140. The plane  $2x + 3y + kz - 7 = 0$  is parallel to the line whose d.r's are  $(2,3,-1)$  then  $k =$

- A. 5
- B. 10
- C. 15
- D. 20

**Answer: B**

 [View Text Solution](#)

141. The centroid of the triangle formed by the points  $(1, 2, 3)$ ,  $(2, 3, 1)$ ,  $(3, 1, 2)$  is

- A.  $(1,1,1)$
- B.  $(2,2,2)$
- C.  $(1,2,2)$

D. (3,1,3)

**Answer: B**



[Watch Video Solution](#)

142.  $\lim_{x \rightarrow 0} \left( \frac{1 - 2 \cos x + \cos 2x}{x^2} \right)$

A. 2

B. 3

C. -1

D. 0

**Answer: C**



[Watch Video Solution](#)

143.  $\lim_{x \rightarrow 2^-} \{x + (x - [x])^2\} =$

A. 3

B. 1

C. 2

D. None

**Answer: A**



**Watch Video Solution**

**144.**

If

$$f(x) = \begin{cases} -2 \sin x, & \text{or } x \leq -\frac{\pi}{2} \\ a \sin x + b, & \text{or } -\frac{\pi}{2} < x < \frac{\pi}{2} \end{cases}$$

everywhere then the ordered pair(a,b) is

A. (-1,1)

B. (1,1)

C. (0,0)

D. None

**Answer: A**



**View Text Solution**

145. If  $y = x \log |x + \sqrt{1 + x^2}| - \sqrt{1 + x^2}$  then  $\frac{dy}{dx} =$

A.  $\cos^{-1} hx$

B.  $\frac{1}{2} \log(x - \sqrt{1 + x^2})$

C.  $\cos ec^{-1} hx$

D.  $\sin^{-1} hx$

**Answer: D**



**Watch Video Solution**

146. The derivative of  $e^{\sin^{-1} x}$  w.r.t  $\log x$  is

A.  $\frac{e^{\sin^{-1} x}}{\sqrt{1 + x^2}}$

- B.  $x \frac{e^{\sin^{-1} x}}{\sqrt{1+x^2}}$
- C.  $\frac{e^{\sin^{-1} x}}{\sqrt{1+x^2}}$
- D.  $x \frac{e^{\sin^{-1} x}}{\sqrt{1-x^2}}$

**Answer: D**



**Watch Video Solution**

147. If  $y = ae^{nx} + be^{-nx}$  then  $y_2 =$

- A.  $-n^2y$
- B.  $ny$
- C.  $n^2y$
- D.  $-ny$

**Answer: C**



**Watch Video Solution**

148. The equation of the tangent to the curve  $y^2 = 4ax$  at  $(at^2, 2at)$  is

A.  $xt + y - 2at - at^3 = 0$

B.  $x + yt = at^2$

C.  $x - yt + at^2 = 0$

D.  $xt - y - 2at - at^3 = 0$

**Answer: C**



[View Text Solution](#)

149. The side of a square is equal to the diameter of a circle. If the side and radius change at the same rate then the ratio of the change of their areas is

A.  $2 : \pi$

B.  $1 : 1$

C.  $\pi : 2$

D.  $1 : \pi$

**Answer: A**



[View Text Solution](#)

150. Verify Rolle's theorem for the function  $f(x) = x(x + 3)e^{-\frac{x}{2}}$  in  $[-3,0]$ .

A. -2

B. 1

C. -1

D. 0

**Answer: A**



[Watch Video Solution](#)



151. Show that when the curved surface of a right circular cylinder inscribed in a sphere of radius  $R$  is maximum, then the height of the cylinder is  $\sqrt{2R}$ .

A.  $\sqrt{3}R$

B.  $\sqrt{2}R$

C.  $2R$

D.  $R$

**Answer: B**



**Watch Video Solution**

152.  $\int \left( \sin 2 \frac{x}{a \cos^2 x} + b \sin^2 x \right) dx =$

A.  $\frac{1}{b} - a \log |a \cos^2 x - b \sin^2 x| + c$

B.  $\frac{1}{b} - a \log |a \cos^2 x + b \sin^2 x| + c$

C.  $\frac{1}{b} + a \log |a \cos^2 x + b \sin^2 x| + c$

$$D. \frac{1}{b} + a \log|a \cos^2 x - b \sin^2 x| + c$$

**Answer: B**

 [View Text Solution](#)

$$153. \int \frac{3x - 4}{\sqrt{2x^2 + 4x + 5}} dx =$$

$$A. \frac{3}{2} \sqrt{2x^2 + 4x + 5} - \frac{7}{\sqrt{2}} \sin^{-1} \left( \sqrt{2} \frac{x - 1}{\sqrt{3}} \right) + c$$

$$B. \frac{3}{2} \sqrt{2x^2 + 4x + 5} + \frac{7}{\sqrt{2}} \sin^{-1} h \left( \sqrt{2} \frac{x + 1}{\sqrt{3}} \right) + c$$

$$C. \frac{3}{2} \sqrt{2x^2 + 4x + 5} - \frac{7}{\sqrt{2}} \sin^{-1} h \left( \sqrt{2} \frac{x + 1}{\sqrt{3}} \right) + c$$

$$D. \frac{3}{2} \sqrt{2x^2 + 4x + 5} + \frac{7}{\sqrt{2}} \sin^{-1} h \left( \sqrt{2} \frac{x - 1}{\sqrt{3}} \right) + c$$

**Answer: C**

 [Watch Video Solution](#)

$$154. \int \frac{1}{2 \cos x + 3 \sin x + 4} dx =$$

- A.  $\frac{2}{\sqrt{3}} \tan^{-1} \left[ \frac{1}{\sqrt{3}} \left( 2 \frac{\tan x}{2} + 3 \right) \right] + c$
- B.  $\frac{2}{\sqrt{3}} \tan^{-1} \left[ \frac{1}{\sqrt{3}} \left( 2 \frac{\tan x}{2} - 3 \right) \right] + c$
- C.  $\frac{1}{\sqrt{3}} \tan^{-1} \left[ \frac{1}{\sqrt{3}} \left( 2 \frac{\tan x}{2} + 3 \right) \right] + c$

D. None

**Answer: A**



**Watch Video Solution**

155.  $\int e^x \left( \frac{2 + \sin 2x}{1 + \cos 2x} \right) dx =$

A.  $e^x \cos 2x + c$

B.  $e^x \cot x + c$

C.  $2e^x \sec^2 x + c$

D.  $e^x \tan x + c$

**Answer: D**



**Watch Video Solution**

156.  $\int_0^{\pi} \sin^3 x \cos^4 x dx =$

A.  $\frac{2}{35}$

B.  $\frac{4}{35}$

C.  $\frac{4}{30}$

D.  $\frac{2}{30}$

**Answer: B**



**Watch Video Solution**

157.  $Lt_{n \rightarrow \infty} \left[ \frac{1}{3n+1} + \frac{1}{3n+2} + \dots + \frac{1}{3n+n} \right] =$

A.  $\log\left(\frac{4}{3}\right)$

B.  $\log\left(\frac{1}{3}\right)$

C.  $\log\left(\frac{3}{2}\right)$

D. 0

**Answer: A**



[Watch Video Solution](#)

**158.** The area of the region bounded by the curves  $y = |x - 1|$  and  $y = 3 - |x|$  is

A. 1

B. 2

C. 3

D. 4

**Answer: D**



[Watch Video Solution](#)

159. The degree of the differential equation  $\left[5 + \frac{d^2y}{dx^2}\right]^{\frac{3}{2}} = \frac{dy}{dx}$

- A. 1
- B. 2
- C. 3
- D. None

**Answer: C**



[Watch Video Solution](#)

160. The solution of  $\frac{dy}{dx} = \frac{x - 2y + 3}{2x - y + 5}$  is

- A.  $x^2 + 4xy - y^2 - 6x + 10y = c$
- B.  $x^2 + 4xy + y^2 - 6x - 10y = c$
- C.  $x^2 - 4xy - y^2 - 6x + 10y = c$
- D.  $x^2 - 4xy + y^2 + 6x - 10y = c$

**Answer: D**



**Watch Video Solution**

## Physics

1. When a big drop of water is formed from  $n$  small drops of water, the energy loss is  $3E$ , where,  $E$  is the energy of the bigger drop. If the radius of the bigger drop is  $R$  and  $r$  is the radius of the smaller drop, then number of smaller drops ( $n$ ) is

A.  $4\frac{R}{r^2}$

B.  $4\frac{R}{r}$

C.  $2\frac{R^2}{r}$

D.  $4\frac{R^2}{r^2}$

**Answer: C**



**Watch Video Solution**

2. Two litre glass flask contains some mercury. It is found that at all temperatures the volume of the air inside the flask remains the same. The volume of mercury inside the flask is

$$\left(\alpha_g = 9 \times 10^{-6} \text{ } ^\circ\text{C}^{-1} \gamma_{Hg} = 1.8 \times 10^{-4} \text{ } ^\circ\text{C}^{-1}\right)$$

A. 1500 CC

B. 150 CC

C. 300 CC

D. 3000 CC

**Answer: B**

 [View Text Solution](#)

3. Two photons of energies twice and thrice the work function of a metal are incident on the metal surface. Then, the ratio of maximum velocities of the photoelectrons emitted in the two cases respectively, is



A.  $\sqrt{2}: 1$

B.  $\sqrt{3}: 3$

C.  $\sqrt{3}: \sqrt{2}$

D.  $1: \sqrt{2}$

**Answer: D**



**Watch Video Solution**

4. A gas is compressed at a constant pressure of  $50N/m^2$ , from a volume  $10m^3$  to a volume of  $4m^3$ . 100 J of heat is added to the gas then its internal energy.

A. Increases by 400 J

B. Increases by 200 J

C. Decreases by 400 J

D. Decreases by 200 J

**Answer: A**



**View Text Solution**

5. An ammeter whose resistance is  $180\Omega$  shows full scale deflection when the current is 2 mA. The shunt required to convert into an ammeter reading 20mA is (in ohm )

A. 18

B. 20

C. 0.1

D. 10

**Answer: D**



**Watch Video Solution**

6. Two bodies of mass 4 kg and 5 kg are moving along East and North directions with velocities 5 m/s and 3 m/s respectively. Magnitude of the velocity of centre of mass of the system is

A.  $\frac{25}{9} \text{ m/s}$

B.  $\frac{9}{25} \text{ m/s}$

C.  $\frac{41}{9} \text{ m/s}$

D.  $\frac{16}{9} \text{ m/s}$

**Answer: A**



[Watch Video Solution](#)

7. The frequency of vibration in a vibration magnetometer of the combination of two bar magnets of magnetic moments  $M_1$  and  $M_2$  is 6 Hz when like poles are tied and it is 2 Hz when the unlike poles are tied together, then the ratio  $M_1 : M_2$  is

A. 4:5

B. 5:4

C. 1:3

D. 3:1

**Answer: B**



[Watch Video Solution](#)

8. A short magnetic needle is pivoted in a uniform magnetic field of induction  $IT$ . Now, simultaneously another magnetic field of induction  $\sqrt{3}T$  is applied at right angles to the first field, the needle deflects through an angle  $\theta$  whose value is

A.  $30^\circ$

B.  $45^\circ$

C.  $90^\circ$

D.  $60^\circ$

**Answer: A**



[Watch Video Solution](#)

9. Two litre glass flask contains some mercury. It is found that at all temperatures the volume of the air inside the flask remains the same. The volume of mercury inside the flask is

$$\left( a_g = 9 \times 10^{-6} \text{ } ^\circ\text{C}^{-1} \gamma_{Hg} = 1.8 \times 10^{-4} \text{ } ^\circ\text{C}^{-1} \right)$$

A. 1500 CC

B. 150 CC

C. 300 CC

D. 3000 CC

**Answer: B**



[View Text Solution](#)

10. A mass  $kg$  is suspended by a weightless string. The horizontal force required to hold the mass at  $60^\circ$  with the vertical is

A.  $Mg$

B.  $Mg\sqrt{3}$

C.  $Mg(\sqrt{3} + 1)$

D.  $M\frac{g}{\sqrt{3}}$

**Answer: B**



[View Text Solution](#)

11. If the equation of motion of a projectile is  $y = 3x - \frac{1}{8}x^2$ , the range and maximum height are respectively ( $y$  and  $x$  are in metres).

A. 18 m and 24 m

B. 24 m and 18 m

C. 24 m and 6 m

D. 12 m and 9 m

**Answer: A**



[View Text Solution](#)

12. If  $n_e$  and  $n_h$  are electron and hole concentrations in an extrinsic semiconductor and  $n_i$  is electron concentration in an intrinsic semiconductor then.

A.  $\left(\frac{n_e}{n_h}\right) = n_i$

B.  $(n_e + n_h) = n_i$

C.  $(n_e - n_h) = n_i^2$

D.  $(n_e n_h) = n_i^2$

**Answer: A**



[View Text Solution](#)

13. A radioactive nucleus can decay by two different processes. The half lives of the first and second decay processes are  $5 \times 10^3$  and  $10^5$  years respectively, Then, the effective half-life of the nucleus is,

A.  $105 \times 10^5$

B. 4762 yrs

C. 104 yrs

D. 47.6 yrs


**Answer: A**



[Watch Video Solution](#)

14. Which logic gate is represented by the following combination of logic gates





- A. OR
- B. NAND
- C. AND
- D. NOR

**Answer: A**

[View Text Solution](#)

15. The FM radio broadcasting band is,

- A. 5 MHz to 30 MHz
- B. 88 MHz to 108 MHz

C. 30 KHz to 300 KHz

D. 3 GHz to 30 GHz

**Answer: B**



[View Text Solution](#)

## Chemistry

1. A metallic carbide on treatment with water gives a colourless gas which burns readily in air and gives a precipitate with ammoniacal silver nitrate solution. The gas evolved

A.  $CH_4$

B.  $C_2H_6$

C.  $C_2H_4$

D.  $C_2H_2$

**Answer: A**

 [Watch Video Solution](#)

2. Chlorination of toluene in presence of light and heat followed by treatment with aqueous NaOH gives

A. o-Cresol

B. p-Cresol

C. 2: 4 dihydroxy toluene

D. Benzoic acid

**Answer: B**

 [Watch Video Solution](#)

3. Molar ionic conductivities of a bivalent electrolyte are 57 and 73. The molar conductivity of the solution will be

A.  $130 \text{ Scm}^2 \text{ mol}^{-1}$

B.  $65 \text{ Scm}^2 \text{ mol}^{-1}$

C.  $260 \text{ Scm}^2 \text{ mol}^{-1}$

D.  $187 \text{ Scm}^2 \text{ mol}^{-1}$

**Answer: C**



**Watch Video Solution**

4. The forces operating between the adsorbate and the adsorbent in physical adsorption are

A. Van der Waals forces

B. Chemical forces

C. Covalent forces

D. All the three

**Answer: A**

 [Watch Video Solution](#)

5. Horn silver is

- A. Carbonate mineral
- B. Chloride mineral
- C. Sulphate mineral
- D. Phosphate mineral

**Answer: B**

 [Watch Video Solution](#)

6. Stainless steel does not rust because

- A. Chromium and nickel combine with iron
- B. Chromium forms an oxide layer and protects iron from rusting
- C. Nickel present in it does not rust.

D. Iron forms a hard chemical compound with chromium present in it

**Answer: B**



[Watch Video Solution](#)

7. A polymer which is commonly used as a packaging material is

A. Polythene

B. Polypropylene

C. PVC





D. Bakelite

**Answer: D**



[Watch Video Solution](#)

8. Which compound/set of compounds is used in the manufacture of Nylon-6,6?



Answer: D



Watch Video Solution

9. The pH value of a solution in which a polar amino acid does not migrate under the influence of electric field is called

A. Isoelectronic point

B. Iso-electric point

C. Neutralisation point

D. None

**Answer: B**



[Watch Video Solution](#)

10. The substances which affect the central nervous system and induce sleep are called

- A. Antipyretics
- B. Tranquilizers
- C. Analgesics
- D. Antibiotics



**Answer: A**



[Watch Video Solution](#)



11. Which is detected by carbylamine test?



D. All of these

**Answer: B**



[Watch Video Solution](#)

12. Acid catalyzed hydration of alkenes except ethene leads to the formation of

A. Primary alcohol

B. Secondary or tertiary alcohol

C. Mixture of primary and secondary alcohols




D. Mixture of secondary and tertiary alcohols

**Answer: D**

 [Watch Video Solution](#)

13. 2, 2-dichloro propane treated with aq.KOH gives an unstable product.

It is



**Answer: B**

 [Watch Video Solution](#)

14. The process that does not yield an amine is

- A. Action of ammonia on RX
- B. Reduction of aldoxime with Na/alcohol
- C. Acid hydrolysis of alkyl cyanide
- D. Reduction of amide with  $LiAlH_4$

**Answer: D**

 [Watch Video Solution](#)

**15.** Which one of the following compounds would undergo nitration with greatest ease?

- A. Benzene
- B. Phenol
- C. Nitrobenzene
- D. Benzoic acid

**Answer: B**

 [Watch Video Solution](#)

16. In the chemical reaction,  $CH_3CH_2NH_2 + CHCl_3 + 3KOH \rightarrow (A) + (B) + 3H_2O$  the compounds (A) and (B) are respectively

- A.  $C_2H_5NC$  and  $3KCl$
- B.  $C_2H_5CN$  and  $3KCl$
- C.  $CH_3CH_2CONH_2$  and  $3KCl$
- D.  $C_2H_5NC$  and  $K_2CO_3$

**Answer: B**

 [Watch Video Solution](#)

17. 'X' grams of calcium carbonate was completely burnt in air. The weight of the solid residue formed is 28 g. What is the value of 'X'(in grams)?

- A. 50
- B. 150
- C. 300
- D. 200

**Answer: D**

 [Watch Video Solution](#)

**18.** 4 g of a hydrocarbon on complete combustion gave 12.571 g of  $\text{CO}_2$  and 5.143 g of water. What is the empirical formula of the hydrocarbon?

- A.  $\text{CH}_2$
- B.  $\text{C}_3\text{H}_3$
- C.  $\text{C}_2\text{H}_3$
- D.  $\text{C}_2\text{H}_4$

**Answer: B**

 [Watch Video Solution](#)

19. 10 grams of  $CaCO_3$  is completely decomposed to x and CaO. 'x' is passed into an aqueous solution containing 0.1mole of sodium carbonate.

What is the number of moles of sodium bicarbonate formed? (mol. wts:

$CaCO_3 = 100, Na_2CO_3 = 106, NaHCO_3 = 84$ )

A. 0.2

B. 10

C. 0.3

D. 5

**Answer: A**

 [Watch Video Solution](#)

20. 50 grams of calcium carbonate was completely burnt in air. What is the weight (in grams) of the residue? Atomic weights of Ca, C and O are

40,12 and 16 respectively)

A. 2.8

B. 28

C. 22

D. 4.4

**Answer: B**



[Watch Video Solution](#)

21. 4 g of a hydrocarbon on complete combustion gave 12.571 g of  $\text{CO}_2$  and 5.143 g of water. What is the empirical formula of the hydrocarbon?

A.  $\text{CH}_2$

B.  $\text{C}_3\text{H}_3$

C.  $\text{C}_2\text{H}_3$

D.  $\text{C}_2\text{H}_4$

**Answer: B**



**Watch Video Solution**