



MATHS

BOOKS - SAI MATHS (TELUGU ENGLISH)

QUESTION PAPER

Math

1. The domain of function $f(x) = \frac{1}{\sqrt{x^2 - [x]} - 2}$ is (Here $[x]$ denotes the greatest integral not exceeding the value of x)

A. $(-\infty, -2) \cup (1, \infty)$

B. $(-\infty, -2) \cup (0, \infty)$

C. $(-\infty, 2) \cup (2, \infty)$

D. $(-\infty, -1) \cup (3, \infty)$

Answer: D



[View Text Solution](#)

2. Let $f: R \rightarrow T$ and $g: R \rightarrow R$ be the function defined by

$$f(x) = \frac{x}{1+x^2}, x \in R, g(x) = \frac{x^2}{1+x^2}, x \in R$$

Then, the correct statement (s) among the following is/are:

- (A) both f, g are one-one
- (B) Both f, g are onto
- (C) both f, g are not one-one as well as not onto
- (D) f and g onto but not one-one

A. A

B. A,B

C. D

D. C

Answer: D



[View Text Solution](#)

3.

If

 $\alpha \in R, n \in N$ and $n + 2(n - 1) + 3(n - 2) + \dots + (n - 1)2 + n.1 =$

A. $\frac{1}{2}$

B. $\frac{1}{3}$

C. $\frac{1}{5}$

D. $\frac{1}{6}$

Answer: D[Watch Video Solution](#)

4. If the function $f \left\{ [a, b] \rightarrow \left[-\frac{\sqrt{3}}{4}, \frac{1}{2} \right] \right\}$ defined by

$$f(x) = \begin{bmatrix} 1 & 1 & 1 \\ 1 & 1 + \sin x & 1 \\ 1 + \cos x & 1 & 1 \end{bmatrix} \text{ is one-one onto, then}$$

A. $a = \frac{-\pi}{4}, b = \frac{\pi}{6}$

B. $a = \frac{-\pi}{2}, b = \frac{\pi}{2}$

C. $a = \frac{-\pi}{6}, b = \frac{\pi}{4}$

D. $a = -\pi, b = \pi$

Answer: A

 [View Text Solution](#)

5. The rank of the matrix $\begin{vmatrix} 3 & 5 & -1 & 4 \\ 2 & 1 & 3 & -2 \\ 8 & 11 & 1 & 6 \\ -7 & 14 & 6 & -14 \end{vmatrix}$ is

A. 1

B. 2

C. 3

D. 4

Answer: B



[View Text Solution](#)

6. A value of e for which the following system of equations has a non-trivial solution is

$$(4 \sin \theta)x - 3y + z = 0, \quad -(6 \cos 2\theta)y + z = 0, \quad 3x - 12y + 4z = 0$$

A. $\tan^{-1}\left(\frac{1}{2}\right)$

B. $\frac{\pi}{4}$

C. $\sin^{-1}\left(\frac{3}{16}\right)$

D. $\frac{\pi}{12}$

Answer: C



[Watch Video Solution](#)

7. If $z = \sqrt{2}\sqrt{1 + \sqrt{3}i}$ represents a point P in the argand plane and P lies in the third quadrant, then the polar form of z is

A. $\left[\cos\left(\frac{-4\pi}{3}\right) + I \sin\left(\frac{-4\pi}{3}\right) \right]$

B. $2 \left[\cos\left(\frac{-5\pi}{6}\right) + I \sin\left(\frac{-5\pi}{6}\right) \right]$

C. $2 \left[\cos\left(\frac{-\pi}{6}\right) + I \sin\left(\frac{-\pi}{6}\right) \right]$

D. $2 \left[\cos\left(\frac{-2\pi}{3}\right) + I \sin\left(\frac{-2\pi}{3}\right) \right]$

Answer: B



Watch Video Solution

8. If $z = x + iy$ represents a point P in the argand plane, then the area of the region represented by the inequality $2 < |z - (1+i)| < 3$ is

A. 49π

B. 36π

C. 25π

D. 5π

Answer: D

 [Watch Video Solution](#)

9. If p is complex number whose modulus is one, then the equation

$$\left(\frac{1+iz}{1-iz}\right)^4 = p \text{ has}$$

- A. real and equal roots
- B. real and distinct roots
- C. two real and two complex roots
- D. all complex roots

Answer: B

 [View Text Solution](#)

10.
$$\sum_{r=1}^{16} \left(\sin \left[\frac{2r\pi}{17} \right] + i \cos \left[\frac{2r\pi}{17} \right] \right) =$$

- A. 1

B. -1

C. i

D. $-I$

Answer: D

 [Watch Video Solution](#)

11. If the maximum value of $2x - 7 - ax^2$ cannot exceed 20, then the minimum value of a is

A. 27

B. $\frac{1}{13}$

C. 13

D. $\frac{1}{27}$

Answer: D

 [Watch Video Solution](#)

12. A student while solving a quadratic equation in x , he copied its constant term incorrectly and got its roots as 5 and 9. Another student copied the constant term and coefficient of x^2 of the same equation correctly as 12 and 4 respectively. If s , p and Δ denote respectively the sum of the roots, the product of the roots and discriminant of the correct equation, then, $\frac{\Delta}{3p + s} =$

A. 48

B. 45

C. 8

D. 16

Answer: C



View Text Solution

13. The product of the real roots of the equation

$$(x + 1)^4 + (x + 3)^4 = 8 \text{ is}$$

A. 0

B. 74

C. $7 - 2\sqrt{3}$

D. $7 + 2\sqrt{3}$

Answer: C



[Watch Video Solution](#)

14. The sum of all the real numbers satisfying the equation

$$x^2 + |x - 3| = 4 \text{ is}$$

A. 0

B. 1

C. 2

Answer: B



Watch Video Solution

15. Consider the following statements:

(i) Number of one-one functions from set A to B, where

$O(A) = m$ and $O(B) = n (m \leq n)$ is given by ${}^n P_m$,

(ii) Number of ways in which 'n' people can be arranged at a circular table

is $\frac{(n-1)!}{2}$.

(iii) Number of ways of selecting atleast one thing out of the given n

distinct things is $2^n - 1$.

(iv) Number of ways in which n distinguishable objects can be distributed

into k distinguishable bins is ${}^n C_{k-1}$.

A. All the statements are true

B. All except (iii) are true

C. Only (i) and (iii) are true

D. Only (ii) is false

Answer: C



[View Text Solution](#)

16. The number of ways in which 3 identical balls can be distributed into 7 distinct bins is

A. 49

B. 84

C. 35

D. 42

Answer: B



[Watch Video Solution](#)

17. For $x = \frac{5}{7}$, If t_k is the first negative term in the expansion of $(1+x)^{7/5}$, then $t_1 + t_2 + \dots + t_k =$

- A. $\frac{13}{7}$
- B. $\frac{107}{14}$
- C. $\frac{104}{49}$
- D. $\frac{921}{28}$

Answer: C



[Watch Video Solution](#)

18. The coefficient of x^6 in the tower series expansion of $\frac{x^4 - 12x^2 + 7}{(x^2 + 1)^3}$ is

- A. 149
- B. -253
- C. -145

Answer: C**Watch Video Solution****19.** Show that

$$\left(1 + \cos. \frac{\pi}{8}\right) \left(1 + \cos. \frac{3\pi}{8}\right) \left(1 + \cos. \frac{5\pi}{8}\right) \left(1 + \cos. \frac{7\pi}{8}\right) = \frac{1}{8}$$

A. $\frac{1 + \sqrt{2}}{2\sqrt{2}}$

B. $\frac{\pi}{8}$

C. $\frac{1}{8}$

D. $\frac{1}{2}$

Answer: C**Watch Video Solution**

20. If A and B are acute angles satisfying

$$3 \cos^2 A + 2 \cos^2 B = 4 \quad \text{and} \quad \frac{3 \sin A}{\sin B} = \frac{2 \cos B}{\cos A},$$

then $A + 2B =$

A. 30°

B. 45°

C. 60°

D. 90°

Answer: D



Watch Video Solution

21.

If

$$A + B + C = \frac{\pi}{3}, \text{ then}$$

$$\sin\left(\frac{\pi - 6A}{6}\right) + \sin\left(\frac{\pi - 6B}{6}\right) + \sin C =$$

A. $-1 + 4 \cos\left(\frac{\pi - 6A}{12}\right) \sin\left(\left(\frac{\pi - 6B}{12}\right) \sin \frac{C}{2}\right)$

B. $4 \sin\left(\frac{\pi + 6A}{12}\right) \sin\left(\frac{\pi + 6B}{12}\right) \cos \frac{C}{2}$

$$\text{C. } 1 - 4 \cos\left(\frac{\pi - 6A}{12}\right) \cos\left(\frac{\pi - 6B}{13}\right) \cos\left(\frac{\pi - 6C}{12}\right)$$

$$\text{D. } 4 \cos\left(\frac{\pi - 6A}{12}\right) \cos\left(\frac{\pi - 6B}{12}\right) \sin \frac{C}{2}$$

Answer: D



View Text Solution

22. If $\sqrt{2} \sin^2 x + (3\sqrt{2} + 1) \sin x + 3 > 0$ and $x^2 - 7x + 10 < 0$, then

x lies in the interval

A. $\left(\frac{-\pi}{4}, \frac{3\pi}{4}\right)$

B. $\left(2, \frac{5\pi}{4}\right)$

C. $\left(0, \frac{3\pi}{2}\right)$

D. $\left(\frac{5\pi}{4}, 5\right)$

Answer: B



Watch Video Solution

23. If $\cos^{-1} x + \cos^{-1} y + \cos^{-1} z = 3\pi$, then

A. $x + y + z - 3 = 0$

B. $x + y + z + 3 = 0$

C. $x + 2y + 3z - 5 = 0$

D. $x - y - z = 0$

Answer: B



Watch Video Solution

24. If $\sinh^{-1}(\sqrt{8}) + \sinh^{-1}(\sqrt{24}) = \alpha$, then $\sinh \alpha =$

A. $6\sqrt{6} - 10\sqrt{2}$

B. $6\sqrt{6} + 10\sqrt{2}$

C. $16\sqrt{6}$

D. $16\sqrt{6} + 4\sqrt{2}$

Answer: B



Watch Video Solution

25. In a triangle ABC, if $\cos A \cos B + \sin A \sin B + \sin C =$

A. $\frac{2 + \sqrt{3}}{2}$

B. $1 + \sqrt{2}$

C. $(2\sqrt{3} - 1)(2)$

D. $\frac{3 + \sqrt{3}}{2}$

Answer: B



View Text Solution

26. In a $\triangle ABC$, if the medians AD and BE are such that $AD = 4$,

$\angle DAB = \frac{\pi}{6}$ and $\angle ABE = \frac{\pi}{3}$ then the area of $\triangle ABC$ (in square

units) is

A. $\frac{16}{3\sqrt{3}}$

B. $\frac{48}{3\sqrt{3}}$

C. $\frac{64}{3\sqrt{3}}$

D. $\frac{32}{3\sqrt{3}}$

Answer: D



Watch Video Solution

27. In a $\triangle ABC$, if $r_1 = 2r_2 = 3r_3$, then $a : b =$

A. 3 : 5

B. 5 : 3

C. 4 : 5

D. 5 : 4

Answer: D



View Text Solution

28. If \bar{a} and \bar{b} are respectively the internal and external bisectors of the angles between the vectors $-\bar{i} + 2\bar{j} - 2\bar{k}$ and $3\bar{i} + 4\bar{j}$ and $|\bar{a}| = \frac{2}{3}\sqrt{6}$, $|\bar{b}| = \frac{2}{3}\sqrt{3}$, then one of the values of $\bar{a} - \bar{b}$ is,

A. $\frac{1}{10}(-8\bar{i} + 11\bar{j} - 2\bar{k})$

B. $\frac{2}{3}(-\bar{i} + 2\bar{j} - 2\bar{k})$

C. $\frac{1}{15}(9\bar{i} - 11\bar{j} + 3\bar{k})$

D. $\frac{1}{12}(2\bar{i} - 3\bar{j} - \bar{k})$

Answer: B



[View Text Solution](#)

29. If the angle between the vectors $2\alpha^2\bar{i} + 4\alpha\bar{j} + \bar{k}$ and $7\bar{i} - 2\bar{j} + \alpha\bar{k}$ is obtuse, then

A. $\alpha > \frac{1}{2}$

B. $0 < \alpha < \frac{1}{2}$

C. $\alpha < 0$

D. $-|\alpha| < \frac{1}{2}$

Answer: B



View Text Solution

30. Let L be the line parallel to the vector $\sqrt{2}\bar{i} - 5\bar{j} + 3\bar{k}$ and passing through the point A given by $\bar{i} + 2\bar{j} - 3\bar{k}$. If the distance between A and a point P on the line L is 18 units, then the position vector of such a point P is

A. $(1 - 3\sqrt{2})\bar{i} + 17\bar{j} - 12\bar{k}$

B. $(1 + 3\sqrt{2})\bar{i} + 17\bar{j} + 12\bar{k}$

C. $(1 + 3\sqrt{2})\bar{i} - 17\bar{j} - 12\bar{k}$

D. $(1 - 3\sqrt{2})\bar{i} - 17\bar{j} + 12\bar{k}$

Answer: A



[View Text Solution](#)

31. Let $\bar{a} = p(\bar{i} + \bar{j} + \bar{k})$, $\bar{b} = \bar{i} + \bar{j} - 2\bar{k}$ and $\bar{c} = 2\bar{i} - \bar{j} + 2\bar{k}$ be three vectors. If the values of $[\bar{a}\bar{b}\bar{c}]$ is not more than 15 and not less than -5 , then p lies in the interval.

A. $\left[\frac{-5}{3}, \frac{5}{9} \right]$

B. $\left(\frac{-5}{9}, \frac{5}{9} \right)$

C. $\left(0, \frac{5}{9} \right)$

D. $\left(\frac{-5}{3}, \frac{5}{9} \right)$

Answer: D



[View Text Solution](#)

32. Let $a = \hat{i} + 2\hat{j} - \hat{k}$ and $b = \hat{i} + \hat{j} + \hat{k}$. If p is a unit vector such that $[abp]$ is maximum then $p =$

A. $\frac{1}{\sqrt{6}}(\bar{i} - 2\bar{j} + \bar{k})$

B. $\frac{1}{\sqrt{3}}(\bar{i} + \bar{j} - \bar{k})$

C. $\frac{1}{\sqrt{14}}(3\bar{i} - 2\bar{j} - \bar{k})$

D. $\frac{1}{\sqrt{14}}(\bar{i} + 2\bar{j} + 3\bar{k})$

Answer: C



Watch Video Solution

33. If $\bar{a} = 2\bar{i} - 3\bar{j} + \bar{k}$, $\bar{b} = \bar{i} - \bar{j} + 2\bar{k}$ and $\bar{c} = 2\bar{i} + \bar{j} + \bar{k}$ are three vectors, then $|(\bar{a} \times \bar{b}) \times \bar{c}| =$

A. $|\bar{a} \times (\bar{b} \times \bar{c})|$

B. $\frac{\sqrt{39}}{\sqrt{11}}|\bar{a} \times (\bar{b} \times \bar{c})|$

C. $\sqrt{\frac{11}{39}}|\bar{a} \times (\bar{b} \times \bar{c})|$

D. $\sqrt{11}|\bar{a} \times (\bar{b} \times \bar{c})|$

Answer: C



[View Text Solution](#)

34. For a group of 100 observations, the arithmetic mean and standard deviation are 8 and $\sqrt{10.5}$ respectively. The mean and standard deviation of 50 items selected from these 100 observations are 10 and 2 respectively. Then the standard deviation of the remaining 50 observation is

A. 2

B. 3

C. 3.5

D. 4

Answer: B



[Watch Video Solution](#)

35. If a number is selected from the first 30 natural numbers, then the probability that the number selected is divisible by 4 or 7, is

A. $\frac{4}{3}$

B. $\frac{1}{5}$

C. $\frac{7}{30}$

D. $\frac{1}{3}$

Answer: D



[Watch Video Solution](#)

36. If 80% of flights depart on time, 70% of flights arrive on time and 65% of flights depart on time and arrive on time, then the probability that a flight that has just departed on time will arrive on time is

A. $\frac{13}{16}$

B. $\frac{11}{16}$

C. $\frac{13}{14}$

D. $\frac{11}{14}$

Answer: A



Watch Video Solution

37. A computer program has two modules X and Y and errors in them occur independently. X has an error with probability 0.1 and Y has an error with probability 0.3. If an error in X alone cause the program to crash with probability 0.5, an error in Y alone causes the program to crash with probability 0.7 and an error in both X and Y cause the program to crash with probability 0.8, then the probability that the program is crashed is

A. $\frac{23}{125}$

B. $\frac{26}{125}$

C. $\frac{29}{125}$

D. $\frac{31}{125}$

Answer: D



Watch Video Solution

38. Two dice are rolled. If a random variable X denotes the sum of the numbers on them and μ denotes the mean of X , then

$$\mu + P(X < 5) + P(X > 9) + P(x = 7) =$$

A. $\frac{15}{2}$

B. 17

C. $\frac{17}{2}$

D. 15

Answer: A



Watch Video Solution

39. A boy rolled a die once. If an even number appear on then the number of chocolates the boy gets is equal to two more then the number appeared. If an odd number appear on that die. Then the number of chocolates he gets is equal to three more than the number'appeared. If a random variable X represents the number of chocolates the boy receive, then the range of X is

A. $\{4, 6, 8\}$

B. $\{3, 5, 7\}$

C. $\{3, 4, 7\}$

D. $\{2, 3\}$

Answer: A



[Watch Video Solution](#)

40. When the origin is shifted to $(1, -2)$ by translation of coordinate axes, the transformed coordinates of $(3, -2)$ are (α, β) If the axes are rotated

about origin through an angle of 45° after the translation, then the transformed coordinates of (α, β) are

A. $(\sqrt{2}, 0)$

B. $(0, \sqrt{2})$

C. $(-\sqrt{2}, \sqrt{2})$

D. $(\sqrt{2}, -\sqrt{2})$

Answer: D



Watch Video Solution

41. If a straight line is passing through the point of intersection of the lines $3x - 4y + 1 = 0$, $5x + y - 1 = 0$ and making equal non-zero intercepts on the coordinate axes, then the area (in sq. units) of the triangle formed by this line with the coordinate axes, is

A. $\frac{121}{1058}$

B. $\frac{121}{529}$

C. $\frac{529}{121}$

D. $\frac{1058}{121}$

Answer: A



Watch Video Solution

42. If the equations of the perpendicular bisectors of the sides AB and AC of a $\triangle ABC$ are $x - y + 5 = 0$ and $x + 2y = 0$ respectively and if A is $(1, -2)$, then the equation of the perpendicular bisector of the side BC is,

A. $3x + 3y + 5 = 0$

B. $9x - 3y + 40 = 0$

C. $6x + 15 = 5$

D. $23x - 14y + 100 = 0$

Answer: D



Watch Video Solution

43. If the lines $x + 3y - 5 = 0$, $5x + 2y - 12 = 0$ and $3x - ky - 1 = 0$ do not form a triangle, then a value of k is

A. $\frac{1}{5}$

B. $\frac{-1}{5}$

C. $\frac{-6}{5}$

D. $\frac{6}{5}$

Answer: C

[Watch Video Solution](#)

44. If the combined equation of the diagonals of the square formed by the pairs of lines $xy + 4x - 5y - 20 = 0$ and $xy - 5x + 4y - 20 = 0$ is $x^2 - y^2 - kx + ly = 0$, then $k+l =$

A. 0

B. 2

C. -1

D. 1

Answer: B



Watch Video Solution

45. The equation of the pair of bisectors of the angles between the pair of straight lines

A. $7(3x + 5)^2 - 2(3x + 5)(y + 5) - 7(y + 5)^2 = 0$

B. $7(3x - 5)^2 - 2(3x - 5)(y - 5) - 7(y - 5)^2 = 0$

C. $7(5x + 3)^2 - 2(5x + 3)(5y + 1) - 7(5y + 1)^2 = 0$

D. $7(5x - 3)^2 - 2(5x - 3)(5y - 1) - 7(5y - 1)^2 = 0$

Answer: C



View Text Solution

46. If $x = \frac{2at}{1+t^2}$, $y = \frac{a(1-t^2)}{1+t^2}$, where t is a parameter, then a is

- A. the length of the latus rectum of a parabola
- B. the radius of a circle
- C. the length of the transverse axis of a hyperbola
- D. the length of semi-major axis of an ellipse

Answer: B



[Watch Video Solution](#)

47. Consider the following statement :

I. $P(x_1, y_1)$ and $Q(x_2, y_2)$ are conjugate points with respect to the circle

$$x^2 + y^2 + 2gx + 2fy + c = 0 \text{ then}$$

$$x_1x_2 + y_1y_2 + g(x_1 + x_2) + f(y_1 + y_2) + c = 0$$

II. The pole of the line $x + y + 1 = 0$ with respect to the circle $x^2 + y^2 = 9$ is

(9.9).

Then, which one of the following is true?

- A. Both I and II are true
- B. Neither I nor II is true
- C. I is false and II is true
- D. I is true and II is false

Answer: D



[Watch Video Solution](#)

48. If P is the point of contact of the circles $x^2 + y^2 + 4x + 4y - 10 = 0$ and $x^2 + y^2 - 6x - 6y + 10 = 0$ and Q is their external centre of similitude, then the equation of the circle with P and Q as the extremities of its diameter is

- A. $x^2 + y^2 + 14x + 14y - 26 = 0$
- B. $x^2 + y^2 + 5x + 5y - 8 = 0$

$$C. x^2 + y^2 - 5x - 5y + 8 = 0$$

$$D. x^2 + y^2 - 14x - 14y + 26 = 0$$

Answer: C::D



[View Text Solution](#)

49. The equation of the circle touching the line $2x + 3y + 1 = 0$ at the point $(1, -1)$ and orthogonal to the circle with has the line segment having end points $(0, -1)$ and $(-1, 3)$ as diameter, is

$$A. x^2 + y^2 - 10x + 5y + 1 = 0$$

$$B. x^2 + y^2 + 5x - 10y - 1 = 0$$

$$C. 2x^2 + 2y^2 + 10x - 5y - 1 = 0$$

$$D. 2x^2 + 2y^2 - 10x - 5y + 1 = 0$$

Answer: A



[View Text Solution](#)

50. The angle between the tangents drawn from the point (1,4) to the parabola $y^2 = 4x$ is

A. $\frac{\pi}{3}$

B. $\frac{\pi}{2}$

C. $\frac{3\pi}{4}$

D. π

Answer: D



Watch Video Solution

51. If the line $y = -x + k$ is normal to the curve $y^2 = 16x$, then k is

A. 21

B. 14

C. 13

D. 12

Answer: D



Watch Video Solution

52. The normal drawn at the point

$\left(\sqrt{9} \cos\left(\frac{\pi}{4}\right), \sqrt{7} \sin\left(\frac{\pi}{4}\right)\right)$ to the ellipse $\frac{x^2}{9} + \frac{y^2}{7} = 1$

intersects its major axis at the point

A. $\left(0, \sqrt{\frac{2}{7}}\right)$

B. $\left(-\sqrt{\frac{2}{9}}, 0\right)$

C. $\left(0 - \sqrt{\frac{2}{7}}\right)$

D. $\left(\sqrt{\frac{2}{9}}, 0\right)$

Answer: C



Watch Video Solution

53. If $(1, -2)$ is the focus and $x + y - 2 = 0$ is the directrix of the ellipse $17x^2 - 2xy + 17y^2 - 32x + 76y + 86 = 0$, then its eccentricity is

- A. $\frac{1}{5}$
- B. $\frac{1}{2}$
- C. $\frac{1}{3}$
- D. $\frac{1}{4}$

Answer: B



[View Text Solution](#)

54. The line $2x + y = 1$ is a tangent to the hyperbola $\frac{x^2}{a^2} - \frac{y^2}{b^2}$ ($a > b$). If this line passes through the point of intersection of a directrix and the positive X-axis, then the eccentricity of that hyperbola is

A. $\sqrt{2}$

B. 2

C. $\sqrt{3}$

D. 3

Answer: A



[View Text Solution](#)

55. If the orthocentre and the centroid of a triangle are at $(5, 2, -6)$ and $(9, 6, -4)$ respectively, then its circumcentre is

A. $(11, 8, -3)$

B. $(8, 8, -3)$

C. $(11, 8, 3)$

D. $(11, -8, -3)$

Answer: A

 [Watch Video Solution](#)

56. If the direction ratios of two lines are given by $a + 2b + c = 0$ and $11bc + 6ca - 14ab = 0$. then the angle between these lines is

A. $\frac{\pi}{3}$

B. $\cos^{-1}\left(\frac{1}{3}\right)$

C. $\cos^{-1}\left(\frac{2}{3}\right)$

D. $\frac{\pi}{2}$

Answer: D

 [Watch Video Solution](#)

57. A plane cuts the coordinate axes X,Y,Z at A,B,C respectively such that the centroid of the ΔABC is (6,6,3). Then the equation of that plane is

A. $x + y + z = 18$

B. $x + 2y + z = 18$

C. $x + y + 2z = 18$

D. $2x + y + z = 18$

Answer: C

 [Watch Video Solution](#)

58. $\lim_{x \rightarrow \infty} x \left(\log \left(1 + \frac{x}{2} \right) - \log \frac{x}{2} \right) =$

A. 0

B. 1

C. 2

D. e

Answer: C

 [Watch Video Solution](#)

59. The set of values of x for which the function $f(x) = \log\left(\frac{x-1}{x+2}\right)$ is continuous, is

A. \mathbb{R}

B. $(-\infty, -1) \cup (0, \infty)$

C. $(-\infty, -2) \cup (1, \infty)$

D. $(-2, -1)$

Answer: C



Watch Video Solution

60. If the function $f: \mathbb{R} \rightarrow \mathbb{R}$, defined by

$$f(x) = \begin{cases} 5 - 3x & , \text{if } x \leq \frac{5}{3} \\ x^2 - 3x + 20, & \text{if } x > \frac{5}{3} \end{cases}, \text{ then 'f' is}$$

A. continuous at $x = \frac{5}{3}$

B. differentiable at $x = \frac{5}{3}$

C. differentiable at $x = 2$

D. discontinuous at $x = -2$

Answer: C

 [Watch Video Solution](#)

61. If t is a parameter and $x = t + \frac{1}{t}$, $y = t - \frac{1}{t}$, then $\frac{d^2y}{dx^2} =$

A. $\frac{4t}{(t^2 + 1)^2}$

B. $\frac{t^2}{t^2 + 1}$

C. $\frac{4t^3}{(t^2 + 1)^3}$

D. $\frac{-4t^3}{(t^2 - 1)^3}$

Answer: D

 [Watch Video Solution](#)

62. If $ax^2 + 2hxy + by^2 = 3$, then $\frac{d^2y}{dx^2} =$

A. $\frac{(hx^2 + by + ax)}{(ax + hy)^2}$

B. $\frac{(axy + hx^2 + byx)}{(ax + by)^2}$

C. $\frac{2(h^2 - ab)}{(hx + by)^3}$

D. $\frac{(ab + h)^2}{(ax + hy)^2} [h(x^2 + y^2) + xy(a + b)]$

Answer: C



[View Text Solution](#)

63. If the error in measuring the side l of an equilateral triangle is 0.01, then the percentage error in the area of the triangle, in terms of its side l is

A. $\frac{2}{l}$

B. $\frac{3}{l}$

C. $\frac{4}{l}$

D. $\frac{6}{l}$

Answer: A



[View Text Solution](#)

64. If at any point (x_1, y_1) on the curve $y = f(x)$, the lengths of the subtangent and subnormal are equal, then the length of the tangent drawn to that curve at that point is

A. $2|y_1|$

B. $\sqrt{2}|y_1|$

C. $\sqrt{5}|y_1|$

D. $\sqrt{2} \left| \frac{y_1}{x_2} \right|$

Answer: B



[View Text Solution](#)

65. The height of a right circular cylinder is decreasing while its diameter is increasing at a rate of 4 cm/sec so as to keep its volume unchanged. The rate of change in its lateral surface area (in $cm^2/sec.$) at the instant when its diameter is 8 cm and height is 12 cm, is

A. 24π

B. -24π

C. 48π

D. -48π

Answer: D



[View Text Solution](#)

66. If $x + y = 60$, $x > 0$, $y > 0$, then the maximum value of xy^3 is

A. $(15)^4 \frac{25}{3}$

B. $45(15)^3$

C. $\frac{(45)^3 9}{5}$

D. $\frac{(45)^4}{3}$

Answer: D

 [View Text Solution](#)

67. $\int \sqrt{1 + 2 \cot x (\cot x + \operatorname{cosec} x)} dx =$

A. $2 \log \left| \sin \frac{x}{2} \right| + c$

B. $2 \log \left| \cos \frac{x}{2} \right| + c$

C. $\log \left| \sin \frac{x}{2} + \cos \frac{x}{2} \right| + c$

D. $2 \log |\sin x + \cos x| + c$

Answer: A

 [View Text Solution](#)

68. If $\int e^{ax} \left(\frac{1 - \beta \sin x}{1 - \cos x} \right) dx = -e^x \cot \frac{x}{2} + c$, then $\frac{\alpha^2 + \beta^2}{2\alpha\beta} =$

A. -1

B. 1

C. 2

D. -2

Answer: B

 [View Text Solution](#)

69. $\int \frac{2x^3 - 4x^2 - x - 3}{x^2 - 2x - 3} dx =$

A. $\frac{7}{2} \log|x - 1| + \frac{3}{2} \log|x + 3| + c$

B. $2 \log|x - 1| + \frac{7}{2} \log|x - 3| + c$

C. $2x + \frac{1}{2} \log|x + 1| + \frac{3}{4} \log|x - 3| + c$

D. $x^2 + 2 \log|x + 1| + 3 \log|x - 3| + c$

Answer: D



View Text Solution

70. $\int \cos e^x \cot^3 x dx =$

A. $\frac{\cos e^x \cot^3 x}{4} - \frac{5}{8} \cos e^x \cot x + \frac{3}{8} \log \left| \tan \frac{x}{2} \right| + c$

B. $-\frac{\cos e^x \cot^3 x}{4} - \frac{5}{8} \cos e^x \cot x + \frac{3}{8} \log \left| \tan \frac{x}{2} \right| + c$

C. $-\frac{\cos e^x \cot x}{4} - \frac{3}{8} \cos e^x \cot x + \frac{3}{8} \log \left| \tan \frac{x}{2} \right| + c$

D. $-\frac{\cos e^x \cot x}{4} - \frac{3}{8} \cos e^x \cot x - \frac{3}{8} \log \left| \tan \frac{x}{2} \right| + c$

Answer: C



View Text Solution

71. $\int_{\frac{1}{3}}^3 \frac{1}{x} \sin \left(\frac{1}{x} - x \right) dx =$

A. 0

B. $\frac{4}{3}$

C. $\frac{5}{3}$

D. $\frac{8}{3}$

Answer: A

 [View Text Solution](#)

72. $\int^{\frac{3}{2}} -(-1)|x \sin(\pi x)| dx =$

A. $\frac{1}{\pi} - \frac{1}{\pi^2}$

B. $\frac{2}{\pi} + \frac{1}{\pi^2}$

C. $\frac{3}{\pi} - \frac{1}{\pi^2}$

D. $\frac{3}{\pi} + \frac{1}{\pi^2}$

Answer: D

 [View Text Solution](#)

73. Area of the region (in square units) bounded by the curve $y = x^2 + 4$ and the line $y = 5x - 2$ is

A. $\frac{1}{2}$

B. $\frac{1}{12}$

C. 2

D. $\frac{1}{6}$

Answer: D



[View Text Solution](#)

74. If m and n are the order and degree of the differential equation of the family of parabolas with focus at the origin and X-axis as its axis, then mn

$$-m + n =$$

A. 1

B. 2

C. 3

D. 4

Answer: C



[View Text Solution](#)

75. The solution of the differential equation

$$\frac{dy}{dx} + \frac{x}{y} \cdot \frac{x^2 + y^2 - 1}{2(x^2 + y^2) + 1} = 0 \text{ is}$$

A. $x^2 + y^2 + 3 \log(x^2 + y^2) = c$

B. $x^2 2y^2 - 3 \log(x^2 + y^2 + 2) = c$

C. $x^2 + y^2 - 3 \log(x^2 + y^2 + 2) = c$

D. $-x^2 - 2y^2 - 3 \log(x^2 + y^2) = c$

Answer: C



[View Text Solution](#)

76. If $x \log x \frac{dy}{dx} + y = \log x^2$ and $y(e) = 0$, then $y(e^2) =$

A. 0

B. 1

C. $\frac{1}{2}$

D. $\frac{3}{2}$

Answer: D

 [View Text Solution](#)

Physics

1. Identify the correct option.

A. The range of the gravitational force is large but not infinite.

B. The range of electromagnetic force is large but lesser than the range of gravitational force.

C. The range of weak nuclear force is smaller than the range of strong nuclear force, gravitation force and electron magnetic force.

D. The range of the weak as well as strong nuclear force is the order of 10^{-19} m

Answer: C

 [Watch Video Solution](#)

2. A current carrying conductor obeys Ohm's law ($V = RI$). If the current passing through the conductor is $I = (5 \pm 0.2)$ A and voltage developed is $V = (60 \pm 6)$ then find the percentage of error in resistance, R

A. 18

B. 6

C. 14

D. 2

Answer: C



[Watch Video Solution](#)

3. A particle covers a distance from A to B over a period of time, the distance versus time plot is shown below. Then which of the following is true for the motion of the particle ?



- A. Both average speed and instantaneous speed are always zero.
- B. Average speed is always non zero but average speed can be zero.
- C. Instantaneous speed is always non zero but average speed can be zero.
- D. Both average speed and instantaneous speed are always non zero.

Answer: B



[View Text Solution](#)

4. A boy standing on a moving truck throws a projectile such that he is able to catch it back after the truck has moved 100 m. If the truck is moving horizontally along a straight line with a constant speed 30 m/s, at what speed (relative to the truck) must the projectile is thrown. (Assume, $g = 10m / s^2$)

A. $\frac{55}{3}m / s$

B. $\frac{43}{2}m / s$

C. $\frac{50}{3}m / s$

D. $\frac{23}{2}m / s$

Answer: C::D



[Watch Video Solution](#)

5. In the pulley system shown in figure, the mass of A is half of that of rod B. The rod length is 500 cm. The mass of pulleys and the threads may be neglected. The mass A is set at the same level as the lower end of the rod

and then released. After releasing the mass A, it would reach the top end of the rod B in time (Assume $g = 10\text{ m/s}^2$)



A. 2.0 s

B. 1.0 s

C. 3.0 s

D. 4.0 s

Answer: B



[View Text Solution](#)

6. A light rigid wire of length 1 m is attached to a ball A of mass 500 g to one end. The other end of the wire is fixed, so that the wire can rotate freely in vertical plane about its fixed end. At the lowest point of the circular motion, the ball is given a horizontal velocity 6 m/s. Determine the radial component of the acceleration of the ball, when this rigid wire

makes an angle 60° with the upward vertical. (Assume $g = 10\text{ m/s}^2$)



A. 10 m/s^2

B. 6 m/s^2

C. 18 m/s^2

D. 25 m/s^2

Answer: D



[View Text Solution](#)

7. A block of mass 2 kg is connected to an ideal spring and is placed on a smooth horizontal surface. The spring is pulled to move the block and at an instant, the speed of end A of the spring and speed of the block were measured to be 6 m/s and 3 m/s respectively. At this moment the potential energy stored in the spring is increasing at a rate 5 J/s. Find the acceleration of the block at this instant.



A. $1.5m / s^2$

B. $3.0m / s^2$

C. $4.5m / s^2$

D. $2.5m / s^2$

Answer: D



View Text Solution

8. If the displacement of a body is given by $x = 3 \cos \left[2\pi t + \frac{\pi}{4} \right]$ m, then the acceleration of the body at $t=2s$ is

A. 0

B. $-6\sqrt{2}\pi^2 m / s^2$

C. $-10\pi^2 m / s^2$

D. $-12\sqrt{2}\pi^2 m / s^2$

Answer: B

 [Watch Video Solution](#)

9. If the acceleration due to gravity g doubles and the radius of earth becomes half that of the present value, then the value of escape velocity is (Assume, $g = 10m / s^2$ and radius of earth, $R = 6400$ km)

- A. $12km / s$
- B. $16\sqrt{2}km / s$
- C. $8\sqrt{2}km / s$
- D. $4\sqrt{2}km / s$

Answer: C

 [Watch Video Solution](#)

10. A uniform rod of length L is rotated in a horizontal plane about a vertical axis through one of its ends. The angular speed of rotation is ω .

Find increase in length of the rod, if ρ and Y are the density and Young's modulus of the rod respectively,

A. $\frac{\rho \omega \eta^3 Y}{4L^2}$

B. $\frac{4\rho\omega^2 L^3}{3Y}$

C. $\frac{\rho\omega^2 L^3}{3Y}$

D. $\frac{\rho\omega^3 L^3}{8Y}$

Answer: C



Watch Video Solution

11. A cylindrical vessel of height 50 cm is filled with water and rests on table. A small hole is made at the height 'h' from the bottom of the vessel so that the water jet could hit the table surface at a maximum distance x_{\max} from the vessel as shown in the figure. The value of x_{\max} will be (Neglect the viscosity of water)



A. 15 cm

B. 35 cm

C. 50 cm

D. 40 cm

Answer: C

 [View Text Solution](#)

12. A cubical block of wood, of length 10 cm, floats at the interface between oil of density $800\text{kg}/\text{m}^3$ and water. The lower surface of the block is 1.5cm below the interface. If the depth of water is 10 cm below the interface and oil is up to 10 cm above the interface then the difference in pressure at the lower and the upper face of the wooden block is (Assume density of water = $1000\text{kg}/\text{m}^3$ and $g = 10\text{m}/\text{s}^2$)



A. 850 Pa

B. 780 Pa

C. 800 Pa

D. 830 Pa

Answer: D



[View Text Solution](#)

13. A thermally insulated vessel containing monatomic gas is moving with a speed of 30 m/s. If the vessel suddenly stops, the increase in gas temperature is (Molar mass of gas = 83 g/mol and $R = 8.3\text{J/K mol}$)

A. 1 K

B. 3 K

C. 4 K

D. 6 K

Answer: B

[Watch Video Solution](#)

14. A string of length 100 cm has three resonant frequencies, 120 Hz, 200 Hz and 280 Hz. If a node is formed at the end of the string, the speed of the transverse wave on this string is:

- A. 60 m/s
- B. 80 m/s
- C. 100 m/s
- D. 120 m/s

Answer: B

[Watch Video Solution](#)

15. Two particles executing simple harmonic motion as described by $y_1 = 30 \sin\left(2\pi t + \frac{\pi}{3}\right)$ and $y_2 = 10(\sin 2\pi t + \sqrt{3} \cos 2\pi t)$ have amplitudes A_1 and A_2 respectively. The ratio $A_1 : A_2$ is

A. 2:1

B. 1:1

C. 3:1

D. $1:\sqrt{3}$

Answer: C



Watch Video Solution

16. Two lenses A and B having focal lengths 2.0 cm and 5.0 cm, respectively are placed 14 cm apart. Lens A is placed to the left of lens B. An object is placed 3 cm to the left of lens A. The distance of the image from lens A will be

A. $\frac{40}{3} \text{ cm}$

B. $\frac{82}{3} \text{ cm}$

C. $\frac{112}{5} \text{ cm}$

D. $\frac{92}{5} \text{ cm}$

Answer: B



Watch Video Solution

17. A spherical drop of liquid carrying charge, Q has potential V_0 at its surface. If two drops of same charge and radius combine to form of single spherical drop, then the potential at the surface of new drop is (Assume, $V = 0$ at infinity.)

A. $2^{1/3}V_0$

B. $4^{1/3}V_0$

C. $6^{1/3}V_0$

D. $2^{-1/3}V_0$

Answer: B



Watch Video Solution

18. Calculate the Voltage across AB terminals in the given circuit



A. $\frac{3}{8}V$

B. $\frac{8}{3}V$

C. $\frac{3}{2}V$

D. $\frac{2}{3}V$

Answer: B



[View Text Solution](#)

19. When subjected to a voltage of 10 V, the current through a resistor at a temperature of $40^\circ C$ is 0.1 A. The temperature coefficient of resistance of the material of the resistor is $2 \times 10^{-4}^\circ C$. The temperature of the resistor in $^\circ C$ when the current drops to 0.098 A is

A. 142

B. 167

C. 181

D. 206

Answer: A



Watch Video Solution

20. A small bar magnet experiences a torque of 0.016 Nm when placed with its axis at 30° with an external field of 0.04 T . If the bar magnet is replaced by a solenoid of cross-sectional area of 1 cm^2 and 1000 turns but having the same magnetic moment as that of bar magnet, then the current flowing through the solenoid is

A. 0.083

B. 0.16

C. 0.25

D. 0.3

Answer: A



[Watch Video Solution](#)

21. A rod of length 80 cm rotates about its mid point with a frequency of 10 rev/s. The potential difference in volts) between two ends of the rod due to a magnetic field, $B=0.5$ T directed perpendicular to the rod is

A. π

B. 1.6π

C. 2π

D. 0.8π

Answer: B



[Watch Video Solution](#)

22. LCR circuit, the resonance frequency of circuit increases two times of the initial circuit by changing C and C' and R from 100Ω to 400Ω while the inductance was kept the same. The ratio C/C' , is

A. 2

B. 8

C. 16

D. 4

Answer: D



[Watch Video Solution](#)

23. The half life of neutron is 693 seconds. What fraction of neutrons will decay when a beam of neutrons, having kinetic energy of 0.084 eV, travells a distance of 1 km? (mass of neutron 1.68×10^{-27} kg, and $\ln 2=0.693$)

A. 60×10^{-5}

B. 15×10^{-5}

C. 25×10^{-5}

D. 50×10^{-5}

Answer: C



Watch Video Solution

24. For a given truth table A,B,C are inputs and Y is the output, then the functional form of the circuit is



A. \bar{A}

B. \bar{B}

C. $\bar{A} + BC$

D. $A + B + C$

Answer: A



[View Text Solution](#)

25. Two diodes are connected in the following fashion. Provision is made to connect either $+5V$ or ground ($0V$) to the points A and B. The output Q will act as



- A. OR gate
- B. AND gate
- C. XOR gate
- D. NAND gate

Answer: C



[View Text Solution](#)

26. Carrier waves are used for communicating signals over long distance, because

- A. Carrier waves can be generated very easily
- B. Low frequencies can not be easily modulated by the carrier waves
- C. Low frequencies can be transmitted over long distnces
- D. Carrier waves that are generatea at higher frequencies can be transmitted over long distances

Answer: D

 [Watch Video Solution](#)

Chemistry

1. In the Millikan's oil drop method, which of the following force does not act on the oil drop?

A. Gravitational force

B. Viscous force

C. Magnetic force

D. Electrostatic force

Answer: C



Watch Video Solution

2. Match the followig



The correct answer is

A. (a) (b) (c) (d)
(iv) (v) (iii) (ii)

B. (a) (b) (c) (d)
(iii) (iv) (v) (ii)

C. (a) (b) (c) (d)
(iv) (ii) (iii) (i)

D. (a) (b) (c) (d)
(v) (iv) (ii) (iii)

Answer: A



[View Text Solution](#)

3. Which one of the following represents Boyle's temperature of a gas?

- A. The temperature at which an ideal gas obeys Boyle's law.
- B. The temperature at which an ideal gas obeys Boyle's is less than 1 for a real gas.
- C. The temperature at which a real gas obeys ideal gas law over an appreciable range of pressure.
- D. The temperature at which the compressibility factor deviates from 1 for an ideal gas.

Answer: C



[Watch Video Solution](#)

4. The order of stability of aromatic hydrocarbons given below is



A. $A < D < B > C$

B. $D < A < B < C$

C. $B > C > D > A$

D. $A > D > B > C$

Answer: D



[View Text Solution](#)

5. match the following.



The correct answer is

A. $\begin{matrix} A & B & C & D \\ IV & II & I & III \end{matrix}$

B. $\begin{matrix} A & B & C & D \\ II & V & I & III \end{matrix}$

- C. $A \quad B \quad C \quad D$
 $III \quad II \quad I \quad V$
- D. $A \quad B \quad C \quad D$
 $IV \quad V \quad I \quad III$

Answer: A



[View Text Solution](#)

6. Arrange the following compounds in the correct order of their acid strength.



A. $a > d > c > d$

B. $a > b > c > d$

C. $b > c > d > a$




D. $b > a > d > c$

Answer: D



[View Text Solution](#)

7. What are the suitable conditions for the following transformation ?



Answer: A



[View Text Solution](#)

8. The compound 'A' decolourises Br_2 / CCl_4 and releases N_2 gas with HNO_2 . The compound 'A' is



C. 

D. 

Answer: A



Watch Video Solution