



MATHS

BOOKS - TS EAMCET PREVIOUS YEAR PAPERS

ONLINE QUESTION PAPER

Mathematics

1. The set of all values of x and the set of all values of a for which the real valued function

$f(x) = \sqrt{\log_a(x - [x])}$ is defined are respectively

A. $R - Z$ and $(0, 1)$

B. Z and $R - \{0, 1\}$

C. Z^- and $(1, \infty)$

D. R and R

Answer: A



View Text Solution

2. A function $f, \mathbb{R} - \{0\} \rightarrow \mathbb{R}$ is defined as

$$f(x) = \begin{cases} x^2 + 3x - 7, & x > 0 \\ h(x), & x < 0 \end{cases}$$

If $f(x)$ is an odd function, then $h(x) =$

A. $x^2 + 3x + 7$

B. $x^2 + 3x - 7$

C. \mathbb{Z}^- and $(1, \infty)$

D. $-x^2 - 3x + 7$

Answer: C



Watch Video Solution

3. $x^n + y^n$ is divisible by

A. $x - y$ for all $n \in N$

B. $x + y$ for all $n \in N$

C. $x + y$ for all $n = 2m - 1, m \in N$

D. $x + y$ for all $n = 2m, m \in N$

Answer: C



Watch Video Solution

4. Consider the set A of all determinants of order 3 with entries 0 or 1 only . Let B be the subset of A containing of all determinants with value 1. Let C be the subset of the set of all determinants with value - 1. Then

A. $n(C) = 0$

B. $n(B) = n(C)$

C. $A = B \cup C$

D. $n(B) = 2n(A)$

Answer: B



Watch Video Solution

5.
$$\begin{vmatrix} 1 & bc + ad & b^2c^2 + a^2d^2 \\ 1 & ca + bd & c^2a^2 + b^2d^2 \\ 1 & ab + cd & a^2b^2 + c^2d^2 \end{vmatrix} =$$

A. $(a - b)(b - c)(c - d)(a - d)(a - c)(d - b)$

B. $(a - b)(a - c)(b - c)(b - d)(a - d)(c - d)$

C. $(a - b)(a - c)(a - d)(b - c)(b - d)(d - c)$

D. $(a - b)(b - c)(c - d)(b - d)$

Answer: B



View Text Solution

6. Let λ and α be real. The set of all values of λ for which the system of linear equations

$$\lambda x + (\sin \alpha)y + (\cos \alpha)z = 0$$

$$x + (\cos \alpha)y + (\sin \alpha)z = 0$$

$$-x + (\sin \alpha)y - (\cos \alpha)z = 0 \text{ has a non trivial solution is}$$

A. $\frac{n\pi}{2} + (-1)^n \frac{\pi}{4} + \frac{\pi}{8}$ (n is an integer)

B. $\frac{n\pi}{2} + (-1)^n \frac{\pi}{8}$ (n is an integer)

C. $\frac{n\pi}{2} + (-1)^n \frac{\pi}{8} - \frac{\pi}{8}$ (n is an integer)

D. $\frac{n\pi}{2} + (-1)^n \frac{\pi}{4} - \frac{\pi}{8}$ (n is an integer)

Answer: C



Watch Video Solution

7. If $\frac{1 - 10i \cos \theta}{1 - 10\sqrt{3}i \sin \theta}$ is purely real, then one of the values of θ is

A. $\frac{\pi}{6}$

B. $\frac{\pi}{4}$

C. $\frac{\pi}{3}$

D. $\frac{\pi}{2}$

Answer: A



Watch Video Solution

8. If z and w are complex numbers such that

$$\bar{z} - \overline{iw} = 0 \text{ and } \text{Arg}(zw) = \frac{3\pi}{4}, \text{ then } \text{Arg } z =$$

A. $\frac{\pi}{16}$

B. $\frac{\pi}{8}$

C. $\frac{\pi}{4}$

D. $\frac{3\pi}{4}$

Answer: B



Watch Video Solution

9. Find all the roots of the equation

$$x^{11} - x^7 + x^4 - 1 = 0$$

A. 2

B. 3

C. 7

D. 9

Answer: A



Watch Video Solution

10. If α is a root of $z^2 - z + 1 = 0$, then

$$\begin{aligned} & \left(\alpha^{2014} + \frac{1}{\alpha^{2014}} \right) + \left(\alpha^{2015} + \frac{1}{\alpha^{2015}} \right)^2 \\ & + \left(\alpha^{2016} + \frac{1}{\alpha^{2016}} \right)^3 + \left(\alpha^{2017} + \frac{1}{\alpha^{2017}} \right)^4 + \\ & \left(\alpha^{2018} + \frac{1}{\alpha^{2018}} \right)^5 = \end{aligned}$$

A. 8

B. 5

C. 3

D. -5

Answer: A



Watch Video Solution

11. Let $E_1 \equiv ax^2 + bx + c$, $E_2 \equiv bx^2 + cx + a$, $E_3 \equiv cx^2 + bx + a$ and $\frac{a^2}{bc} + \frac{b^2}{ca} + \frac{c^2}{ab} = 3$. If these quadratic expressions have a common zero, then the quadratic expression having zeros that are common to E_3 and different from the zeros of E_1 is

A. $x^2 - \frac{a(b+c)}{bc}x + bc$

B. $ax^2 + bx + c$

C. $x^2 - b(c+a)x + ac$

D. $x^2 - \frac{a(b+c)}{bc} + \frac{a^2}{bc}$

Answer: D



View Text Solution

12. If for any real x , $\frac{11x^2 + 12x + 6}{x^2 + 4x + 2} = y$ is such that $y < a$ or $y \geq b$, then a, b are

A. 3, 5

B. $-5, 3$

C. $-4, 5$

D. $-6, 4$

Answer: B



Watch Video Solution

13. Given that the roots of $x^3 + 3px^2 + 3qx + r = 0$ are in harmonic progression Then,

A. $2q^3 = r(3pq - r)$

B. $q^3 = r(3pq - r)$

C. $q^3 = -r(3pq - r)$

D. $q^3 = r(r + 3pq)$

Answer: A



Watch Video Solution

14. If α, β, γ are the roots of $x^3 + px^2 + qx + r = 0$, then $\alpha^3 + \beta^3 + \gamma^3 =$

A. $p^3 - 3pq + r$

B. $p^2 - 2pq + r$

C. $3pq - 3r - p^3$

D. $3pq + 3r + p^3$

Answer: C



[Watch Video Solution](#)

15. The number of proper divisors of the number obtained by dividing $13!$

With 100 is

A. 216

B. 430

C. 214

D. 790

Answer: B



[Watch Video Solution](#)

16. In an admission test, there are 15 multiple choice questions. Each question is followed by 4 alternatives to choose. Out of these there may be one or more than one correct answers. If a student attempts all the 15 question and marks the answers randomly the question paper is

A. $4 \times 15 C_4$

B. 15^{15}

C. 4^{15}

D. $4!. 15!$

Answer: B



Watch Video Solution

17. The absolute value of the numerically greatest term in the expansion of $(2x - 3y)^{12}$ when $x = 3, y = 2$ is

A. $^{12}C_5 6^{12}$

B. $^{12}C_6 6^{12}$

C. $^{12}C_4 6^{12}$

D. $^{12}C_9 6^{12}$

Answer: B

 [Watch Video Solution](#)

18. The sum to infinite terms of the series

$$\frac{3}{10} + \frac{3.7}{10.15} + \frac{3.7.9}{10.15.20} + \dots \text{ to } \infty \text{ is}$$

A. $\sqrt[4]{125} - 1$

B. $\frac{5\sqrt{5}}{3\sqrt{3}} - \frac{8}{5}$

C. $\sqrt[3]{4} - \frac{4}{3}$

D. $\sqrt{\frac{5}{3}} - \frac{6}{5}$

Answer: B

 [Watch Video Solution](#)

19.

$$\text{if } \frac{3x + 2}{(x + 1)(2x^2 + 3)} = \frac{A}{x + 1} + \frac{Bx + C}{2x^2 + 3}, \text{ then } A - B + C =$$

A. 1

B. 2

C. 3

D. 5

Answer: B



Watch Video Solution

20.

$\sin 10^\circ \sin 50^\circ \sin 60^\circ \sin 70^\circ = m$ and $\tan 20^\circ \tan 40^\circ \tan 60^\circ \tan 80^\circ = r$

then $\frac{n}{m} =$

A. $\frac{3\sqrt{3}}{16}$

B. $16\sqrt{3}$

C. $\frac{16}{\sqrt{3}}$

D. $8\sqrt{3}$

Answer: B

[Watch Video Solution](#)

21. Assertion (A) If $\alpha = 12^\circ, \beta = 15^\circ, \gamma = 18^\circ$, then

$\tan 2\alpha \tan 2\beta + \tan 2\beta \tan 2\gamma + \tan 2\gamma \tan 2\alpha = 1$ Reason (R) In

$$\triangle ABC, \tan \frac{A}{2} \tan \frac{B}{2} + \tan \frac{B}{2} \tan \frac{C}{2} + \tan \frac{C}{2} \tan \frac{A}{2} =$$

Which of the following is true?

- A. Both (A) and (R) are true and (R) is the correct explanation of (A)
- B. Both (A) and (R) are true, but (R) is not the correct explanation of (A)
- C. (A) is true, but (R) is false
- D. (A) is false, but (R) is true

Answer: A

[View Text Solution](#)

22. $\cot \theta - \tan \theta - 2 \tan 2\theta - 4 \tan 4\theta =$

A. $4 \cot 8\theta - \tan 6\theta$

B. $\cot 8\theta + \tan 3\theta$

C. $\cot 8\theta + \cot 6\theta$

D. $8 \cot 8\theta$

Answer: D



Watch Video Solution

23. If the general solution of

$\sin x + 3 \sin 3x + \sin 5x = 0$ is $x = y$ then the set of all values of $\cos y$ is

A. $\left\{ -1, -\frac{\sqrt{3}}{2}, \frac{\sqrt{3}}{2}, 1 \right\}$

B. $\left\{ -1, \frac{1}{2}, 1 \right\}$

C. $\left\{ -\frac{\sqrt{3}}{2}, 0, 1, \frac{\sqrt{3}}{2} \right\}$

D. $\left\{ -1, -\frac{1}{2}, \frac{1}{2}, 1 \right\}$

Answer: D



Watch Video Solution

24. If $\cos^{-1} 2x + \cos^{-1} 3x = \frac{\pi}{3}$, then $x =$

A. $\frac{\sqrt{3}}{2\sqrt{7}}$

B. $\frac{\sqrt{3}}{\sqrt{7}}$

C. $\frac{\sqrt{2}}{\sqrt{5}}$

D. $\frac{\sqrt{3}}{2\sqrt{5}}$

Answer: A



Watch Video Solution

25. If $\cosh \beta = \sec \alpha \cos \theta$, $\sinh \beta = \operatorname{cosec} \alpha \sin \theta$, then $\sinh^2 \beta =$

A. $\sin \alpha \cos \alpha$

B. $\cos^2 \alpha$

C. $\cos^2 \alpha$

D. $\sin \alpha + \cos \alpha$

Answer: C



Watch Video Solution

26. If $\sin A$, $\sin B$ are roots of $c^2x^2 - c(a+b)x + ab = 0$, then the Δ is

A. the triangle is acute angled

B. the triangle is obtuse angled

C. $\sin C = \frac{\sqrt{3}}{2}$

D. $\sin A + \cos A = \frac{a+b}{c}$

Answer: D



Watch Video Solution

27. Let ABC be an isosceles triangle with BC as its base. Then, $rr_1 =$

A. a^2

B. $\frac{a^2}{2}$

C. $R^2 \sin^2 A$

D. $R^2 \sin^2 2B$

Answer: C



Watch Video Solution

28. In a $\triangle ABC$, $a^4 + b^4 + c^4 = 2b^2c^2 + 2a^2b^2$, then B =

A. $\frac{\pi}{4}$ or $\frac{3\pi}{4}$

B. $\frac{\pi}{2}$

C. $\frac{\pi}{3}$ or $\frac{2\pi}{3}$

D. $\frac{\pi}{6}$ or $\frac{5\pi}{6}$

Answer: A



Watch Video Solution

29. If $a = 2\hat{i} + 3\hat{j} + \hat{k}$, $b = \hat{i} - 3\hat{j} - 5\hat{k}$ and $c = 3\hat{i} - 4\hat{k}$, then match the items of List - I with those of List - II.

List-I	List-II
A Unit vector in the direction opposite to that $\mathbf{a} - \mathbf{b}$ is	(i) $5\hat{i} + 3\hat{j} - 3\hat{k}$
B If $\mathbf{AB} = \mathbf{a}$, $\mathbf{BC} = \mathbf{b}$, then $\mathbf{CA} =$	(ii) $2\hat{i} - \frac{8}{3}\hat{k}$
C If \mathbf{a} , \mathbf{b} , \mathbf{c} are the position vectors of the vertices a triangle then, its centroid is	(iii) $-3\hat{i} + 4\hat{k}$
D In D part it is $\mathbf{b} + \mathbf{d}$ instead of $\mathbf{b} - \mathbf{d}$ If \mathbf{d} is a vector of magnitude $2\sqrt{14}$ and parallel to the vector \mathbf{a} , then $\mathbf{b} + \mathbf{d} =$	(iv) $-\frac{\hat{i}}{\sqrt{73}} - \frac{6\hat{j}}{\sqrt{73}} - \frac{6\hat{k}}{\sqrt{73}}$
	(v) $\frac{3}{\sqrt{43}}\hat{i} + \frac{5}{\sqrt{43}}\hat{j} - \frac{3}{\sqrt{43}}\hat{k}$

A. $\begin{matrix} A & B & C & D \\ iv & iii & ii & i \end{matrix}$

- B. $\begin{matrix} A & B & C & D \\ iv & iii & ii & v \end{matrix}$
- C. $\begin{matrix} A & B & C & D \\ iv & iii & i & ii \end{matrix}$
- D. $\begin{matrix} A & B & C & D \\ i & ii & iii & v \end{matrix}$

Answer: A



Watch Video Solution

30. If $2\hat{i} - \hat{j} + 3\hat{k}$, $-12\hat{i} - \hat{j} - 3\hat{k}$, $-\hat{i} + 2\hat{j} - 4\hat{k}$ and $\lambda\hat{i} + 2\hat{j} - \hat{k}$ are the position vectors of four coplanar points, then $\lambda =$

- A. -2
- B. 6
- C. 3
- D. -6

Answer: B



Watch Video Solution

31. A point lying on the plane that passes through the point

$\hat{i} - \hat{j} + \hat{k}$, $\hat{i} - 2\hat{j} + 3\hat{k}$ and $\hat{i} + 2\hat{j} - 3\hat{k}$ is

A. $-\hat{i} + 2\hat{j} - 3\hat{k}$

B. $-\hat{i} + \hat{j} - \hat{k}$

C. $\hat{i} + \hat{j} - \hat{k}$

D. $4\hat{i} + 2\hat{j} + 3\hat{k}$

Answer: C



Watch Video Solution

32. A non-zero vector a is parallel to the line of intersection of the plane

determined by the vectors \hat{i} , $\hat{i} + \hat{j}$ and the plane determined by vectors

$\hat{i} - \hat{j}$, $\hat{j} + \hat{k}$. The angle between a and

$(\hat{i} - 2\hat{j} + 2\hat{k})$ is

A. $\frac{\pi}{6}$

B. $\frac{\pi}{4}$

C. $\frac{\pi}{3}$

D. $\frac{2\pi}{5}$

Answer: D



Watch Video Solution

33. The equation of the plane passing through the point with position vectors $A(2\hat{i} + 6\hat{j} - 6\hat{k})$, $B(-3\hat{i} + 10\hat{j} - 9\hat{k})$ and $C(-5\hat{i} - 6\hat{k})$ is

A. $r \cdot (2\hat{i} - \hat{j} - 2\hat{k}) = 2$

B. $r \cdot (\hat{i} - 2\hat{j} - \hat{k}) = 1$

C. $r \cdot (2\hat{i} + \hat{j} - 2\hat{k}) = 3$

D. $r \cdot (\hat{i} + 2\hat{j} - 2\hat{k}) = 3$

Answer: A



Watch Video Solution

34. If a , b and c are three vectors with magnitudes 1, 1 and 2 respectively and $a \times (a \times c) + b = 0$, then the angle between a and c is

A. $\frac{2\pi}{5}$

B. $\frac{\pi}{3}$

C. $\frac{\pi}{4}$

D. $\frac{\pi}{6}$

Answer: D



Watch Video Solution

35. The coefficient of variation of the first n natural numbers is

A. $\frac{100}{\sqrt{3}}(n-1)$

B. $\frac{100}{\sqrt{3}}\sqrt{\frac{n+1}{n-1}}$

C. $\frac{\sqrt{3}}{100}\sqrt{\frac{n+1}{n-1}}$

D. $\frac{100}{\sqrt{3}}\sqrt{\frac{n-1}{n+1}}$

Answer: D



View Text Solution

36. Two distributions A and B have the same mean. If their coefficients of variation are 6 and 2 respectively and σ_A, σ_B are their standard deviations, then

A. $\sigma_A = 3\sigma_B$

B. $3\sigma_A = \sigma_B$

C. $\sigma_A = 2\sigma_B$

D. $2\sigma_A = \sigma_B$

Answer: A



Watch Video Solution

37. From a certain population, the probability of choosing a colour blind man is $\frac{1}{20}$ and that of a colour blind woman is $\frac{1}{10}$. If a randomly chosen person is found to be colour blind, then the probability that the person is a man is

A. $\frac{2}{9}$

B. $\frac{2}{3}$

C. $\frac{1}{3}$

D. $\frac{1}{9}$

Answer: C



Watch Video Solution

38. From a lot containing n good and m bad articles, if 2 articles are picked at random in succession without replacement, then the probability that the second article picked is bad is

A. $\frac{m}{m+n}$

B. $\frac{m-1}{m+n}$

C. $\frac{(n-1)(m-1)}{(m+n)^2}$

D. $\frac{mn}{(m+n)^2}$

Answer: A



Watch Video Solution

39. In a classroom 5% of the boys and 2% of the girls taller than 1.6 m. The class consists of 60% girl student. The probability that a randomly selected student is taller than 1.6 m is

A. $\frac{121}{125}$

B. $\frac{5}{8}$

C. $\frac{3}{8}$

D. $\frac{4}{125}$

Answer: D



Watch Video Solution

40. The distribution of a random variable X is given below

$X = x$	1	2	3	4
$P(X = x)$	$2c$	$4c$	$6c$	$8c$

Then, the standard deviation of X is

A. 4

B. $\frac{3}{2}$

C. 2

D. 1

Answer: D



Watch Video Solution

41. The probability of securing a succes in an trial is three times that of a failure. The probability of getting atleast 4 successes in 5 trial is

A. $\frac{649}{1024}$

B. $\frac{81}{128}$

C. $\frac{27}{64}$

D. $\frac{243}{1024}$

Answer: B



Watch Video Solution

42. If the line jjoining the point $A(b \cos \alpha, b \sin \alpha)$ and $B(a \cos \beta, a \sin \beta)$ is extended to the point $N(x, y)$ such that $AN : NB = b : a$, then

A. Option1 $x \cos \frac{\alpha - \beta}{2} + y \sin \frac{\alpha + \beta}{2} = 0$

B. Option2 $x \cos \frac{\alpha - \beta}{2} + y \sin \frac{\alpha - \beta}{2} = 0$

C. Option3 $x \cos \frac{\alpha - \beta}{2} + y \sin \frac{\alpha + \beta}{2} = 0$

D. Option4 $x \cos \frac{\alpha + \beta}{2} + y \sin \frac{\alpha + \beta}{2} = 0$

Answer: C



Watch Video Solution

43. A straight line $x - 2y - 4 = 0$ is shifted parallel to it by 3 units away from the origin and then rotated by an angle of 30° in the anti-clockwise direction. If the slope of the new line formed is m , then the integral part of ' m ' is

A. -1

B. 0

C. 1

D. 2

Answer: C



View Text Solution

44. A straight line $x - 2y - 4 = 0$ is shifted parallel to it by 3 units away from the origin and then rotated by an angle of 30° in the anti-clockwise direction. If the slope of the new line formed is m , then the integral part of ' m ' is

A. $-\frac{13\pi}{12}$

B. $\frac{29\pi}{12}$

C. $-\frac{11\pi}{12}$

D. $\frac{35\pi}{12}$

Answer: D



View Text Solution

45. The straight lines $x + 3y - 4 = 0$, $x + y - 4 = 0$ and $3x + y - 4 = 0$

- A. from an isosceles triangle
- B. are concurrent
- C. from an equilateral triangle
- D. from a right angled isosceles triangle

Answer: A



Watch Video Solution

46. The combined equation of the straight lines passing through $(1, 1)$ and making an angle of 45° with the straight line $x + y - 1 = 0$ is

- A. $2x^2 + 3xy - 2y^2 - 7x + y + 1 = 0$
- B. $xy - x - y + 1 = 0$
- C. $xy + 2y^2 - x - 5y - 3 = 0$

D. $2x^2 - xy - 3x + y + 1 = 0$

Answer: B



Watch Video Solution

47. The centroid of the triangle formed by the pair of straight lines $12x^2 - 20xy + 7y^2 = 0$ and the line $2x - 3y + 4 = 0$ is (α, β) . Then, $\alpha + 2\beta =$

A. $-\frac{4}{3}$

B. 2

C. 8

D. $-\frac{8}{3}$

Answer: C



Watch Video Solution

48. The equation of the bisectors of the angles between the lines joining the origin to the points of intersection of the curve

$$x^2 + xy + y^2 + x + 3y + 1 = 0 \text{ and the line}$$

$$x + y + 2 = 0 \text{ is}$$

A. $2x^2 - 4xy + y^2 = 0$

B. $x^2 - 4xy + y^2 = 0$

C. $2x^2 + 4xy + y^2 = 0$

D. $x^2 + 4xy - y^2 = 0$

Answer: D



Watch Video Solution

49. Consider the following statements.

I. The intercept made by the circle

$$x^2 + y^2 - 2x - 4y + 1 = 0 \text{ on Y-axis is } 2\sqrt{3}$$

II. The intercept made by the circle

$$x^2 + y^2 - 4x - 2y + 6 = 0 \text{ on X-axis is } 2\sqrt{2}$$

III. The straight line $y = 2x + 1$ cuts the circle $x^2 + y^2 = 9$ at two distinct points

Then which one of the following option is correct?

- A.

<i>I</i>	<i>II</i>	<i>III</i>
True	True	True
- B.

<i>I</i>	<i>II</i>	<i>III</i>
True	True	False
- C.

<i>I</i>	<i>II</i>	<i>III</i>
True	False	True
- D.

<i>I</i>	<i>II</i>	<i>III</i>
False	False	True

Answer: C



Watch Video Solution

50. If the circle

$x^2 + y^2 + 2kx - 4y + 1 = 0$ and $x^2 + y^2 - 8x - 12y + 43 = 0$ touch each other

A. 2

B. 1

C. -1

D. -2

Answer: C



Watch Video Solution

51. For all real values of k , the point which lies on the polar of $(k, k + 1)$ with respect to the circle

$$x^2 + y^2 + 4x - 8y - 5 = 0 \text{ is}$$

A. $(3, -1)$

B. $(3, 1)$

C. $(2, -2)$

D. $(2, 3)$

Answer: A



Watch Video Solution

52. The number of common tangents to the circles

$$x^2 + y^2 + 4x = 0 \text{ and } x^2 + y^2 - 2x = 0 \text{ is}$$

A. 4

B. 3

C. 2

D. 1

Answer: B



Watch Video Solution

53. If $\frac{2}{\sqrt{5}}$ is the length of the common chord of the circles

$$x^2 + y^2 + 2x + 2y + 1 = 0 \quad \text{and}$$

$$x^2 + y^2 + \alpha x + 3y + 2 = 0, \alpha \neq 0, \text{ then } \alpha =$$

A. 4

B. 3

C. 2

D. 1

Answer: A



Watch Video Solution

54. Match the item of list-I with those of List-II

List-I	List-II
A Equation of the tangent drawn at $(2, \sqrt{8})$ on the curve $y^2 = 4x$ is	(i) -36
B Equation of the normal to the curve $y^2 = 16x$, that makes and angle of 45° with its axis is	(ii) 4
C The chord joining the points (x_1, y_1) and (x_2, y_2) on the curve $y^2 = 12x$ is a focal chord if $y_1 y_2 =$	(iii) 8
D A value of k for which $x - 3 = 0$ is the directrix of the curve $y^2 - kx + 16 = 0$ is	(iv) $x - \sqrt{2}y + 2 = 0$
	(v) $x + y - 12 = 0$
	(vi) $x - y - 12 = 0$

Then, which of the following is correct ?

- A. $\begin{matrix} A & B & C & D \\ v & iv & iii & ii \end{matrix}$
- B. $\begin{matrix} A & B & C & D \\ iv & v & ii & i \end{matrix}$
- C. $\begin{matrix} A & B & C & D \\ iv & vi & i & ii \end{matrix}$
- D. $\begin{matrix} A & B & C & D \\ iv & vi & ii & iii \end{matrix}$

Answer: C



View Text Solution

55. An equilateral is inscribed in the parabola $y^2 = 8x$ with one of its vertices is the vertex of the parabola. Then the length of the side of that triangle is

- A. $(8a, 0)$
- B. $(16a, 0)$
- C. $(32a, 0)$
- D. $(48a, 0)$

Answer: C



Watch Video Solution

56. If the straight line $8x + 3\sqrt{2}y = 36$ touches the

ellipse $\frac{x^2}{9} + \frac{y^2}{4} = 2$ at (a, b) , then $a + \sqrt{2b} =$

A. $\frac{36}{5\sqrt{2}}$

B. $\frac{8}{3}$

C. $\frac{12 + 2\sqrt{2}}{3}$

D. $\frac{16}{3}$

Answer: D



Watch Video Solution

57. The eccentricity of an ellipse, with its centre at the origin, is $1/2$. If one of the directrices is $x = 4$, then the equation of the ellipse is

A. $3x^2 + 4y^2 = 12$

B. $3x^2 + 4y^2 = 49$

C. $3x^2 + 4y^2 = 1$

D. $4x^2 + 3y^2 = 12$

Answer: A



Watch Video Solution

58. If the product of the slopes of the tangents drawn from an external point P to the hyperbola $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$ is a constant k^2 , then the locus of P is

A. $y^2 + b^2 = k^2(x^2 - a^2)$

B. $y^2 - b^2 = k^2(x^2 - a^2)$

C. $x^2 + b^2 = k^2(y^2 - a^2)$

D. $x^2 - b^2 = k^2(y^2 - a^2)$

Answer: A



View Text Solution

59. Let A, B, C be three points on \overline{OX} , \overline{OY} , \overline{OZ} respectively at the distances 3, 6, 9 from origin.

Let Q be the point (2, 5, 8) and P be the point equidistant from O,A,B,C. Then, the coordinates of the point R which divides PQ in the ratio : 2 is

A. $\left(\frac{17}{10}, \frac{29}{5}, \frac{43}{10}\right)$

B. $\left(\frac{7}{5}, \frac{16}{5}, 5\right)$

C. $\left(\frac{9}{5}, \frac{21}{5}, \frac{33}{5}\right)$

D. $\left(\frac{8}{5}, \frac{19}{5}, 6\right)$

Answer: C



View Text Solution

60. If the direction cosines of two lines are such

that $l + m + n = 0$, $l^2 + m^2 - n^2 = 0$, then the angle between them is

A. $\frac{\pi}{6}$

B. $\frac{\pi}{4}$

C. $\frac{\pi}{3}$

D. $\frac{\pi}{2}$

Answer: C



Watch Video Solution

61. If the line joining the points A (1, 0, 0) and B(0, 0, 1) is a normal to the plane (π) which passes through the point A, then the angle between the planes (π) and $x + y + z = 6$ is

A. $\frac{\pi}{6}$

B. $\frac{\pi}{4}$

C. $\frac{\pi}{3}$

D. $\frac{\pi}{2}$

Answer: A



Watch Video Solution

62. The number of points at which the function

$$f(x) = \frac{\sqrt{11 + |x|} - 6\sqrt{2 + |x|}}{6 - 2\sqrt{2 + |x|}}$$

is discontinuous in

$(-\infty, \infty)$ is

A. 1

B. 0

C. 2

D. 3

Answer: C



View Text Solution

63. If $f(x) = \begin{cases} \frac{\sqrt{1+px} - \sqrt{1-px}}{x}, & -1 \leq x < 0 \\ \frac{2x+1}{x-2}, & 0 \leq x \leq 1 \end{cases}$

continuous on $[-1,1]$, then $p =$

A. $-\frac{1}{2}$

B. $-\frac{1}{4}$

C. $\frac{1}{2}$

D. 2

Answer: A



Watch Video Solution

64. If $y = \tan^{-1}(\sin \sqrt{x}) + \operatorname{cosec}^{-1}(e^{2x+1})$, then $\frac{dy}{dx} =$

A. $\frac{1}{\sqrt{x}(1 + \sin^2 \sqrt{x})} + \frac{1}{\sqrt{e^{4x+2} + 1}}$

B. $\frac{\cos \sqrt{x}}{2\sqrt{x}(1 + \sin^2 \sqrt{x})} - \frac{2}{\sqrt{e^{4x+2} - 1}}$

$$\text{C. } \frac{\cos \sqrt{x}}{(1 + \sin^2 \sqrt{x})} + \frac{2}{\sqrt{e^{4X+2} + 1}}$$

$$\text{D. } \frac{1}{2\sqrt{x}} \frac{\cos \sqrt{x}}{1 + \sin^2 \sqrt{x}} - \frac{1}{\sqrt{e^{2X+1} - 1}}$$

Answer: B



Watch Video Solution

65. If $f : R \rightarrow R$ is such that $f(3) = 16, f'(3) = 4$, then

$$\lim_{x \rightarrow 3} \frac{xf(3) - 3f(x)}{x - 3} =$$

A. 4

B. 6

C. 8

D. 12

Answer: A



Watch Video Solution

66. If $y = e^x(\log x)$, then $xy_2 + (x - 1)y_1 =$

A. $(2x - 1)y_1$

B. $(x - 1)y_1$

C. $(4 - 2x)y_1$

D. $(3x - 1)y_1$

Answer: A



Watch Video Solution

67. Let $f(x) = e^x \cos x + 1$. Which of the following statements is always true?

A. Between any two consecutive roots of $f(x) = 0$

there is always a root of $e^x \sin x + 1 = 0$

B. Between any two consecutive roots of $f(x) = 0$

there is always a root of $e^x \sin x - 1 = 0$

C. Between any two consecutive roots of $f(x) = 0$

there is always a root of $e^x \cos x = 0$

D. Between any two consecutive roots of $f(x) = 0$

there is always a roots of $e^x \sin x = 0$

Answer: D



View Text Solution

68. The radius of a circular plate is increasing at the rate of 0.01 cm/sec when the radius is 12 cm. Then the rate at which the area increases is

A. 60π

B. 24π

C. 1.2π

D. 0.24π

Answer: B

[Watch Video Solution](#)

69. Let $f(x) = x^2 e^{-2x}$, $x > 0$. The maximum value of $f(x)$ is

A. 0

B. $\frac{1}{e^2}$

C. $\frac{1}{4e^2}$

D. $\frac{1}{2e}$

Answer: B

[Watch Video Solution](#)

70. Let $f(x)$ be differentiable on $[1, 6]$ and $f(1) = -2$.

If $f(x)$ has only one root in $(1, 6)$ then there exists $c \in (1, 6)$ such that

A. $f'(c) = \frac{1}{10}$

B. $f'(c) < \frac{2}{5}$

C. $f'(c) < \frac{1}{5}$

D. $f'(c) > \frac{2}{5}$

Answer: D



View Text Solution

71. If $\int \phi(x) dx = \Psi(x)$, then $\int (\phi \circ h)(x) h(x) h'(x) dx =$

A. $(\phi \circ h)(x) \phi'(x) - \int (\phi \circ h)(x) h'(x) dx + c$

B. $(\Psi \circ h)(x) h(x) - \int (\Psi \circ h)(x) h'(x) dx + c$

C. $(\Psi \circ h)(x) \phi(x) - \int (\Psi \circ h)(x) \phi'(x) dx + c$

D. $(\Psi \circ \Phi)(x) h(x) - \int (\Psi \circ \phi)(x) h'(x) dx + c$

Answer: B



Watch Video Solution

72. If $f\left(\frac{t+1}{2t+1}\right) = t+1$, then $\int f(x)dx =$

A. $\frac{x^2}{2} + c$

B. $\log(2x-1) + \frac{1}{2}\log(x+1) + c$

C. $\frac{1}{2}\log(2x-1) + c$

D. $\frac{x}{2} + \frac{1}{4}\log(2x-1) + c$

Answer: D



Watch Video Solution

73. $\int \frac{x^8 - 9x^2 + 18}{x^4 - 3x^2 + 13} dx =$

A. $\frac{X^4}{4} + X^3 + 6X^2 + c$

B. $\frac{X^5}{5} + \frac{X^4}{4} + 6X + c$

C. $\frac{X^5}{5} + X^3 + 6X + c$

D. $\frac{X^5}{5} + X^3 + 6X + c$

Answer: C



View Text Solution

74. $\int (\cot x \cot(x + \alpha) + 1) dx =$

A. $\cot \alpha \log \left(\left| \frac{\sin x}{\sin(x + \alpha)} \right| \right) + c$

B. $\log |\sin x \sin(x + \alpha)| + x + c$

C. $\log |\sin x \cos(x + \alpha)| + x + c$

D. $\tan \alpha \log \left(\left| \frac{\cos x}{\sin(x + \alpha)} \right| \right) + c$

Answer: A



View Text Solution

75. If $\lim_{n \rightarrow \infty} \sum_{r=1}^n \frac{4r^3}{r^4 + n^4} = p$, then $e^4 =$

A. 4

B. 3

C. 2

D. 1

Answer: C



View Text Solution

76. $\int_0^{\pi/4} \left[\sqrt{\frac{1 - \sin 2x}{1 + \sin 2x}} \right] dx =$

A. 1

B. $2\log 2$

C. $2\log \sqrt{2}$

D. 2

Answer: D



Watch Video Solution

77. The area bounded by the curve

$y = x^3 - 3x^2 + 2x$ and the X-axis is (in square units)

A. $\frac{1}{2}$

B. $\frac{5}{2}$

C. 1

D. 4

Answer: A



Watch Video Solution

78. The general solution of the differential equation $\cos(x + y)dy = dx$ is

A. $y = \sec(x + y) + c$

B. $y - \tan \frac{y}{2} = x + c$

C. $y = \tan \left(\frac{x + y}{2} \right) + c$

D. $y = \frac{1}{2} \tan(x + y) + c$

Answer: C



Watch Video Solution

79. The general solution of the differential equation $(x^3 - 3xy^2)dx = (y^3 - 3x^2y)dy$ is where c is an arbitrary constant

A. $c^2(X^2 + y^2) = (y^2 - X^2)$

B. $c^2(X^2 + y^2) = (y^2 - X^2)$

C. $c^2(X^2 + y^2)^2 = (y^2 - X^2)$

D. $c^2(X^2 - y^2)^2 = (y^2 - X^2)$

Answer: C



Watch Video Solution

80. The differential equation corresponding to all the circle lting in the first quadrant and touching the coordinate axes is

A. $(x - y)^2 \left[1 + \left(\frac{dy}{dx} \right)^2 \right] = \left(x + y \frac{dy}{dx} \right)^2$

B. $(x - y)^2 \left[1 + \frac{dy}{dx} \right]^2 = \left(x + y \frac{dy}{dx} \right)^2$

C. $(x - y)^2 \left[1 + \left(\frac{dy}{dx} \right)^2 \right] = x + y \left(\frac{dy}{dx} \right)^2$

D. $(x - y)^2 \left[1 + \frac{dy}{dx} \right] = \left(x + y \frac{dy}{dx} \right)^{\frac{1}{2}}$

Answer: A



Watch Video Solution

81. If $f: [1, \infty) \rightarrow [5, \infty)$ is given by $f(x) = 3x + \frac{2}{x}$, then $f^{-1}(x) =$

A. $\frac{1}{6} \left[x + \sqrt{x^2 - 24} \right]$

B. $\frac{x}{3x^2 + 2}$

C. $\frac{1}{6} \left[x - \sqrt{x^2 - 24} \right]$

D. $\frac{1}{2} \left[1 + \sqrt{x^2 - 4} \right]$

Answer: A



Watch Video Solution

82. The domation of $f(x) = \log \left[(25)^{3-x^2} - (0.4)^{x+9} \right]$ is

A. $(-4, 3)$

B. $(-3, 4)$

C. $(3, 4)$

D. $(0, \infty)$

Answer: B



View Text Solution

83. Let $n \in \mathbb{N}$ which one of the following is true?

A. $47^n + 16n - 1$ is divisible by 4

B. $2(4^{2n+1}) - 3^{3n+1}$ is divisible by 9

C. $4^n - 3n - 1$ is divisible by 11

D. $3(5^{2n+1}) + 2^{3n+1}$ is divisible by 17

Answer: D



View Text Solution

84. If $A = \begin{bmatrix} 1 & 2 & 2 \\ 2 & 1 & 2 \\ 2 & 2 & 1 \end{bmatrix}$, then $A^{-1} =$

A. $4I - A$

B. $A - 4I$

C. $\frac{1}{5}(A - 4I)$

D. $\frac{1}{5}(4I - A)$

Answer: C

[View Text Solution](#)

85. If x, y are any two non-zero real numbers, $a_{ij} = \xi + yj$, $A = (a_{ij})_{m \times n}$

and P, Q are two $n \times n$ matrices such that $A = xP + yQ$, then

- A. P is singular and Q is non-singular
- B. $P + Q$ is symmetric and $P - Q$ is skew symmetric
- C. Both $P + Q$ and $P - Q$ are singular
- D. Both $P + Q$ and $P - Q$ are non-singular

Answer: B

[View Text Solution](#)

86. If the system $\begin{bmatrix} 2 & 8 \\ 3 & 7 \end{bmatrix} \begin{bmatrix} a \\ b \end{bmatrix} = k \begin{bmatrix} a \\ b \end{bmatrix}$ has non-trivial, solution then the positive value of k and a solution of the system for that value of k are

- A. 9, $\begin{bmatrix} 3 \\ -8 \end{bmatrix}$

B. $10, \begin{bmatrix} -8 \\ 3 \end{bmatrix}$

C. $6, \begin{bmatrix} -1 \\ -1 \end{bmatrix}$

D. $10, \begin{bmatrix} 1 \\ 1 \end{bmatrix}$

Answer: D



View Text Solution

87. The modulus-amplitude form of $\frac{(1-i)^3(2-i)}{(2+i)(1+i)}$ is

A. $2cis\left(\pi \tan^{-1} \frac{4}{3}\right)$

B. $2cis\left(-\tan^{-1} \frac{4}{3}\right)$

C. $2cis\left(-\pi + \tan^{-1} \frac{4}{3}\right)$

D. $2cis\left(\tan^{-1} \frac{4}{3}\right)$

Answer: A



View Text Solution

88. If $z_1 = 2 - 3i$ and $z_2 = -1 + i$, then the locus of a point P represented by $z = x + iy$ in the argand plane satisfying the equation $\arg \left(\frac{z - z_1}{z - z_2} \right) = \frac{\pi}{2}$ is

A. $x^2 + y^2 - x + 2y - 5 = 0$

B. $x^2 + y^2 - x + 2y - 5 = 0$ and $4x + 3y + 1 < 0$

C. $4x + 3y + 1 = 0$ and $x^2 + y^2 - x + 2y - 5 > 0$

D. $x^2 + y^2 - x + 2y - 5 = 0$ and $4x + 3y + 1 > 0$

Answer: D



View Text Solution

89. $\left(\frac{1 + \cos \frac{\pi}{8} - i \sin \frac{\pi}{8}}{1 + \cos \frac{\pi}{8} + i \sin \frac{\pi}{8}} \right)^{12} =$

A. -1

B. i

C. $-i$

D. 2

Answer: B



Watch Video Solution

90. If the complex number a is such that $|a| = 1$ and $\arg(a) = \theta$, then the roots of the equation $\left(\frac{1 + iz}{1 - iz}\right)^4 = a$ are $z =$

A. $\tan\left(\frac{2k\pi + \theta}{4}\right), k = 0, 1, 2, 3$

B. $\tan\left(\frac{k\pi + \theta}{8}\right), k = 0, 1, 2, 3$

C. $\tan\left(\frac{3k\pi + \theta}{4}\right), k = 0, 1, 2, 3$

D. $\tan\left(\frac{2k\pi + \theta}{8}\right), k = 0, 1, 2, 3$

Answer: D



View Text Solution

91. If $x^2 + 2px - 2p + 8 > 0$ for all real values of x , then the set of all possible values of p is

A. $(2, 4)$

B. $(-\infty, -4)$

C. $(2, \infty)$

D. $(-4, 2)$

Answer: D



Watch Video Solution

92. If the roots of the equation $(p - 3)x^2 + 2(p - 3)x + 2p - 5 = 0$ are real number distinct for $\alpha < p < \beta$ and $(\beta - \alpha)$ is maximum, then the extreme value of the quadratic expression $-(\alpha + \beta)x^2 + \alpha\beta x + (\alpha - \beta)$ is

A. Option A $-\frac{4}{5}$

B. OptionB 5

C. OptionC -1

D. OptionD $\frac{4}{5}$

Answer: D



Watch Video Solution

93. If the equation $x^3 - 7x^2 + 14x - 8 = 0$ is transformed to $y^3 + py - \frac{20}{27} = 0$ when its roots are diminished by k , then $p =$

A. Option1 $\frac{8}{3}$

B. Option2 $\frac{7}{3}$

C. Option3 $\frac{-7}{3}$

D. Option4 $\frac{-8}{3}$

Answer: C



Watch Video Solution

94. If one root of the equation $x^3 - 9x^2 + 26x - 24 = 0$ is twice the other. Then, the sum of the cubes of those two roots is

A. 72

B. 253

C. 9

D. $\frac{9}{64}$

Answer: A



Watch Video Solution

95. There are 10 points in a plane of which no three points are collinear except 4. Then, the number of distinct triangles that can be formed by joining these points such that atleast one of the vertices of every triangle formed is formed the given 4 collinear points is

A. 116

B. 96

C. 120

D. 100

Answer: B



View Text Solution

96. The number of ways of arranging 8 boys and 8 girls in a row so that boys and girls sit alternately is

A. 9!

B. $(9!)(8!)$

C. $(8!)^2$

D. $2!(8!)^2$

Answer: D

97. If the coefficients of $(2\alpha + 4)th$ and $(\alpha - 2)th$ terms in the expansion $(1 + x)^{2019}$ are equal, then $\alpha =$

A. 673

B. 674

C. 675

D. 676

Answer: A

98. If n is a positive integer, then the coefficient of x^6 in the expansion of $(1 - 2x + 3x^2 - 4x^3 + \dots)^{-n}$ is

A. ${}^{(2n)}C_4$

B. ${}^nC_{12}$

C. ${}^{(2n)}C_6$

D. nC_6

Answer: C



View Text Solution

99. $\frac{d}{dx} \left(\frac{x+5}{(x+1)^2(x+2)} \right) =$

A. $\frac{8}{(x+2)^2} - \frac{3}{(x+1)^2} + \frac{3}{(x+1)^3}$

B. $\frac{3}{(x+1)^2} - \frac{3}{(x+2)^2} - \frac{8}{(x+1)^3}$

C. $\frac{3}{(x+2)^2} - \frac{3}{(x+1)^3} - \frac{8}{(x+1)^2}$

D. $\frac{8}{(x+2)^2} - \frac{3}{(x+1)^3} + \frac{3}{(x+1)^2}$

Answer: B



View Text Solution

100. If the period of the function $f(x) = \sin 5x \cos 3x$ is α , then $\cos \alpha =$

A. 1

B. $\frac{1}{\sqrt{2}}$

C. $-\frac{1}{2}$

D. -1

Answer: D



View Text Solution

101. If $\cos \frac{\pi}{15} \cos \frac{2\pi}{15} \cos \frac{4\pi}{15}$
 $\cos \frac{5\pi}{15} \cos \frac{7\pi}{15} \cos \frac{30\pi}{15} = x$, then $\frac{1}{8x} =$

A. 4

B. $\frac{1}{4}$

C. 8

D. $\frac{4}{3}$

Answer: A



View Text Solution

102. If $A + B + C = 2S$, then $\sin(S - A) + \sin(S - B) - \sin C =$

A. $-4\sin\frac{S-A}{2}\sin\frac{S-B}{2}\sin\frac{C}{2}$

B. $4\sin\frac{S-A}{2}\sin\frac{S-B}{2}\sin\frac{C}{2}$

C. $-4\sin\frac{S-A}{2}\sin\frac{S-B}{2}\cos\frac{C}{2}$

D. $4\sin\frac{S-A}{2}\sin\frac{S-B}{2}\cos\frac{C}{2}$

Answer: B



View Text Solution

103. When a is irrational, the number of solutions satisfying the equation

$$1 + \sin^2 ax = \cos x \text{ is}$$

A. 1

B. 0

C. 2

D. Infinite

Answer: A



Watch Video Solution

104. $\tan^{-1}\left(\frac{1}{2\sqrt{2}}\right) + \sin^{-1}\left(\frac{1}{\sqrt{3}}\right) = \cos^{-1}x$, then $x =$

A. $\frac{1}{\sqrt{3}}$

B. $\frac{1}{\sqrt{2}}$

C. $\frac{2}{\sqrt{3}}$

D. $\frac{1}{2\sqrt{2}}$

Answer: A



View Text Solution

105. $\tanh^{-1}\left(\frac{1}{3}\right) + \coth^{-1}(3) =$

A. $\log_e\left(\frac{2}{\sqrt{3}}\right)$

B. $\log_e 2\sqrt{3}$

C. 0

D. $\log_e 3\sqrt{3}$

Answer: B



Watch Video Solution

106. IF the angles of a triangle are in the ratio 1:2:3, the corresponding sides are in the ratio

A. $1:\sqrt{2}:1$

B. $1:\sqrt{3}:2$

C. $1:\sqrt{3}:1$

D. $1:1:\sqrt{2}$

Answer: B



Watch Video Solution

107. Two ships leave a port from a point at the same time. One goes with a velocity of 3 km/h along North-East making an angle of 45° with East direction and the other travels with a velocity of 4 km/h along South-East making an angle of 15° with East direction. Then, the distance between the ships at the end of two hours is

A. $2\sqrt{13}$

B. $\sqrt{13}$

C. 5

D. 10

Answer: A

[View Text Solution](#)

108. In $\triangle ABC$, $r_1 + r_2 + r_3 =$

A. $4R$

B. $4R + r$

C. $4R - r$

D. $4R + s^2$

Answer: B

[View Text Solution](#)

109. In a quadrilateral PQRS, A divides SR in the ratio 1 : 3 and B is the mid-point of PR. If $3SR - QR - 3PS - PQ = kAB$, then k=

A. 2

B. 4

C. 6

D. 8

Answer: D



View Text Solution

110. It is given that a, b, c are vectors of lengths 6, 8, 10 respectively. If a is perpendicular to $(b + c)$, b is perpendicular to $(c + a)$, and c is perpendicular to $(a + b)$, then the length of the vector of $|a + b + c|$ is

A. $6\sqrt{2}$

B. $12\sqrt{2}$

C. $5\sqrt{2}$

D. $10\sqrt{2}$

Answer: D



Watch Video Solution

111. If the direction cosines of two lines are given by $l + 3m + 5n = 0$ and $5lm - 2mn + 6ln = 0$, then the angle between the lines is

A. $\cos^{-1}\left(\frac{1}{6}\right)$

B. $\cos^{-1}\left(\frac{1}{3}\right)$

C. $\cos^{-1}\left(\frac{1}{5}\right)$

D. $\sin^{-1}\left(\frac{1}{6}\right)$

Answer: A



Watch Video Solution

112. If a, b, c, d are 4 vector such that $a \cdot b = 0$, $|a \times c| = |a||c|$, $|a \times d| = |a||d|$, then $[bcd] =$

A. $|a||b||c|$

B. $|b||c||d|$

C. $\frac{1}{6}$

D. 0

Answer: D



View Text Solution

113. If a, b, c are three non-coplanar vectors and d is any unit vector, then

$$|(a \cdot d)(b \times c) + (b \cdot d)(c \times a) + (c \cdot d)(a \times b)| =$$

A. $2|[abc]|$

B. $\frac{1}{2} |[abc]|$

C. $|[abc]|$

D. $\frac{1}{6} |[abc]|$

Answer: C



View Text Solution

114. If the line $r = a + lb$ is parallel to the plane $r = c + ld + me$, then

A. $[abc]=0$

B. $[bcd]=0$

C. $[cde]=0$

D. $[bde]=0$

Answer: D



[View Text Solution](#)

115. The mean derviation about the mean for the following data, is

$x,$	2	4	5	7	9
$f,$	2	4	10	8	6

A. 6.3

B. 1.5

C. 2.83

D. 1.733

Answer: D



View Text Solution

116. If the coefficients of variation of two distributions are 40 and 20 and their variances are 144 and 64 respectively, then the mean of their arithmetic means is

A. 40

B. 12

C. 30

D. 35

Answer: D

[View Text Solution](#)

117. A number n is chosen at random from the natural numbers 2 to 1001. The probability that n is a number that leaves remainder 1 when divided by 7, is

A. $\frac{73}{500}$

B. $\frac{71}{1000}$

C. $\frac{143}{1000}$

D. $\frac{71}{500}$

Answer: D

[Watch Video Solution](#)

118. If A and B are two independent events such that $P(B) = \frac{2}{7}$ and $P(A \cup B^C) = 0.8$, then $p(A \cup B) =$

A. $\frac{29}{35}$

B. $\frac{39}{70}$

C. $\frac{1}{2}$

D. $\frac{41}{105}$

Answer: C



Watch Video Solution

119. In a certain recruitment test with multiple choice questions, there are four options to answer each question, out of which only one is correct. An intelligent student knows 90% correct answers while a weak student known only 20% correct answers. If an intelligent student gets the correct answer for a question, then the probability that he was guessing it, is

A. 0.03

B. 0.27

C. 0.4

D. 0.50

Answer: D



Watch Video Solution

120. If the mean and variance of a Binomial variable X are $\frac{5}{2}$ and $\frac{5}{4}$ respectively, then $P(X > 1) =$

A. $\frac{3}{16}$

B. $\frac{11}{16}$

C. $\frac{13}{16}$

D. $\frac{15}{16}$

Answer: C



Watch Video Solution

121. If a random variable X follows a poisson distribution such that

$$P(X = 1) = 3P(X = 2), \text{ then } P(x = 3) =$$

A. $\frac{4}{81}e^{-\frac{2}{3}}$

B. $\frac{2}{81}e^{-\frac{2}{3}}$

C. $\frac{2}{27}e^{-\frac{2}{3}}$

D. $\frac{4}{81}e^{-\frac{1}{3}}$

Answer: A



View Text Solution

122. Let $Q(x_1, y_1)$ be a variable point and $R(1, 0)$ be a point on the circle

$x^2 + y^2 = 1$ and P be the mid-point of QR . Then, the locus of the point

P is

A. $x^2 + y^2 - 2x = 0$

B. $x^2 + y^2 + x = 0$

C. $x^2 + y^2 + 2x = 0$

D. $x^2 + y^2 - x = 0$

Answer: D



View Text Solution

123. The point $P(3,2)$ undergoes the following transformations successively

(i) Reflection about the line $y=x$

(ii) Translation to a distance of 3 units in the positive direction of X-axis

(iii) Rotation through an angle $\frac{\pi}{4}$ about the origin in the anti-clockwise direction

Then , the final position of that point is

A. A (2,4)

B. B $(\sqrt{2}, 4\sqrt{2},)$

C. C $\left(\frac{1}{\sqrt{2}}, \sqrt{2}\right)$

D. D $(\sqrt{2}, 2\sqrt{2})$

Answer: B



Watch Video Solution

124. The equation of the straight line which is perpendicular to the line $5x - 2y = 7$ and passing through the point of intersection of the lines $2x + 3y - 1 = 0$ and $3x + 4y - 6 = 0$ is

A. $2x + 5y - 17 = 0$

B. $2x + 5y + 17 = 0$

C. $2x + 5y + 47 = 0$

D. $2x + 5y - 47 = 0$

Answer: B



Watch Video Solution

125. The angle between the line joining the points $(1,2)$, $(3,2)$ and the line

$$x + 2y - 7 = 0 \text{ is}$$

A. 0

B. $\frac{\pi}{4}$

C. $\frac{\pi}{2}$

D. π

Answer: C



Watch Video Solution

126. The vertices of a triangle are $A(1, 7)$, $B(-5, -1)$ and $C(-1, 2)$.

Then, the equation of a bisector of the $\angle ABC$ is

A. $x - y + 4 = 0$

B. $x + y + 4 = 0$

C. $2x - 3y + 6 = 0$

D. $x - 2y + 4 = 0$

Answer: A



Watch Video Solution

127. Show that two pairs of lines $3x^2 + 8xy - 3y^2 = 0$ and $3x^2 + 8xy - 3y^2 + 2x - 4y - 1 = 0$ forms a square.

A. $\frac{1}{2}$

B. $\frac{1}{4}$

C. $\frac{1}{8}$

D. $\frac{1}{16}$

Answer: D



Watch Video Solution

128. The equation of pair of lines passing through origin and forming an equilateral triangle with the line $3x + 4y - 5 = 0$ is

A. $39x^2 + 11y^2 - 96xy = 0$

B. $x^2 + y^2 - 4xy = 0$

C. $x^2 - 7xy + 12y^2 = 0$

D. $2x^2 + 6xy + y^2 = 0$

Answer: A



Watch Video Solution

129. From a point A(1,0) on the circle $x^2 + y^2 - 2x + 2y + 1 = 0$, a chord AB is drawn and it is extended to a point P such that $AP = 3AB$. The equation of the locus of P is

A. $x^2 + y^2 - 2x + 6y + 1 = 0$

B. $x^2 + y^2 - 2x + 4y + 1 = 0$

C. $x^2 + y^2 - 2x + 8y - 8 = 0$

D. $x^2 + y^2 - 2x + 3y + 1 = 0$

Answer: A



Watch Video Solution

130. Show that the tangent at $(-1, 2)$ of the

circle $x^2 + y^2 - 4x - 8y + 7 = 0$

touches

the circle $x^2 + y^2 + 4x + 6y = 0$ and also

find its point of contact.

A. $\left(0, \frac{1}{3}\right)$

B. $\left(-\frac{1}{3}, 1\right)$

C. $\left(\frac{2}{3}, \frac{1}{3}\right)$

D. $(-1, -1)$

Answer: B

[Watch Video Solution](#)

131. If C_1 and C_2 are the centres of similitude with respect to the circles $x^2 + y^2 - 14x + 6y + 33 = 0$ and $x^2 + y^2 + 30x - 2y + 1 = 0$, then the equation of the circle with C_1C_2 as diameter is

A. $2x^2 + 2y^2 + 30x - 33y - 17 = 0$

B. $2x^2 + 2y^2 - 14x + 9y - 13 = 0$

C. $2x^2 + 2y^2 - 39x + 14y + 74 = 0$

D. $2x^2 + 2y^2 - 24x + 8y - 5 = 0$

Answer: C

[View Text Solution](#)

132. If tangents are drawn to the circle $x^2 + y^2 = 12$ at the points of intersection with the circle $x^2 + y^2 - 5x + 3y - 2 = 0$, then the ordinate of the point of intersection of these tangents is

A. $-\frac{18}{5}$

B. $-\frac{12}{5}$

C. $-\frac{9}{5}$

D. $-\frac{3}{5}$

Answer: A



Watch Video Solution

133. From a point P on the line $4x - 5y = 6$ two tangents are drawn to the circle $x^2 + y^2 - 6x - 4y + 4 = 0$. If the angle between these tangents is $\tan^{-1}\left(\frac{24}{7}\right)$, then P =

A. $\left(1, \frac{-2}{3}\right)$

B. $\left(2, \frac{2}{3}\right)$

C. $\left(-1, \frac{-10}{3}\right)$

D. (6,6)

Answer: D



View Text Solution

134. If $(-1, -1)$ is the focus and $x + y + 3 = 0$ is the directrix of a parabola, then its vertex is

A. A. $\left(-\frac{3}{2}, -\frac{3}{2}\right)$

B. B. $\left(-\frac{5}{2}, -\frac{5}{2}\right)$

C. C. $\left(-\frac{1}{4}, -\frac{1}{4}\right)$

D. D. $\left(\frac{1}{4}, \frac{1}{4}\right)$

Answer: A



Watch Video Solution

135. If a normal chord of a parabola $y^2 = 4ax$ subtends a right angle at the origin, then the slope of that normal chord is

A. ± 2

B. $\pm 2\sqrt{2}$

C. $\pm \frac{1}{\sqrt{2}}$

D. $\pm \sqrt{2}$

Answer: D



Watch Video Solution

136. If origin is the centre, X-axis is the major axis and $\sqrt{\frac{2}{5}}$ is the eccentricity of an ellipse which passes through $(-3, 1)$ then the equation of that ellipse is

A. $3x^2 + 5y^2 = 32$

B. $2x^2 + y^2 = 19$

C. $x^2 + 23y^2 = 32$

D. $x^2 + 2y^2 = 11$

Answer: A



Watch Video Solution

137. The slope of a common tangent to the ellipse $\frac{x^2}{49} + \frac{y^2}{4} = 1$ and the circle $x^2 + y^2 = 16$ is

A. $\frac{5}{\sqrt{11}}$

B. $\frac{4}{\sqrt{11}}$

C. $\frac{3}{\sqrt{11}}$

D. $\frac{2}{\sqrt{11}}$

Answer: D



Watch Video Solution

138. The distance between the tangents drawn to the hyperbola $3x^2 - y^2 = 3$, that are parallel to the line $y = 2x + 4$ is

A. A. $\frac{4}{\sqrt{5}}$

B. B. $\frac{2}{\sqrt{5}}$

C. C. $\frac{2}{3}$

D. D. 1

Answer: B



Watch Video Solution

139. If the distance between two points A and B is d , and the lengths of the projections of AB on the coordinate planes are d_1, d_2, d_3 , then

A. $2d^2 = d_1^2 + d_2^2 + d_3^2$

B. $d_1 + d_2 + d_3 = 0$

C. $d_1^2 + d_2^2 + d_3^2 = d^2$

D. $d_1 + d_2 + d_3 = d$

Answer: A

140. L is a line passing through the point $A(1, 0, -3)$ and parallel to a line having direction ratios $0, 1, -2$, P is a point on the line L which is at a minimum distance from the plane $2x + 3y + 5z = 1$. Then, the equation of the plane through P and perpendicular to AP is

A. $y + 2z = 12$

B. $y - 2z + 4 = 0$

C. $x + y - 2z = 12$

D. $2y - z = 16$

Answer: B

141. Let π_1 be the plane passing through the points $(0,1,2)$, $(1,0,-2)$, $(-2, 1,0)$ and p_2 be the plane passing through the point $(1, 2,3)$ and perpendicular

to the planes $x + y + z = 1$ and $2x - 3y + z = 5$. If θ is the acute angle between the planes π_1 and π_2 then $\cos \theta =$

A. $\frac{\sqrt{14}}{9}$

B. $\frac{\pi}{3}$

C. $\frac{13}{3\sqrt{22}}$

D. $\frac{\pi}{4}$

Answer: A



View Text Solution

142. $\lim_{x \rightarrow 0} \frac{\cos 4x - 4 \cos 2x + 3}{x^4} =$

A. 4

B. 8

C. $\frac{1}{4}$

D. $\frac{1}{8}$

Answer: B



Watch Video Solution

143. If $\alpha = \lim_{x \rightarrow 0} \frac{x \cdot 2^x - x}{1 - \cos x}$ and $\beta = \lim_{x \rightarrow 0} \frac{x \cdot 2^x - x}{\sqrt{1+x^2} - \sqrt{1-x^2}}$, then

A. $\alpha = \beta$

B. $\alpha = 2\beta$

C. $\alpha = \frac{\beta}{2}$

D. $\alpha = 3\beta$

Answer: B



View Text Solution

144. If $f(x) = \frac{2}{4 + 3|x|}$, $x \in \mathbb{R}$, then $f(0) =$

A. A. 0

B. B. $\frac{1}{4}$

C. C. $\frac{1}{2}$

D. D. $\frac{3}{4}$

Answer: C



Watch Video Solution

145. If f is a real function such that $f(4) = 4$ and

$f'(4) = 16$, then $\lim_{x \rightarrow 4} \frac{\sqrt{f(x)} - 2}{\sqrt{x} - 2} =$

A. 16

B. 12

C. 8

D. 2

Answer: A



Watch Video Solution

146. If $y = (\sin^{-1} 2x)^2 + (\cos^{-1} 2x)^2$, then $(1 - 4x^2)y_2 - 4xy_1 =$

A. 0

B. 4

C. 16

D. 12

Answer: C



View Text Solution

147. Let $f(x) = x^3 + 2x^2 - x$ be a real valued function. Then, the value of Lagrange's constant C in $(-1, 2)$ is

A. $\frac{-4 + \sqrt{76}}{3}$

B. $\frac{-2 + \sqrt{19}}{3}$

C. $\frac{-4 + \sqrt{19}}{6}$

D. $\frac{-2 + \sqrt{19}}{6}$

Answer: B



View Text Solution

148. $\int \tan^{-1} \left(\frac{2x}{1-x^2} \right) dx =$

A. $2x \tan^{-1} \left(\frac{2x}{1-x^2} \right) + \log(1+x^2) + c$

B. $x \tan^{-1} \left(\frac{2x}{1-x^2} \right) - \log(1-x^2) + c$

C. $x \tan^{-1} \left(\frac{2x}{1-x^2} \right) - \log(1-x^2) + c$

D. $x^2 \tan^{-1} \left(\frac{x}{1-x^2} \right) + \log(1-x^2) + c$

Answer: C



Watch Video Solution

149. The angle between the curves $y = \sin 2x$ and $y = \cos 2x$ is

A. $\tan^{-1} \sqrt{2}$

B. $\tan^{-1} 2\sqrt{2}$

C. $\tan^{-1} \left(\frac{1}{\sqrt{2}} \right)$

D. $\tan^{-1} \left(\frac{1}{2\sqrt{2}} \right)$

Answer: B



View Text Solution

150. The ratio between the length of substangent at any point other than origin on the parabola $y^2 = 16ax$ and the abscissa of that point is

A. 1:3

B. 1:4

C. 1:2

D. 2:1

Answer: D

[Watch Video Solution](#)

151. $\int \frac{dx}{\sqrt{(x-1)(x-2)}}$

A. $\sin^{-1}(2x+5) + c$

B. $(\sinh^{-1}(2x-5) + c$

C. $\cosh^{-1}(2x-3) + c$

D. $\sin^{-1}(3-2x) + c$

Answer: C

[Watch Video Solution](#)

152. If $\int \frac{x^4+1}{x^6+1} dx = A \tan^{-1} x + B \tan^{-1} x^3 + c$, then $(A,B) =$

A. $\left(1, \frac{1}{3}\right)$

B. $\left(1, \frac{1}{4}\right)$

C. $\left(1, \frac{1}{6}\right)$

D. $\left(1, \frac{4}{3}\right)$

Answer: A



Watch Video Solution

153. If $\int x(1+x)\log(1+x^2)dx = F(x)$

$\log(1+x^2) - \frac{2}{3}\tan^{-1}x - \frac{2x^3}{9} - \frac{x^2}{2} + \frac{2x}{3} + c$, then $F(x) =$

A. $\frac{x^2}{2} + \frac{x^3}{3}$

B. $\frac{x^2}{2} + \frac{x^3}{3} - \frac{1}{3}$

C. $\frac{x^2}{2} + \frac{x^3}{3} + \frac{1}{2}$

D. $\frac{x^2}{2} + \frac{x^3}{3} - \frac{2}{3}$

Answer: C



View Text Solution

154. If $I_n = \int \cos^n x dx$, then $6I_6 - 5I_4 =$

A. $-\cos^5 x \sin^2 x$

B. $\cos^6 x \sin^2 x$

C. $\cos^3 x \sin^2 x$

D. $\cos^5 x \sin x$

Answer: D



Watch Video Solution

155. If $f(x) = \frac{|\log x|}{x^2}$, then $\int_{1/e}^e f(x) dx =$

A. e

B. $1 - \frac{1}{e}$

C. $e^2 \left(1 - \frac{1}{e}\right)$

D. $2 \left(1 - \frac{1}{e}\right)$

Answer: D



View Text Solution

156. The area enclosed by the curves

$y = 8x - x^2$ and $8x - 4y + 11 = 0$ is

A. $\frac{125}{6}$

B. $\frac{32}{3}$

C. 36

D. $\frac{9}{2}$

Answer: A



View Text Solution

157. If $I = \int_0^{\pi/2} \frac{dx}{5 + 3 \sin x} = \lambda \tan^{-1} \left(\frac{1}{2} \right)$, then $\lambda =$

A. $\frac{1}{4}$

B. 1

C. $\frac{1}{2}$

D. $\frac{1}{3}$

Answer: C



View Text Solution

158. The general solution of the differential equation

$$\left(\frac{1}{x^2} + x\right) \frac{dy}{dx} + 3y = 1 \text{ is}$$

A. $y = \frac{1}{x^2} + 3c$

B. $(3y - 1)x^3 + 3y = c$

C. $\log y - xy = c$

D. $(1 + x^3)y = x^3 + c$

Answer: B

[View Text Solution](#)

159. A family of curves whose equation is general solution of a differential equation having order 1 and degree 3, is

(g, a, c are arbitrary constants)

A. $x^2 + y^2 + 2gx + 4y + 2 = 0$

B. $x^2 = a^2(1 + y^2)$

C. $y^2 = 2c(x + \sqrt{c})$

D. $y^2 = 4ax$

Answer: C

[View Text Solution](#)

160. The general solution of the differential equation $\frac{dy}{dx} = \frac{1}{x + y + 1}$ is

(k, c are arbitrary constant)

A. $y = \log_e \left(\frac{x + y + 2}{k} \right)$

B. $x = \log_e \left(\frac{x + y + 2}{k} \right)$

C. $x = ce^y + y + 2$

D. $y = ce^x + x + 2$

Answer: A



View Text Solution

161. If $f: R \rightarrow R$ is defined by $f(x)=[2x]-2[x]$ for $x \in R$, then the range of f is (Here $[x]$ denotes the greatest integer not exceeding x)

A. Z , the set of all integers

B. N , the set of all natural numbers

C. R the set of all real numbers

D. $(0,1)$

Answer: D

162. Given that a, b and c are real numbers such that $b^2 = 4ac$ and $a > 0$.

The maximal possible set $D \subset \mathbb{R}$ on which the function $f: D \rightarrow \mathbb{R}$ given

by

$f(x) = \log(ax^3 + (a+b)x^2 + (b+c)x + c)$ is defined, is

A. $\mathbb{R} - \left\{ -\frac{b}{2a} \right\}$

B. $\mathbb{R} - \left(\left\{ -\frac{b}{2a} \right\} \cup (-\infty - 1) \right)$

C. $\mathbb{R} - \left(\left\{ -\frac{b}{2a} \right\} \cup (x, x \geq 1) \right)$

D. $\mathbb{R} - (\{ -\frac{b}{2a} \} \cup (-\infty - 1))$

Answer: D

163. For any natural number n , $(15 \times 5^{2n}) + (2 \times 2^{3n})$ is divisible by

A. 7

B. 11

C. 13

D. 17

Answer: D



Watch Video Solution

164. For the matrix $A = \begin{bmatrix} 3 & -3 & 4 \\ 2 & -3 & 4 \\ 0 & -1 & 1 \end{bmatrix}$ $A^{-1} =$

A. A

B. A^2

C. A^3

D. A^4

Answer: C

[Watch Video Solution](#)

165. $A = \begin{bmatrix} k/2 & 0 & 0 \\ 0 & l/3 & 0 \\ 0 & 0 & m/4 \end{bmatrix}$ and $A^{-1} = \begin{bmatrix} 1/2 & 0 & 0 \\ 0 & 1/3 & 0 \\ 0 & 0 & 1/4 \end{bmatrix}$, then

$k+l+m=$

- A. 1
- B. 9
- C. 14
- D. 29

Answer: D

[Watch Video Solution](#)

166. If A and B are the two real values of k for which the system of equations $x + 2y + z = 1, x + 3y + 4z = k, x + 5y + 10z = k^2$ is consistent, then $A+B =$

A. 3

B. 4

C. 5

D. 7

Answer: A



Watch Video Solution

167. Let $z=x+iy$ and a point P represent z in the Argand plane. If the real part of $\frac{z-1}{z+i}$ is 1, then a point that lies on the locus of P is

A. (2016, 2017)

B. (-2016, 2017)

C. (-2016, -2017)

D. (2016, -2017)

Answer: D

 [Watch Video Solution](#)

168. If $13e^{i \tan^{-1} \frac{5}{12}} = a + ib$, then the ordered pair (a,b) =

A. (12,5)

B. (5,12)

C. (24,10)

D. (10,24)

Answer: A

 [Watch Video Solution](#)

169. If

$z_1 = 1 - 2i$, $z_2 = 1 + i$ and $z_3 = 3 + 4i$, then $\left(\frac{1}{z_1} + \frac{3}{z_2} \right) \frac{z_3}{z_2} =$

A. 13-6i

B. 13-3i

C. $6 - \frac{13}{2}i$

D. $\frac{13}{2} - 3i$

Answer: D



Watch Video Solution

170. If $1, \omega, \omega^2$ are the cube roots of unity, then prove that

$$\frac{1}{2 + \omega} - \frac{1}{1 + 2\omega} = \frac{1}{1 + \omega}.$$

A. 1

B. ω

C. ω^2

D. 0

Answer: D



Watch Video Solution

171. The number of integral values of x satisfying

$$5x - 1 < (x + 1)^2 < 7x - 3 \text{ is}$$

A. 0

B. 1

C. 2

D. 3

Answer: B



Watch Video Solution

172. For real number x , if the minimum value of $f(x) = x^2 + 2bx + 2c^2$ is greater than the maximum value of $g(x) = -x^2 - 2cx + b^2$, then

A. $c^2 > 2b^2$

B. $c^2 < 2b^2$

C. $b^2 = 2c^2$

D. $c^2 = 2b^2$

Answer: A



Watch Video Solution

173. If a, b and c are the roots of $x^3 + qx + r = 0$, then

$$(a - b)^2 + (b - c)^2 + (c - a)^2 =$$

A. $-6q$

B. $-4q$

C. $6q$

D. $4q$

Answer: A



Watch Video Solution

174. If the sum of two roots of the equation $x^3 - 2px^2 + 3qx - 4r = 0$ is zero, then the value of r is

A. $\frac{3pq}{2}$

B. $\frac{3pq}{4}$

C. pq

D. $2pq$

Answer: A



Watch Video Solution

175. The sum of the four digit even numbers that can be formed with the digits 0,3,5,4 with out repetition is

A. 14684

B. 43536

C. 46526

D. 52336

Answer: B



Watch Video Solution

176. If x is the number of ways in which six women and six men can be arranged to sit in a row such that no two women are together and if y is the number of ways they are seated around a table in the same manner, then $x:y=$

A. 12: 1

B. 42: 1

C. 16: 1

D. 6: 1

Answer: B



Watch Video Solution

177. The number of 5-letter words that can be formed by using the letters of the word SARANAM is

A. 1120

B. 6720

C. 480

D. 720

Answer: C



Watch Video Solution

178. The number of rational terms in the binomial expansion of

$$\left(\sqrt[4]{5} + \sqrt[5]{4}\right)^{100} \text{ is}$$

A. 50

B. 5

C. 6

Answer: C



Watch Video Solution

179. The numerically greatest term in the binomial expansion of $(2a - 3b)^{19}$ when $a = \frac{1}{4}$ and $b = \frac{2}{3}$ is

A. ${}^{19}C_5, 2^{11}$

B. ${}^{19}C_3, 2\frac{1}{11}$

C. ${}^{19}C_4, 2\frac{1}{13}$

D. ${}^{19}C_3, 2^{13}$

Answer: D



Watch Video Solution

180. If $\frac{x^2 + 5x + 7}{(x - 3)^3} = \frac{A}{(x - 3)} + \frac{B}{(x - 3)^2} + \frac{C}{(x - 3)^3}$, then the equation of the line having slope A and passing through the point (B,C) is

A. $x + y - 20 = 0$

B. $x - y + 20 = 0$

C. $x + y + 20 = 0$

D. $x - y - 20 = 0$

Answer: B



Watch Video Solution

181. If $\left(x - \frac{\pi}{3}\right), \cos x \cos\left(x + \frac{\pi}{3}\right)$ are in a harmonic progression, then

$\cos x =$

A. $\frac{3}{2}$

B. 1

C. $\frac{\sqrt{3}}{2}$

D. $\sqrt{\frac{3}{2}}$

Answer: D



Watch Video Solution

182. $\cos^3 110^\circ + \cos^3 10^\circ + \cos^3 130^\circ =$

A. $\frac{3}{4}$

B. $\frac{3}{8}$

C. $\frac{3\sqrt{3}}{8}$

D. $\frac{3\sqrt{3}}{4}$

Answer: C



Watch Video Solution

183. If the general solution of $\sin 5x = \cos 2x$ is of the form $a_n, \frac{\pi}{2}$ or $n = 0, \pm 1, \pm 2, \dots$, then $a_n =$

A. $\frac{2n}{5 + 2(-1)^n}$

B. $\frac{2n + (-1)^n}{5 + 2(-1)^n}$

C. $\frac{2n + 1}{5 + 2(-1)^n}$

D. $\frac{2n - 1}{5 + 2(-1)^n}$

Answer: B



Watch Video Solution

184. $ax + b(\sec(\tan^{-1} x)) = c$ and $ay + b(\sec(\tan^{-1} y)) = c$

The value of $\frac{x + y}{1 - xy}$ is

A. $\frac{2ab}{a^2 - b^2}$

B. $\frac{2ac}{a^2 + c^2}$

C. $\frac{2ab}{a^2 + b^2}$

D. $\frac{2ac}{a^2 - c^2}$

Answer: D



Watch Video Solution

185. $\tanh^{-1} \frac{1}{2} + \coth^{-1} 3 =$

A. $\log \sqrt{6}$

B. $\log 6$

C. $-\log \sqrt{6}$

D. $-\log 6$

Answer: A



Watch Video Solution

186. If the median of a $\triangle ABC$ through A is perpendicular to AC, then

$$\frac{\tan A}{\tan C} =$$

A. $1 + \sqrt{2}$

B. $-\frac{1}{\sqrt{3}} + 1$

C. -2

D. $1 + \frac{2}{\sqrt{3}}$

Answer: C



View Text Solution

187. In $\triangle ABC$, $\tan \frac{A}{2} + \tan \frac{B}{2} =$

A. $\frac{c \cot \frac{C}{2}}{4s}$

B. $\frac{2c \cot \frac{C}{2}}{a + b + c}$

C. $\frac{2c \tan \frac{C}{2}}{s}$

D. $\frac{c \tan \frac{C}{2}}{a + b + c}$

Answer: B



Watch Video Solution

188. In a $\triangle ABC$, D , E and F respectively are the points of contact of the incircle with the sides AB , BC and CA such that $AD = \alpha$, $BE = \beta$ and $CF = \gamma$, then $\frac{\alpha\beta\gamma}{\alpha + \beta + \gamma} =$

A. R^2

B. $2R$

C. $2r$

D. r^2

Answer: D



Watch Video Solution

189. Let a, b and c be three non-coplanar vectors. The vector equation of a line which passes through the point of intersection of two lines, one joining the points $a+2b-5c$, $-a-2b-3c$ and the other joining the points $-4c$, $6a-4b+4c$ is

A. $r = 2a - 4b + 3c + \mu(a - 6b + 4c)$

B. $r = 3a + 6b - c + \mu(a + 2b + c)$

C. $r = 2a + 3b - c + \mu(a + b + c)$

D. $r = -2b + 3c + \mu(a - 4b + 3c)$

Answer: B



View Text Solution

190. In $\triangle PQR$, M is the mid-point of QR and C is the mid-point of PM . If QC when extended meets PR at N , then $\left| \frac{\overline{QN}}{\overline{CN}} \right| =$

A. 1

B. 2

C. 3

D. 4

Answer: D



Watch Video Solution

191. If $a = \hat{i} - 2\hat{j} - 3\hat{k}$, $b = 2\hat{i} + \hat{j} - k$, $c = \hat{i} + 3\hat{j} - 2\hat{k}$, then

$$[(a \times b) \times (b \times c)(b \times c) \times (c \times a)(c \times a) \times (a \times b)] =$$

A. 160000

B. -8000

C. 400

D. -40

Answer: A



View Text Solution

192. If $a = \hat{i} + 2\hat{j} + 3\hat{k}$, $b = -\hat{i} + 2\hat{j} + \hat{k}$, $c = \hat{i} + 2\hat{j} - 2\hat{k}$, n is perpendicular to both a and b and θ is the angle between c and n then $\sin \theta =$

A. $\sqrt{\frac{2}{3}}$

B. $\frac{\sqrt{2}}{3\sqrt{3}}$

C. $\frac{2}{\sqrt{3}}$

D. $\frac{\sqrt{3}}{2}$

Answer: B



Watch Video Solution

193. If a , b and c are mutually perpendicular vectors of the same magnitude, then the cosine of the angle between a and $a + b + c$ is

A. $\frac{1}{\sqrt{2}}$

B. $\frac{1}{\sqrt{3}}$

C. $\frac{1}{2}$

D. $\frac{\sqrt{3}}{2}$

Answer: B



Watch Video Solution

194. If a, b and c are non-coplanar vectors and the four points with position vectors

$2a + 3b - c$, $a - 2b + 3c$, $3a + 4b - 2c$ and $ka - 6b + 6c$ are coplanar, then $k =$

A. 0

B. 1

C. 2

D. 3

Answer: B



Watch Video Solution

195. The mean and the standard deviation of data of 8 items are 25 and 5 respectively. If two items 15 and 25 are added to this data, then the variance of the new data is

A. 29

B. 24

C. 26

D. $\sqrt{29}$

Answer: A



Watch Video Solution

196. A die is rolled three times, the probability of getting a larger number than the previous number each time is

A. $\frac{15}{216}$

B. $\frac{5}{54}$

C. $\frac{13}{216}$

D. $\frac{1}{18}$

Answer: B



Watch Video Solution

197. A man is known to speak the truth 2 out of 3 times . He throws a die and reports that it is a six . The probability that it is actually a six is

A. $\frac{3}{8}$

B. $\frac{1}{7}$

C. $\frac{2}{7}$

D. $\frac{4}{5}$

Answer: B



Watch Video Solution

198. If the probability function of a random variable X is defined by $P(X = k) = a \left(\frac{k+1}{2^k} \right)$ for $k=0,1,2,3,4,5$, then the probability that X takes a prime value is

A. $\frac{13}{20}$

B. $\frac{23}{60}$

C. $\frac{11}{20}$

D. $\frac{19}{60}$

Answer: B



Watch Video Solution

199. If X is a binomial variate with mean 6 and variance 2, then the value of

$P(5 \leq X \leq 7)$ is

A. $\frac{4762}{6561}$

B. $\frac{4672}{6561}$

C. $\frac{5264}{6561}$

D. $\frac{5462}{6651}$

Answer: B



Watch Video Solution

200. Let $A(2,3)$, $B(3,-6)$, $C(5,7)$ be three points. If P is a point satisfying the condition $PA^2 + PB^2 = 2PC^2$, then a point that lies on the locus of P is

A. $(2, -5)$

B. $(-2, 5)$

C. $(13, 10)$

D. $(-13, -10)$

Answer: D



Watch Video Solution

201. If the coordinates of a point P changes to $(2, -6)$ when the coordinate axes are rotated through an angle of 135° , then the coordinates of P in the original system are

A. $(-2, 6)$

B. $(-6, 2)$

C. $(2\sqrt{2}, 4\sqrt{2})$

D. $(\sqrt{2} - \sqrt{2})$

Answer: C



Watch Video Solution

202. If the portion of a line intercepted between the coordinates axes is divided by the point $(2,-1)$ in the ratio of $3:2$, then the equation of that line is

A. $5x - 2y - 20 = 0$

B. $2x - y - 5 = 0$

C. $3x - y - 7 = 0$

D. $x - 3y - 5 = 0$

Answer: D



Watch Video Solution

203. The equation of the line passing through the point of intersection of the lines $2x + y - 4 = 0$, $x - 3y + 5 = 0$ and lying at a distance of $\sqrt{5}$ units from the origin, is

A. $x - 2y - 5 = 0$

B. $x + 2y - 5 = 0$

C. $x + 2y + 5 = 0$

D. $x - 2y + 5 = 0$

Answer: B



View Text Solution

204. The equation of the line joining the centroid with the orthocentre of the triangle formed by the points $(-2, 3)$, $(2, -1)$, $(4, 0)$ is

A. $x + y - 2 = 0$

B. $11x - y - 14 = 0$

C. $x - 11y + 6 = 0$

D. $2x - y - 2 = 0$

Answer: B



Watch Video Solution

205. The lines represented by the equations

$$23x^2 - 48xy + 3y^2 = 0 \text{ and } 2x + 3y + 4 = 0 \text{ form}$$

- A. an isosceles triangle
- B. a right angled triangle
- C. an equilateral triangle
- D. a scalene triangle

Answer: C



View Text Solution

206. If the line $x+2y=k$ intersects the curve

$x^2 - xy + y^2 + 3x + 3y - 2 = 0$ at two points A and B and if O is the origin. Then the condition for $\angle AOB = 90^\circ$ is

A. $k^2 + k + 1 = 0$

B. $k^2 - 2k + 10 = 0$

C. $2k^2 + 9k - 10 = 0$

D. $3k^2 + 8k - 1 = 0$

Answer: C



Watch Video Solution

207. If $2x^2 + 3xy - 2y^2 = 0$ represents two sides of a parallelogram and $3x+y+1=0$ is one of its diagonals, then the other diagonal is

A. $x - 3y + 1 = 0$

B. $x - 3y + 2 = 0$

C. $x - 3y = 0$

D. $3x - y = 0$

Answer: C



Watch Video Solution

208. If the lengths of the tangents drawn from P to the circles $x^2 + y^2 - 2x + 4y - 20 = 0$ and $x^2 + y^2 - 2x - 8y + 1 = 0$ are in the ratio 2:1, then the locus P is

A. $x^2 + y^2 + 2x + 12y + 8 = 0$

B. $x^2 + y^2 - 2x + 12y + 8 = 0$

C. $x^2 + y^2 + 2x - 12y + 8 = 0$

D. $x^2 + y^2 - 2x - 12y + 8 = 0$

Answer: D



Watch Video Solution

209. The equation of a circle touching the coordinate axes and the line $3x - 4y = 12$ is

A. $x^2 + y^2 + 6x + 6y + 9 = 0$

B. $x^2 + y^2 + 6x + 6y - 9 = 0$

C. $x^2 + y^2 - 6x - 6y + 9 = 0$

D. $x^2 + y^2 - 6x - 6y - 9 = 0$

Answer: C



Watch Video Solution

210. The pole of the straight line $9x+y-28=0$ with respect to the circle

$2x^2 + 2y^2 - 3x + 5y - 7 = 0$ is

A. (3,1)

B. (3,-1)

C. (-3,1)

D. (4,-8)

Answer: B



Watch Video Solution

211. The point of intersection of the direct common tangents drawn to the circles $(x + 11)^2 + (y - 2)^2 = 225$ and $(x - 11)^2 + (y + 2)^2 = 25$ is

A. $\left(\frac{-11}{2}, 1\right)$

B. $(-22, 4)$

C. $\left(\frac{11}{2}, -1\right)$

D. $(22, -4)$

Answer: D



Watch Video Solution

212. If the radical axis of the circles $x^2 + y^2 + 2gx + 2fy + c = 0$ and $2x^2 + 2y^2 + 3x + 8y + 2c = 0$ touches the circle $x^2 + y^2 + 2x + 2y + 1 = 0$, then $(4g - 3)(f - 2) =$

A. $g = \frac{3}{4}$ or $f = 2$

B. $g \neq \frac{3}{4}, f = 2$

C. $g = \frac{3}{4}$ or $f \neq 2$

D. $g = \frac{2}{5}$ or $f = 1$

Answer: A



Watch Video Solution

213. The line $y=6x+1$ touches the parabola $y^2 = 24x$. The coordinates of a point P on this line, from which the tangent to $y^2 = 24x$ is perpendicular to the line $y=6x+1$. is

A. $(-1,-5)$

B. $(-2,-11)$

C. $(-6,-35)$

D. $(-7,-41)$

Answer: C



Watch Video Solution

214. A point on the parabola whose focus is $S(1,-1)$ and whose vertex is $A(1,1)$ is

A. $\left(3, \frac{1}{2}\right)$

B. $(1, 2)$

C. $\left(2, \frac{1}{2}\right)$

D. $(2, 2)$

Answer: D



View Text Solution

215. An ellipse having the coordinate axes as its axes and its major axis along Y-axis, passes through the point $(-3,1)$ and has eccentricity $\sqrt{\frac{2}{5}}$.

Then its equation is

A. $3x^2 + 5y^2 - 15 = 0$

B. $5x^2 + 3y^2 - 32 = 0$

C. $3x^2 + 5y^2 - 32 = 0$

D. $5x^2 + 3y^2 - 48 = 0$

Answer: C



Watch Video Solution

216. The product of the perpendicular distances drawn from the points $(3,0)$ and $(-3,0)$ to the tangent of the ellipse $\frac{x^2}{36} + \frac{y^2}{27} = 1$ at $\left(3, \frac{9}{2}\right)$ is

A. 36

B. 27

C. 9

D. 63

Answer: B



View Text Solution

217. The equation of the hyperbola whose asymptotes are $3x+4y-2=0$, $2x+y+1=0$ and which passes through the point $(1, 1)$ is

A. $6x^2 + 11xy + 4y^2 - 30x + 2y + 7 = 0$

B. $6x^2 + 11xy + 4y^2 - x + 2y - 22 = 0$

C. $6x^2 + 11xy + 4y^2 - x + 2y + 22 = 0$

D. $6x^2 + 11xy + 4y^2 - 3x - 7y - 11 = 0$

Answer: B



Watch Video Solution

218. If the orthocentre and the centroid of a triangle are $(-3,5,2)$ and $(3,3,4)$ respectively, then its circumcentre is

- A. $(6,2,5)$
- B. $(6,2,-5)$
- C. $(6,-2,5)$
- D. $(6,-2,-5)$

Answer: A



Watch Video Solution

219. A plane cuts the coordinate axes X,Y,Z at A,B,C respectively such that the centroid of the $\triangle ABC$ is $(6,6,3)$. Then the equation of that plane is

- A. $x + y + z - 6 = 0$
- B. $x + 2y + z - 18 = 0$
- C. $2x + y + z - 18 = 0$

D. $x + y + 2z - 18 = 0$

Answer: D



Watch Video Solution

220. If the foot of the perpendicular drawn from the origin to a plane is $(1,2,3)$, then a point on that plane is

A. $(3,2,1)$

B. $(7,2,1)$

C. $(7,3,-1)$

D. $(6,-3,4)$

Answer: B



Watch Video Solution

221. If $[x]$ denotes the greatest integer $\leq x$, then

$$\lim_{n \rightarrow \infty} \frac{1}{n^3} \{ [1^2 x] + [2^2 x] + [3^2 x] + \dots + [n^2 x] \} =$$

A. $\frac{x}{2}$

B. $\frac{x}{3}$

C. $\frac{x}{6}$

D. 0

Answer: B



Watch Video Solution

$$222. f(x) = \begin{cases} \frac{1 - \sqrt{2} \sin x}{\pi - 4x} & \text{if } x \neq \frac{\pi}{4} \\ k & \text{if } x = \frac{\pi}{4} \end{cases}$$

continuous at $x = (\pi)/(4)$, then $K =$

A. $\frac{1}{4}$

B. 1

C. $\frac{-1}{4}$

D. 2

Answer: A



Watch Video Solution

223. The derivative of $f(x) = x^{\tan^{-1} x}$ with respect to $g(x) = \sec^{-1}\left(\frac{1}{2x^2 - 1}\right)$ is

A. $\frac{1}{2}\sqrt{1-x^2}x^{\tan^{-1} x}\left[\frac{\log x}{1+x^2} + \frac{\tan^{-1} x}{x}\right]$

B. $-\frac{1}{2}\sqrt{1-x^2}x^{\tan^{-1} x}[\log(\tan^{-1} x) + x(1+x^2)\tan^{-1} x]$

C. $\frac{-2\tan^{-1} x\left[\frac{\log x}{1+x^2} + \frac{\tan^{-1} x}{x}\right]}{\sqrt{1-x^2}}$

D. $-\frac{1}{2}\sqrt{1-x^2}x^{\tan^{-1} x}\left[\frac{\log x}{1+x^2} + \frac{\tan^{-1} x}{x}\right]$

Answer: D



View Text Solution

224. If $x = 3 \cos t$ and $y = 4 \sin t$, then $\frac{d^2y}{dx^2}$ at the point $(x_0, y_0) = \left(\frac{3}{2}\sqrt{2}, 2\sqrt{2}\right)$, is

- A. $\frac{4\sqrt{2}}{9}$
- B. $-\frac{4\sqrt{2}}{9}$
- C. $\frac{8\sqrt{2}}{9}$
- D. $-\frac{8\sqrt{2}}{9}$

Answer: D



View Text Solution

225. If $y = \frac{2}{\sqrt{a^2 - b^2}} \tan^{-1} \left[\sqrt{\frac{a-b}{a+b}} \tan \frac{x}{2} \right]$

- A. $\frac{d^2y}{dx^2} \Big|_{x=\frac{\pi}{2}} =$
- B. $\frac{b}{2a^2}$
- C. $\frac{b}{a^2}$

D. $\frac{2b}{a}$

Answer: B



View Text Solution

226. If $f(x) = x^3 + ax^2 + bx + 5\sin^2 x$ is an increasing function on \mathbb{R} , then

A. $a^2 - 3b - 15 < 0$

B. $a^2 - 3b + 15 < 0$

C. $a^2 - 3b - 15 > 0$

D. $a^2 + 3b + 15 > 0$



Watch Video Solution

227. The approximate value of $\cos 31^\circ$ is (Take $1^\circ = 0.0174$)

A. 0.7521

B. 0.866

C. 0.7146

D. 0.8573

Answer: D



View Text Solution

228. If x and y are two positive numbers such that $x+y=32$, then the minimum value of $x^2 + y^2$ is,

A. 500

B. 256

C. 1024

D. 512

Answer: D

229. The constant 'c' of Lagrange's mean value theorem for the function

$f(x) = \frac{2x + 3}{4x - 1}$ defined on $[1, 2]$ is

A. $\frac{1 + \sqrt{15}}{3}$

B. $\frac{1 + \sqrt{21}}{4}$

C. $\frac{5}{3}$

D. $\frac{3}{2}$

Answer: B

230. $\int \frac{\sin 2x dx}{\sin^4 x + \cos^4 x} = \tan^{-1}(f(x)) + c$, then $f\left(\frac{\pi}{3}\right) =$

A. 1

B. 2

C. 3

D. $\frac{1}{3}$

Answer: C



Watch Video Solution

231. $\int \left\{ \frac{(\log x - 1)}{(1 + (\log x)^2)} \right\}^2 dx =$

A. $\frac{\log x}{1 + (\log x)^2} + c$

B. $\frac{x}{x^2 + 1} + c$

C. $\frac{x}{1 + (\log x)^2} + c$

D. $\frac{-x}{1 + (\log x)^2} + c$

Answer: C



Watch Video Solution

232. $\int \frac{dx}{x^3 + 3x^2 + 2x} =$

A. $\log|x| - \log|x + 1| + \log|x + 2| + c$

B. $\log|x| - \log|x + 1| + \log|x + 2| + c$

C. $\frac{1}{2}[\log|x| + \log|x + 1| + \log|x + 2| + c$

D. $\frac{1}{2}\log\left(\frac{[x^2 + 2x]}{(x + 1)^2}\right) + c$

Answer: D



Watch Video Solution

233. For $n \geq 2$, If $I_n = \int \sec^n x dx$, then $I_4 - \frac{2}{3}I_2 =$

A. $\sec^2 x \tan x + c$

B. $\frac{1}{3}\sec^2 x \tan x + c$

C. $\frac{2}{3}\sec^2 x \tan x + c$

D. $\frac{1}{2}\log|\sec x + \tan x| + c$

Answer: B



Watch Video Solution

234. $\lim_{n \rightarrow \infty} \left(\frac{\sqrt{1} + 2\sqrt{2} + 3\sqrt{3} + \dots + n\sqrt{n}}{n^{5/2}} \right) =$

A. 1

B. $\frac{5}{2}$

C. 0

D. $\frac{2}{5}$

Answer: D



Watch Video Solution

235. $\int_0^{\alpha/3} \frac{f(x)}{f(x) + f\left(\frac{\alpha - 3x}{3}\right)} dx =$

A. $\frac{2\alpha}{3}$

B. $\frac{\alpha}{2}$

C. $\frac{\alpha}{3}$

D. $\frac{\alpha}{6}$

Answer: D



Watch Video Solution

236. The area (in sq. units) of the region bounded by the X-axis and the curve $y = 1 - x - 6x^2$ is

A. $\frac{125}{216}$

B. $\frac{125}{512}$

C. $\frac{25}{216}$

D. $\frac{25}{512}$

Answer: A

[Watch Video Solution](#)

237. If m and n are respectively the order and degree of the differential equation of the family of parabolas with focus at the origin and X-axis as its axis, then $mn - m + n =$

A. 1

B. 4

C. 3

D. 2

Answer: C

[Watch Video Solution](#)

238. Solve $\left(1 + e^{\frac{x}{y}}\right)dx + e^{\frac{x}{y}}\left(1 - \frac{x}{y}\right)dy = 0$

A. $Ye^{\frac{y}{x}} + x = c$

B. $ye^{\frac{x}{y}} - x = c$

C. $ye^{\frac{x}{y}} + y = c$

D. $ye^{\frac{x}{y}} + x = c$

Answer: D



Watch Video Solution

239. If $f: R \rightarrow [-1, 1]$ and $g: R \rightarrow A$ are two surjective mappings and

$$\sin\left(g(x) - \frac{\pi}{3}\right) = \frac{f(x)}{2} \sqrt{4 - f^2(x)}, \text{ then } A =$$

A. $\left[0, \frac{2\pi}{3}\right]$

B. $[-1, 1]$

C. $\left(\frac{-\pi}{2}, \frac{\pi}{2}\right)$

D. $(0, \pi)$

Answer: A



Watch Video Solution

240. The domain of the function $f(x) = \sqrt{\frac{4-x^2}{[x]+2}}$ where $[x]$ denotes the greatest integer not more than x , is

- A. $(-\infty, -2) \cup (1, 2)$
- B. $(-\infty, -2) \cup (-1, 2)$
- C. $(-\infty, -2) \cup [-1, 2]$
- D. $(-\infty, -1) \cup (1, 2)$

Answer: C



Watch Video Solution

241. if $a_n = \sqrt{7 + \sqrt{7 + \sqrt{7 + n \dots}}}$ times, then which one of the following is true ?

- A. $a_n > 7, \forall n \geq 1$
- B. $a_n > 3, \forall n \geq 1$

C. $a_n < 4, \forall n \leq 1$

D. $a_n < 3, \forall n \leq 1$

Answer: C



Watch Video Solution

242. If A is square matrix of order 2 and $A^2 + A + 2I = 0$ then

A. A can not be a skew-symmetric matrix

B. $|A + I| = 0$

C. A is one singular and $A^{-1} = (A + I)^{-1}$

D. $|A||A + I| = 2$

Answer: A



View Text Solution

243. If A is square matrix of order 3, then consider the following statements

I. If $|A|=0$, then $|\text{Adj}A|=0$

II. If $|A| \neq 0$, then $|A^{-1}| = |A|^{-1}$

Which of the above statements is / are true ?

A. Both I and II

B. Neither I nor II

C. I only

D. II only

Answer: A



View Text Solution

244. The system of equations $x-2y+3z=5$, $2x-2y+z=0$, $-x+2y-3z=6$ has

A. infinitely many solutions

B. exactly tow solutions

C. unique solution

D. no solution

Answer: D



Watch Video Solution

245. The amplitude of $\sin \frac{\pi}{5} + I\left(1 - \cos \frac{\pi}{5}\right)$ is

A. $\frac{\pi}{15}$

B. $\frac{\pi}{10}$

C. $\frac{\pi}{5}$

D. $\frac{2\pi}{5}$

Answer: B



Watch Video Solution

246. If a point p represents a complex number $z=x+iy$ in the argand plane and if $\frac{z+1}{z+i}$ is a purely real number, then the locus of p is

A. $x + y + 1 = 0$

B. $x^2 + y^2 + x + y = 0$

C. $x^2 + y^2 + 2x + 1 = 0, (x, y) \neq (0, -1)$

D. $x + y + 1 = 0, (x, y) \neq (0, -1)$

Answer: D



Watch Video Solution

247. If ω is complex cube root of unity, the

$$\left[\frac{51 + 73\omega + 87\omega^2}{73 + 87\omega + 51\omega^2} + \frac{51 + 73\omega + 87\omega^2}{87 + 51\omega + 73\omega^2} \right]^{15} =$$

A. 1

B. -1

C. 0

D. 2

Answer: B



Watch Video Solution

248. If $z \in C$ and $iz^3 + 4z^2 - z + 4i = 0$, then a complex root of this euquation having minimum magnitude is

A. $4i$

B. $\frac{1-i}{\sqrt{2}}$

C. $\frac{\sqrt{3}+I}{2}$

D. $\frac{1+i}{\sqrt{2}}$

Answer: B



Watch Video Solution

249. If α and β are the roots of the equation $x^2 - 4x + 5 = 0$ then the quadratic equation whose roots are $\alpha^2 + \beta$ and $\alpha + \beta^2$ is

A. $x^2 + 10x + 34 = 0$

B. $x^2 - 10x + 34 = 0$

C. $x^2 - 10x - 34 = 0$

D. $x^2 = 10x - 34 = 0$

Answer: B



Watch Video Solution

250. $f(x)$ is quadratic expression such that $f(x)$ is negative when $x \in \left(-\infty, -\frac{5}{3}\right) \cup (3, \infty)$ and positive when $x \in \left(-\frac{5}{3}, 3\right)$ $g(x)$ is another quadratic expression such that $g(x)$ is negative when $x \in \left(3, \frac{9}{4}\right)$ and positive when $x \in \mathbb{R} - \left[3, \frac{9}{2}\right]$ Then, the sign of $f(x)g(x)$ is $[0, 5]$ is

- A. positive in $\left[0, \frac{9}{2}\right]$ and negative in $\left(\frac{9}{2}, 5\right)$
- B. positive in $[0, 3] \cup \left(3, \frac{9}{2}\right)$ and negative in $\left(\frac{9}{2}, 5\right]$
- C. positive in $[0, 3) \cup \left(3, \frac{9}{2}\right)$ and negative in $\left(\frac{9}{5}, 5\right]$
- D. positive in $(0, 3) \cup \left(3, \frac{9}{2}\right) \cup \left(\frac{9}{2}, 5\right]$

Answer: B



View Text Solution

251. If $a, b, c \in \mathbb{R}$ are such that $4a + 2b + c > 0$ and $ax^2 + bx + c = 0$ has no real roots, then the value of $(c+a)(c+b)$ is

- A. greater than ab
- B. less than bc
- C. greater than ca
- D. less than $ab + bc + ca$

Answer: A

 [View Text Solution](#)

252. The minimum degree of polynomial equation with rational coefficients having $\sqrt{3} + \sqrt{27}$, $\sqrt{2} + 5i$ as two of its roots is

A. 8

B. 6

C. 4

D. 2

Answer: B

 [Watch Video Solution](#)

253. If all the digits in the number 53426 are permuted in all possible ways and are arranged in decreasing order, then the number having rank 89, is

A. 34265

B. 34256

C. 43526

D. 43265

Answer: A



Watch Video Solution

254. Three parallel straight lines L_1 , L_2 and L_3 lie on the same plane. Consider 5 points on L_1 , 7 points on L_2 and 9 points on L_3 . Then the maximum possible number of triangles formed with vertices these points, is

A. 1330

B. 1200

C. 1201

D. 129

Answer: C



Watch Video Solution

255. If $a > 0$ and the coefficient of x^2 in the expansion of $\left(ax^3 + \frac{c}{x}\right)^6$ is 60, then $ac^2 =$

A. 2

B. 3

C. 4

D. 5

Answer: A



Watch Video Solution

256. If $x = \frac{3}{4.8} + \frac{3.5}{4.8.12} + \frac{3.5.7}{4.8.12.16} + \dots$, then $2x^2 + 5x =$

A. $\frac{7}{8}$

B. 7

C. $\frac{7}{16}$

D. $\frac{7}{4}$

Answer: A



Watch Video Solution

257. If $\frac{3x^2 + 1}{(x^2 + 1)(x^2 + 2)^2} = \frac{Ax + B}{x^2 + 1} + \frac{Cx + D}{x^2 + 2} + \frac{Ex + F}{(x^2 + 2)^2}$, then A

+C +E=

A. 0

B. $\frac{7}{3}$

C. 1

D. $\frac{4}{3}$

Answer: A

[View Text Solution](#)

258. If $\tan\left(\frac{\pi}{4} + \frac{y}{2}\right) = \tan^3\left(\frac{\pi}{4} + \frac{x}{2}\right)$ then $\frac{3 \sin x + \sin^3 x}{1 + 3 \sin^2 x} =$

A. 0

B. 1

C. $\sin 2y$

D. $\sin Y$

Answer: D

[View Text Solution](#)

259.

$$(\cos 252^\circ - \sin 126^\circ)(\cos 252^\circ + \sin 126^\circ)(\sin^2 126^\circ + \sin^2 186^\circ + \sin^2 66^\circ)$$

A. $\frac{3\sqrt{5}}{8}$

B. $\frac{-3\sqrt{5}}{8}$

C. $\frac{-3(\sqrt{5})}{4}$

D. $3\frac{\sqrt{5}}{4}$

Answer: B



Watch Video Solution

260. If α, β, γ are any three angles, then

$$\cos \alpha + \cos \beta - \cos \gamma - \cos(\alpha + \beta + \gamma) =$$

A. $4 \frac{\cos(\alpha + \beta)}{2} \frac{\cos(\beta + \gamma)}{2} \frac{\cos(\gamma + \alpha)}{2}$

B. $4 \frac{\cos(\alpha + \beta)}{2} \frac{\sin(\beta + \gamma)}{2} \frac{\sin(\gamma + \alpha)}{2}$

C. $4 \frac{\cos(\alpha + \beta)}{2} \frac{\sin(\beta - \gamma)}{2} \frac{\sin(\gamma - \alpha)}{2}$

D. $4 \frac{\sin(\alpha + \beta)}{2} \frac{\cos(\gamma + \alpha)}{2}$

Answer: B



Watch Video Solution

261. The general solution of the equation $\sqrt{3 - 5 \sin x + \sin^2 x} + \cos x = 0$ is

A. $n\pi + (-1)^n \frac{\pi}{6}, n \in \mathbb{Z}$

B. $2n\pi \pm \frac{\pi}{6}, n \in \mathbb{Z}$

C. $(2n + 1)\pi - \frac{\pi}{6}, n \in \mathbb{Z}$

D. $2n\pi \pm \left(5\frac{\pi}{6}\right), n \in \mathbb{Z}$

Answer: C



Watch Video Solution

262. Consider the following statements.

I. $\sin^{-1}(y^2 - 4y + 6) + \cos^{-1}(y^2 - 4y + 6) = \frac{\pi}{2}, \forall y \in \mathbb{R}$

II. $\sec^{-1}(y^2 - 4y + 6) + \csc^{-1}(y^2 - 4y + 6) = \frac{\pi}{2}, \forall y \in \mathbb{R}$

Which of the above statements (s) is / are true ?

A. Only I

B. Only II

C. Both I and II

D. Neither I nor II

Answer: B



Watch Video Solution

263. If $\sec \theta \cos h y = \operatorname{cosec} x$ and $\operatorname{cosec} \theta \sinh y = \sec x$, then $\sinh^2 y =$

A. $\cos^2 x$

B. $\cos x$

C. $\sin^2 x$

D. $\sin x$

Answer: A



Watch Video Solution

264. Consider the following statements.

I. In $\triangle ABC$, if $c=6$ and $\cos C = \frac{-11}{25}$ then, $R = \frac{25}{2\sqrt{14}}$

II. In $\triangle ABC$ if $a = 3$, $b = 4$, $c = 6$, then ABC is acute angled triangle

Which of the above statements is / are true ?

A. Only I

B. Only II

C. Both I and II

D. Neither I nor II

Answer: A



Watch Video Solution

265. In $\triangle ABC$, if $a=3$, $b=4$, $c=6$ then $\frac{\cot \frac{A}{2} + \cot \frac{B}{2} + \cot \frac{C}{2}}{\cot A + \cot B + \cot C} =$

A. $\frac{13}{61}$

B. $\frac{169}{61}$

C. $\frac{61}{169}$

D. $\frac{61}{13}$

Answer: B



Watch Video Solution

266. If the reciprocals of the lengths of the sides of a $\triangle ABC$ are in harmonic progression, then its ex-radii r_1, r_2, r_3 are in

A. Arithmetic progression

B. Geometric progression

C. Harmonic progression

D. Arithmetico-geometric progression

Answer: C



Watch Video Solution

267. If P and Q are two points on the curve $y = 2^{x+2}$ such that $OP \cdot \hat{i} = -1$ and $QQ \cdot \hat{i} = 2$ then the magnitude of $(OQ - 4OP)$ is

A. 10

B. 1

C. 5

D. 100

Answer: A



Watch Video Solution

268. P and Q are points on the straight line passing to the vector $2\hat{i} - \hat{j} - \hat{k}$ and parallel to the vector $2\hat{i} - \hat{j} + 2\hat{k}$, If

$AP = AQ = 3$, then the vector equation of the plane OPQ is

A. $r = (s + 5t)\hat{i} + 2s\hat{j} + (t - 3s)\hat{k}$

B. $r = (3\hat{i} + \hat{j} - \hat{k}) + s(2\hat{i} - \hat{j} + 2\hat{k}) + t(5\hat{i} + \hat{k})$

$$\text{C. } r = (s + 5t)\hat{i} + 2s\hat{j} + (5s + t)\hat{k}$$

$$\text{D. } r = (3t - s)\hat{i} + 2s\hat{j} + (t - 3s)\hat{k}$$

Answer: A



View Text Solution

269. Let m be the unit vector orthogonal to the vector $\hat{i} - \hat{j} + \hat{k}$ and coplanar with the vectors $2\hat{i} + \hat{j}$ AND $\hat{j} - \hat{k}$, If $a = \hat{i} - \hat{k}$, then the length of the perpendicular from the origin to the plane $r \cdot m = a \cdot m$ is

$$\text{A. } \frac{1}{\sqrt{26}}$$

$$\text{B. } \frac{1}{\sqrt{5}}$$

$$\text{C. } \frac{5}{\sqrt{26}}$$

$$\text{D. } 1$$

Answer: C



Watch Video Solution

270. If a, b, c are non-coplanar unit vectors such that $a \times (b \times c) = \frac{b + c}{\sqrt{2}}$, then the angle between a and b is

A. $\frac{\pi}{6}$

B. $\frac{\pi}{4}$

C. $\frac{\pi}{2}$

D. $\frac{3\pi}{4}$

Answer: D



Watch Video Solution

271. If a and b are two unit vectors such that $c = (a \times b) + b$ then the maximum value of $[a \ b \ c]$ is

A. 1

B. $\frac{1}{2}$

C. $\frac{3}{2}$

D. 2

Answer: B



View Text Solution

272. If α, β, γ are non-zero vectors such that $|\beta| = |\gamma| = 1$ and $|\alpha| = 10$, then

A. 10

B. 1

C. 0

D. 12

Answer: C



View Text Solution

273. The arithmetich mean and standard deviation of a data of nine numbers are 13 and 5 respectively. If 3 is incoluded as the 10th item of the data, then the variance of the data of ten number is

- A. 23.5
- B. 21.5
- C. 31.5
- D. 27

Answer: C

 Watch Video Solution

274. The variance of the following distribution is

Marks	1-3	3-5	5-7	7-9
Number of students	40	30	20	10

- A. 2

B. 4

C. 6

D. 8

Answer: B



Watch Video Solution

275. A and B are two events such that $P(A)=0.58$ $P(B) =0.32$ and $P(A \cap B) = 0.28$. Then the probability that neither A nor B occurs is

A. 0.38

B. 0.62

C. 0.72

D. 0.9

Answer: A



Watch Video Solution

276. Two dice are thrown simultaneously. If A is event of getting the sum of the numbers on two dice as greater than or equal to 8 and B is the event of getting a number less than or equal to 3 on at least one of the die. Then $P(B/A)=$

A. $\frac{5}{15}$

B. $\frac{6}{15}$

C. $\frac{7}{15}$

D. $\frac{8}{15}$

Answer: B



Watch Video Solution

277. A bag contains 6 balls .if 4 are drawn at a time and all of them are found to be red, then probability that exactly 5 of the balls in the bag are red is

A. $\frac{10}{19}$

B. $\frac{5}{21}$

C. $\frac{1}{21}$

D. $\frac{5}{7}$

Answer: B



Watch Video Solution

278. If the probability distribution of a random variable X is given by

$X = x_i$	0	1	2	3
$P(X = x_i)$	$\frac{1}{8}$	$\frac{3}{8}$	$3K$	K

A. 3

B. $\frac{9}{4}$

C. $\frac{3}{2}$

D. $\frac{3}{4}$

Answer: D



View Text Solution

279. A manufacturer of locks knows that 2 % of his product is defective . If he sells the locks in boxes each with 100 locks and gurantees that not more than 2 locks will be defectiave in a box, then the probability that a box will fail to meet the guranteed quality is

A. $1 - 5e^{-2}$

B. $\sum_{k=2}^{100} {}^{100}C_k \left(\frac{1}{50}\right)^k \left(\frac{49}{50}\right)^{100-k}$

C. 0.02

D. $1 - 3e^{-2}$

Answer: A



Watch Video Solution

280. The equation of the locus of point $(2 \cos \theta - 3, 3 \sin \theta - 4)$ is

A. $9x^2 + 4y^2 + 54x + 32y + 181 = 0$

B. $4x^2 + 2y^2 - 1 = 0$

C. $9x^2 + 4y^2 + 54x + 32y + 109 = 0$

D. $9x^2 + 4y^2 + 54x + 32y + 109 = 0$

Answer: D



Watch Video Solution

281. When the origin is shifted to the point $(2,3)$ and then the coordinate axes are rotated through an angle $\frac{\pi}{3}$ in the counter clockwise sense,

then the transformed equation of

$3x^2 + 2xy + 3y^2 - 18x - 22y + 50 = 0$ is

A. $3x^2 + 3y^2 - 1 = 0$

B. $(6 + \sqrt{3})x^2 - 2xy + (6 - \sqrt{3})y^2 - 2 = 0$

C. $4x^2 + 2y^2 - 1 = 0$

D. $(6 - \sqrt{3})x^2 + (6 + \sqrt{3})y^2 + 2xy = 0$

Answer: B



View Text Solution

282. A straight line L with negative slope passes through the point (1,1) and cuts the positive coordinates axes at the points A and B . If O is the origin , then the minimum value of $OA + OB$ as L varies , is

A. 1

B. 2

C. 3

D. 4

Answer: D



Watch Video Solution

283. If the straight line $L = 3x + 4y - k = 0$ cuts the line segment joining the points $P(2,-1)$ and $Q(1,1)$ in the ratio $4:1$, then the equation of the line parallel to the line $y=x$ and concurrent with the lines PQ and $L=0$ is

A. $2x - 2y + 7 = 0$

B. $x - y + 1 = 0$

C. $5x - 5y - 3 = 0$

D. $y = x + 3$

Answer: C



Watch Video Solution

284. The orthocentre and the centroid of $\triangle ABC$ are $(5, 8)$ and $\left(3, \frac{14}{3}\right)$ respectively. The equation of the side BC is $x-y=0$, Given that the image of the orthocentre of a triangle with respect to any side lies on the

circumcircle of that triangle, then the diameter of the circumcircle of $\triangle ABC$ is

A. $\sqrt{10}$

B. $2\sqrt{10}$

C. $4\sqrt{10}$

D. $8\sqrt{10}$

Answer: C



Watch Video Solution

285. If a pair of perpendicular lines through the origin together with the straight line $2x + 3y = 6$ form an isosceles triangle, then the area of that triangle (in sq units) is

A. $\frac{6}{\sqrt{13}}$

B. $\frac{6}{13}$

C. $\frac{36}{13}$

D. $\frac{27}{13}$

Answer: C



Watch Video Solution

286. If the equation $3x^2 + 7xy + 2y^2 + 2gx + 2fy + 2 = 0$ represents a pair of intersecting lines and the square of the distance of their point of intersection from the origin is $\frac{2}{5}$, then $f^2 + g^2 =$

A. $\frac{25}{4}$

B. 25

C. 50

D. $\frac{25}{2}$

Answer: D



View Text Solution

287. Tangents are drawn to $x^2 + y^2 = 16$ from the point P(0,h) These tangents meet the x- axis at A and B . If the area of triangle P AB is minimum , then

A. $x^2 + y^2 = 6\sqrt{2}$

B. $x^2 + y^2 = 64$

C. $x^2 + y^2 = 32$

D. $x^2 + y^2 = 4\sqrt{2}$

Answer: C



Watch Video Solution

288. If the angle between the tangents drawn to the circle $x^2 + y^2 - 12x - 16y = 0$ at the points where the line $5y = 5x = k$ cut the circle is 60° , then the value of k is

A. $5 + \sqrt{2}$

B. $5(2 \pm 5\sqrt{2})$

C. $2 \pm 5\sqrt{2}$

D. $5 \pm \sqrt{2}$

Answer: B



View Text Solution

289. If a circle S with radius 5 touches the circle $x^2 + y^2 - 6x - 4y - 12 = 0$ at $(-1, -1)$, then the length of the tangent from the centre of the circle S to the given circle is

A. $5\sqrt{3}$

B. $\sqrt{65}$

C. 10

D. $3\sqrt{11}$

Answer: A



Watch Video Solution

290. If a circle S passing through the point (3,4) cuts the circle $x^2 + y^2 = 36$ orthogonally then locus of the centre of S is

A. $x^2 + y^2 - 6x - 8y + 11 = 0$

B. $6x + 8y - 61 = 0$

C. $x^2 + y^2 - 8x - 6y + 11 = 0$

D. $6x + 8y + 11 = 0$

Answer: B



Watch Video Solution

291. The line $x-2=0$ cuts the circle $x^2 + y^2 - 8x - 2y + 8 = 0$ at A and B .

The equation of the circle passing through the points A and B and having least radius is

A. $x^2 + y^2 - 4x + 2y - 1 = 0$

B. $x^2 + y^2 - 4x - 2y = 0$

C. $x^2 + y^2 - 4x - 2y + 1 = 0$

D. $x^2 + y^2 - 4x + 4y = 0$

Answer: B



Watch Video Solution

292. If a perpendicular drawn through the vertex O of the parabola $y^2 = 4ax$ to any of its tangent meets the tangent at N and the parabola at M, then ON.OM=

A. $4a^2$

B. $3a^2$

C. $2a^2$

D. a^2

Answer: A

[View Text Solution](#)

293. Let α_1 and α_2 be the ordinates of two points A and B on a parabola $y^2 = 4ax$ and let α_3 be the ordinate of the point of intersection of its tangents at A and B. Then $\alpha_3 - \alpha_2 =$

A. $\alpha_3 - \alpha_1$

B. $\alpha_3 + \alpha_2 =$

C. α_1

D. $\alpha_1 - \alpha_3$

Answer: D

[Watch Video Solution](#)

294. Equations of the latus recta of the ellipse $9x^2 + 4y^2 - 18x - 8y - 23 = 0$ are

A. $x = -1 \pm \sqrt{5}$

B. $y = 1 \pm \sqrt{5}$

C. $x = 1 \pm \frac{2\sqrt{5}}{3}$

D. $y = 2 \pm \sqrt{5}$

Answer: B



Watch Video Solution

295. The equation of the tangent of the ellipse $4x^2 + 9y^2 = 36$ at the end of the latusrectum lying in the second quadrant, is

A. $\sqrt{5}x - 3y + 1 = 0$

B. $x - 3y + \sqrt{5} = 0$

C. $\sqrt{5}x - 3y + 3 = 0$

D. $\sqrt{5}x - 3y + 9 = 0$

Answer: A

296. If the product of the lengths of the perpendiculars from any point on the hyperbola $16x^2 - 25y^2 = 400$ to its asymptotes is ρ and the angle between the two asymptotes is θ then $\rho \tan \frac{\theta}{2} =$

A. $\frac{400}{41}$

B. $\frac{320}{41}$

C. $\frac{4}{5}$

D. $\frac{25}{16}$

Answer: B

297. A(3,2,-1), B(4,1,1), C(6,2,5) and D(3,3,3) are four points G_1, G_2, G_3 and G_4 respectively are the centroids of the triangle

$\Delta BCD, \Delta CDA, \Delta DAB, \Delta ABC$. The point of concurrence of the lines

AG_1, BG_2, CG_3 , and DG_4 is

A. (4,2,2)

B. (2,4,2)

C. (2,2,4)

D. (2,2,2)

Answer: A



View Text Solution

298. If the direction cosines of two lines satisfy the equation

$l + m + n = 0$ and $2lm + 2ln - mn = 0$, then the acute angle

between those two lines is

A. $\frac{\pi}{6}$

B. $\frac{\pi}{4}$

C. $\frac{\pi}{3}$

D. $\frac{2\pi}{5}$

Answer: C



Watch Video Solution

299. A variable plane passes through a fixed point (α, β, γ) and meets the coordinate axes in A,B, and C. let P_1P_2 and P_3 be the planes passing through A,B,C and parallel to the coordinate planes YZ,ZX,XY respectively . Then, the locus of the points of intersection of the planes P_1P_2 and P_3 is

A. $\alpha x + \beta y + \gamma z = 1$

B. $\frac{\alpha}{x}, \frac{\beta}{y} + \frac{\gamma}{z} = 1$

C. $\alpha x^2 + \beta y^2 + \gamma z^2 = 1$

D. $\alpha\beta x + \beta\gamma y + \alpha\gamma z = 1$

Answer: B



View Text Solution

300. $\lim_{x \rightarrow \frac{\pi}{2}} \frac{1 + \cos 2x}{\cot 3x (3^{\sin 2x} - 1)} =$

A. $\frac{1}{3 \log 9}$

B. $\frac{2}{3 \log 3}$

C. $\frac{1}{3 \log 3}$

D. $\frac{3}{\log 3}$

Answer: C



Watch Video Solution

301. $\lim_{n \rightarrow \infty} n^{-nk}$

$$\left\{ (n+1) \left(n + \frac{1}{2} \right) \left(n + \frac{1}{2} \right) \dots \left(n + \frac{1}{2^{k-1}} \right) \right\}^n =$$

A. 2

B. $e^{2 \left(1 - \frac{1}{2^k} \right)}$

C. $\left(1 - \frac{1}{2^k}\right)$

D. e^2

Answer: B



View Text Solution

302. If a and b ($a > b$) are points of discontinuity of the function

$$f(x) = \begin{cases} 3 - 2x^2 & \text{for } x \leq 0 \\ 2x + 3 & \text{for } 0 < x \leq 1 \\ 2x^2 - 3x & \text{for } 1 < x < 2 \\ 2x - 3 & \text{for } 2 \leq x < 3 \\ |x| & \text{for } \geq 3 \end{cases}$$

then $3a - b =$

A. 3

B. 7

C. 5

D. 1

Answer: C

[View Text Solution](#)

303. For $-1 < x < 1$, if $f(x) = \cos^2\left(\tan^{-1}\sqrt{\frac{1-x}{1+x}}\right)$ then $f'(x) =$

A. $\frac{1}{2}$

B. 1

C. -1

D. $-\frac{1}{2}$

Answer: A

[Watch Video Solution](#)

304. $f: \mathbb{R} \rightarrow \mathbb{R}$ is a function such that $f(0)=1$ and for all $x, y \in \mathbb{R}$ $f(xy+1) = f(xy+1) = f(x)f(y) - f(y) - x + 2$ then $\frac{df}{dx}$ at $x=e$ is

A. 0

B. -1

C. e

D. 1

Answer: D



Watch Video Solution

305. $y = \sin(\log(x^2 + 2x + 1))$

$$\Rightarrow (x + 1)^2 \frac{d^2 y}{dx^2} + (x + 1) \frac{dy}{dx} =$$

A. y

B. $-4y$

C. $4y$

D. $-y$

Answer: B



Watch Video Solution

306. The angle between the curves $y^2 = 4x$ and $x^2 = 4y$ is

A. $\tan^{-1}\left(\frac{1}{2}\right)$

B. $\sin^{-1}\left(\frac{3}{5}\right)$

C. $\cos^{-1}\left(\frac{1}{3}\right)$

D. $\tan^{-1}\left(\frac{2}{3}\right)$

Answer: B



Watch Video Solution

307. If $x > 0$, then $\frac{x}{1+x} - \log(1+x)$

A. is less than zero

B. is greater than zero

C. is equal to zero

D. takes all the real values

Answer: A



Watch Video Solution

308. On the curve $y = x^3$, the point at which the tangent line is parallel to the chord joining the points $(-1, -1)$ and $(2, 8)$ is

A. $(1, 1)$

B. $(2, 8)$

C. $(1, 1)$

D. $(3, 27)$

Answer: C



Watch Video Solution

309. If the petrol burnt in driving a motor boat varies as the cube of the velocity, then the speed (in km/hour) of the boat going against a water

floe of C kms/hour so that the quantity of petro burnt is minimum is

A. $\frac{2C}{3}$

B. $\frac{3C}{2}$

C. $\frac{4C}{3}$

D. $\frac{3C}{4}$

Answer: B



View Text Solution

310. For $x < 1$, $\int \frac{x - x^2}{\sqrt{1 - x}} dx =$

A. $\frac{4}{3}(1 - x)^{3/2} - \frac{2}{5}(1 - x)^{5/2} - 2\sqrt{1 - x} + C$

B. $\frac{4}{3}(1 - x)^{3/2} - \frac{2}{3}(1 - x)^{5/2} - 2\sqrt{1 - x} + C$

C. $\frac{2}{3}(1 - x)^{3/2} - \frac{2}{3}(1 - x)^{5/2} - 2\sqrt{1 - x} + C$

D. $-\frac{2}{15}(1 - x)^{3/2}(2 + 3x) + C$

Answer: D



Watch Video Solution

311. If $\int \frac{dx}{(2 \sin x + \sec x)^4} = At^5 + Bt^6 + Ct^7 + k,$ where $t = (1 + \tan x)^{-1},$ then $A + B + C =$

A. $\frac{-86}{105}$

B. $\frac{-1}{105}$

C. $\frac{-26}{105}$

D. $\frac{-16}{105}$

Answer: D



Watch Video Solution

312. $\int \frac{2x^2 - 1 + x^2\sqrt{x^2 + 4}}{x^2(x^2 + 4)} dx$

A. $\frac{9}{8} \frac{\tan^{-1} x}{2} + \frac{1}{4x} + \frac{\cosh^{-1} x}{2} + C$

B. $\frac{9}{8} \tan^{-1} \frac{x}{2} + \frac{1}{4x} + \sinh^{-1} \frac{x}{2} + c$

C. $\frac{9}{16} \log \left| \frac{x+2}{x-2} \right| + \frac{1}{4x} + \log \left| \frac{x + \sqrt{x^2 + 4}}{2} \right| + C$

D. $\frac{-9}{16} \log \left| \frac{2-x}{2+x} \right| + \frac{1}{4x} + \cosh^{-1} \frac{x}{2} + C$

Answer: B



View Text Solution

313. For $n \geq 2$, let $I_n = \int_0^{\pi/4} \tan^n x dx$ and $F_n = I_n + I_{n-2}$. Then

$F_n - F_{n+1} =$

A. $\frac{1}{n}$

B. $\frac{1}{n-1}$

C. $\frac{1}{n(n-1)}$

D. $1 + n$

Answer: C

[Watch Video Solution](#)

314.

$$\lim_{n \rightarrow \infty} n \left[\frac{1}{(3n^2 + 8n + 4)} + \frac{1}{3n^2 + 16n + 16} + \dots + \frac{1}{15n^2} \right] =$$

A. $\frac{1}{2} \log \frac{9}{5}$

B. $\frac{1}{4} \log \frac{9}{5}$

C. $2 \log \frac{9}{5}$

D. $\frac{1}{4} \log \frac{5}{9}$

Answer: B

[View Text Solution](#)

315. $\int_0^3 |x^2 - 3x + 2| dx$

A. $\frac{3}{2}$

B. $\frac{1}{6}$

C. $\frac{11}{6}$

D. $\frac{11}{2}$

Answer: C



Watch Video Solution

316. OABC is a unit square where O is origin and $B = (1,1)$. The curves $y^2 = x$ and $x^2 = y$ divide the area of the square into three three parts OABO, OBO and OBCO, if a_1, a_2, a_3 are the areas (in sq units) of these parts respectively, then $a_1 + 2a_2 + 3a_3 =$

A. 1

B. 2

C. 6

D. 64

Answer: C



Watch Video Solution

317. The differential equation corresponding to the family of parabola

$y^2 = 4a(x + a)$ where a is the parameter, is

A. $y \left(\frac{dy}{dx} \right)^2 + 2x \frac{dy}{dx} + y = 0$

B. $y \left(\frac{dy}{dx} \right)^2 - 2x \frac{dy}{dx} - y = 0$

C. $y \left(\frac{dy}{dx} \right)^2 + 2x \left(\frac{dy}{dx} \right) - y = 0$

D. $y = 2x \frac{dy}{dx}$

Answer: C



Watch Video Solution

318. The general solution of $\frac{dy}{dx} + y \tan x = 2x + x^2 \tan x$

A. $y - x^2 = c \sec x$

B. $y \cos x = x^2 \sec x + c$

C. $y \sec x = x^2 + c \cos x$

D. $y = x^2 + c \cos x$

Answer: D



Watch Video Solution

319. If $g(x) = x^2 + x - 2$ and $\frac{1}{2}g \circ f(x) = 2x^2 - 5x + 2$, then which is $f(x)$

A. $2x - 3$

B. $2x + 3$

C. $2 + 2x$

D. $2x^2 - 3x - 1$

Answer: A



Watch Video Solution

320. If $f: R \rightarrow A$ defined by $f(x) = \frac{1}{x^2 + 2x + 2}$, $\forall x \in R$ is surjective, then $A =$

A. $[1, \infty]$

B. $(1, \infty)$

C. $[0, 1]$

D. $(0, 1]$

Answer: D



Watch Video Solution

321. If $2(4^{2n+1}) + 3^{3n+1}$ is divisible by k , $k > 1$ for all $n \in N$, then the value of k is

A. 19

B. 17

C. 11

D. 13

Answer: C



Watch Video Solution

322. If $A = \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix}$, then the incorrect option among the following is

A. $A^3 - I = A(A - I)$

B. $(A^3 + I) = A(A^3 - I)$

C. $A^4 - I = A^2 + I$

D. $A^2 + I = A(A^2 - I)$

Answer: D



View Text Solution

323. If $a \neq 1, b \neq -1, c \neq -1$ and the system of equations, $x = a(y + z), y = b(z + x), z = c(x + y)$ has a non-trivial solution, then.

A. $\frac{a}{a+1} + \frac{b}{b+1} + \frac{c}{c+1} = 0$

B. $\frac{a}{a+1} + \frac{b}{b+1} + \frac{c}{c+1} = 1$

C. $\frac{abc}{(a+1)(b+1)(c+1)} = 1$

D. $\frac{a+b+c}{(a+1)(b+1)(c+1)} = 2$

Answer: B



View Text Solution

324. The rank of the following matrix A is

$$A = \begin{bmatrix} 1 & -2 & 3 & -4 \\ 2 & 9 & 4 & 5 \\ 4 & 5 & 10 & -3 \\ 1 & 11 & -1 & 9 \end{bmatrix}$$

A. 4

B. 3

C. 2

D. 1

Answer: B



Watch Video Solution

325. If a complex number z satisfies $|z|^2 + 1 = |z^2 - 1|$, then the locus of z is

A. a circle

B. the real axis

C. the imaginary axis

D. the straight line $y = x$

Answer: C



Watch Video Solution

326. If $\frac{z-i}{z+1}$ is purely imaginary then the locus of $z = x + iy$ is

- A. the circle with centre $\left(\frac{1}{2}, \frac{1}{2}\right)$ and radius $\frac{1}{\sqrt{2}}$
- B. the circle with centre $\left(-\frac{1}{2}, -\frac{1}{2}\right)$ and radius $\frac{1}{\sqrt{2}}$
- C. the point on the circle with centre $\left(\frac{1}{2}, \frac{1}{2}\right)$ and radius $\frac{1}{\sqrt{2}}$,
excluding the point (1, 0) and (0, 1)
- D. the points on the circle with centre $\left(-\frac{1}{2}, -\frac{1}{2}\right)$ and radius $\frac{1}{\sqrt{2}}$,
excluding the origin

Answer: C



Watch Video Solution

327. If a, b, c are non-zero real number with $c \neq 1$ such that

$a^2 + b^2 + c^2 = c$ and if $\alpha = \frac{a + ib}{1 - c}$, then $a^2 + b^2 =$

- A. $\frac{|\alpha|^2}{(1 + |\alpha|^2)^2}$

- B. $\frac{|\alpha|^4}{(1 + |\alpha|^2)^2}$
- C. $\frac{|\alpha|}{1 + |\alpha|^2}$
- D. $\frac{|\alpha|}{1 + |\alpha|}$

Answer: A



Watch Video Solution

328. For $n \in \mathbb{Z}^+$,

$$(1 + \sin \theta + i \cos \theta)^n + (1 + \sin \theta - i \cos \theta)^n =$$

- A. $2^{n+1} \cdot \cos^n \left(\frac{\pi}{4} - \frac{\theta}{2} \right) \cos \left(\frac{n\pi}{4} - \frac{\theta}{2} \right)$
- B. $2^{n+1} \cdot \cos^n \left(\frac{\pi}{4} - \frac{\theta}{2} \right) \sin \left(\frac{n\pi}{4} - \frac{\theta}{2} \right)$
- C. $2^{n+1} \cdot \cos^n \left(\frac{\pi}{4} - \frac{\theta}{2} \right) \cos \left(\frac{n\pi}{4} - \frac{n\theta}{2} \right)$
- D. $2^{n+1} \cdot \cos^n \left(\frac{\pi}{4} - \frac{\theta}{2} \right) \sin \left(\frac{n\pi}{4} - \frac{n\theta}{2} \right)$

Answer: C



Watch Video Solution

329. If the quadratic equation formed by eliminating x from $x^2 + \alpha x + \beta = 0$ and $xy + l(x + y) + m = 0$ has the same roots as that of the given quadratic equation, then the set of values of β is

A. $\{m, \alpha l - m\}$

B. $\{m, l + m\}$

C. $\{m, \alpha l + m\}$

D. $\{m, l - m\}$

Answer: A



Watch Video Solution

330. If x is real, then the range of $\frac{x^2 + 2x + 1}{x^2 + 2x + 7}$ is

A. $[0, 1)$

B. $(-\infty, 0) \cup (1, \infty)$

C. (0, 1)

D. R

Answer: A



Watch Video Solution

331. If $[x]$ denotes the greatest integer not exceeding x , then the values of x satisfying

$$[x]^2 - 7[x] + 12 \leq 0 \text{ are}$$

A. $1 \leq x < 4$

B. $3 \leq x < 5$

C. $-5 < x \leq -3$

D. $2 \leq x \leq 4$

Answer: B



Watch Video Solution

332. If the equation whose roots are p times the roots of $x^4 - 2ax^3 + 4bx^2 + 8ax + 16 = 0$ is a reciprocal equation, then $|p| =$

A. 3

B. 2

C. $\frac{1}{2}$

D. $\frac{1}{3}$

Answer: C



Watch Video Solution

333. The number of three digit numbers in which 9 appears only in one place is

A. 243

B. 234

C. 217

D. 225

Answer: D



Watch Video Solution

334. ${}^{34}C_{10} + 3 \cdot ({}^{34}C_9) + 3 \cdot ({}^{34}C_8) + {}^{34}C_7 =$

A. ${}^{38}C_{10}$

B. ${}^{36}C_{10}$

C. ${}^{37}C_{10}$

D. ${}^{35}C_{10}$

Answer: C



Watch Video Solution

335. The coefficient of x^{50} in the expansion of $(1+x)^{100} + 2x(1+x)^{99} + 3x^2(1+x)^{98} + \dots + 101x^{100}$, is

A. $^{100}C_{50}$

B. $^{101}C_{50}$

C. $^{102}C_{50}$

D. $^{103}C_{50}$

Answer: C



View Text Solution

336. If $\alpha = \frac{5}{2! \times 3} + \frac{5 \times 7}{3! \times 3^2} + \frac{5 \times 7 \times 9}{4! \times 3^3} + \dots$, then $\alpha^2 + 4\alpha =$

A. 21

B. 23

C. 25

D. 27

Answer: B



View Text Solution

337. If $\frac{x^4 + 24x^2 + 28}{(x^2 + 1)^3} = \frac{A}{(x^2 + 1)} + \frac{B}{(x^2 + 1)^2} + \frac{C}{(x^2 + 1)^3}$, then A + C =

A. 12

B. 10

C. 9

D. 6

Answer: D



Watch Video Solution

338. If $\tan \theta = \frac{\cos 25^\circ + \sin 25^\circ}{\cos 25^\circ - \sin 25^\circ}$ and θ is in the third quadrant, then θ
=

A. 200°

B. 205°

C. 225°

D. 250°

Answer: D



Watch Video Solution

339. If

$\cos A = \frac{7}{25}$ and $\frac{3\pi}{2} < A < 2\pi$, then $\cos \frac{A}{4} + \cos \frac{A}{2} - \cos 2A =$

A. $\frac{1}{\sqrt{10}} + \frac{27}{625}$

B. $\frac{3}{\sqrt{10}} - \frac{27}{625}$

C. $\frac{3}{\sqrt{10}} + \frac{27}{625}$

D. $\frac{1}{\sqrt{10}} - \frac{27}{625}$

Answer: A



Watch Video Solution

340. In $\triangle ABC$, if $\cos 3A + \cos 3B + \cos 3C + \cos 3\pi = 0$, then the least value of the sum of two of its angles is

A. $\frac{\pi}{6}$

B. $\frac{2\pi}{3}$

C. $\frac{\pi}{3}$

D. $\frac{\pi}{4}$

Answer: C



Watch Video Solution

341. If θ in the interval $\left(0, \frac{\pi}{2}\right)$ satisfying the equation $\cos 2\theta \cdot \sec^4 \theta + \sec^2 \theta = 0$, then $\sin^2 \theta =$

A. $\frac{1}{3}$

B. $\frac{3}{4}$

C. $\frac{1}{2}$

D. $\frac{2}{3}$

Answer: D



Watch Video Solution

342. $\lim_{n \rightarrow \infty} \sum_{r=1}^n \tan^{-1} \left(\frac{2r}{r^4 + r^2 + 2} \right) =$

A. $\frac{\pi}{4}$

B. $\frac{\pi}{2}$

C. $\frac{-\pi}{4}$

D. $\frac{-\pi}{2}$

Answer: A



View Text Solution

343. $\operatorname{sech}^2\left(\tanh^{-1}\frac{1}{2}\right) + \operatorname{cosech}^2\left(\coth^{-1}3\right) =$

A. $\frac{35}{9}$

B. $\frac{3}{2}$

C. $\frac{25}{4}$

D. $\frac{35}{4}$

Answer: D



Watch Video Solution

344. In $\triangle ABC$, if $a \cos^2 \frac{C}{2} + c \cos^2 \frac{A}{2} = \frac{3b}{2}$ then a,b,c are in

A. $2b = a + c$

B. $b^2 = ac$

C. $\frac{1}{b} = \frac{1}{a} + \frac{1}{c}$

D. $a = c$

Answer: A



Watch Video Solution

345. If s is the semi-perimeter of $\triangle ABC$ and if

$$\frac{s-a}{4} = \frac{s-b}{5} = \frac{s-c}{6}, \text{ then } \sum \sin^2\left(\frac{A}{2}\right) =$$

A. $\frac{74}{25}$

B. $\frac{25}{74}$

C. $\frac{74}{33}$

D. $\frac{25}{33}$

Answer: D

[Watch Video Solution](#)

346. If I is incentre of the triangle ABC , P_1, P_2, P_3 are radii of circum circles of $\Delta^{le}IBC, ICA, IAB$ respectively

A. $2Rr$

B. $2Rr^2$

C. $2R^2r$

D. $\frac{4R}{r}$

Answer: C

[Watch Video Solution](#)

347. If a and b are two unit vectors such that $a + b$ is also a unit vector, then $|a - b|^2 =$

A. 1

B. 2

C. 3

D. 0

Answer: C



Watch Video Solution

348. If the line joining the points $\hat{i} + \hat{j}$ and $3\hat{i} + \hat{j} - \hat{k}$ meets the plane that passes through the point $2\hat{i} + 4\hat{j}$ and parallel to the vectors $3\hat{i} + 5\hat{k}$ and $3\hat{i} - \hat{k}$ at, P, then the position vector of the point P is

A. $-27\hat{i} + \hat{j} + 14\hat{k}$

B. $29\hat{i} + \hat{j} - 14\hat{k}$

C. $-14\hat{i} + 89\hat{j} + 3\hat{k}$

D. $2\hat{i} + 5\hat{j} - 7\hat{k}$

Answer: B

[View Text Solution](#)

349. If A, B, C and D are four points in the plane such that

$$|AB|^2 + |CD|^2 = |BC|^2 + |DA|^2 = 100, \text{ then } AC \cdot BD =$$

A. 10

B. 0

C. $\frac{1}{10}$

D. 1

Answer: B

[View Text Solution](#)

350. If $a = \hat{i} + 2\hat{j} + 3\hat{k}$, $b = 2\hat{i} + 3\hat{j} + 2\hat{k}$, and c is a vector perpendicular to b, then

$$\left| \left\{ \frac{a \cdot (b \times c)}{|b \times c|^2} \right\} (b \times c) + \left\{ \frac{a \cdot b}{|b|^2} \right\} b + \left\{ \frac{a \cdot c}{|c|^2} \right\} \right| =$$

A. $\sqrt{14}$

B. 14

C. 13

D. $\sqrt{17}$

Answer: A



View Text Solution

351. If a, b and c are three vectors such that $|a|=1$, $|b|=2$, $|c|=3$ and $a \cdot b = b \cdot c = c \cdot a = 0$, then $|[a \ b \ c]|$ is equal to

A. 15

B. 14

C. 12

D. 8

Answer: A

352. If $a = \hat{i} + \hat{j}$, $b = \hat{j} + \hat{k}$, $c = \hat{i} + \hat{k}$, then

$$\frac{[(b \times c) \times (c \times a)(c \times a) \times (a \times b)(a \times b) \times (b \times c)]}{[b + c \quad c + a \quad a + b][b \times c \quad c \times a \quad a \times b]}$$

A. 0

B. 1

C. $\sqrt{3}$

D. $\sqrt{2}$

Answer: B

353. The standard deviation of the scores 505, 510, 515, 520,, 595 is

A. $500 + 5\sqrt{30}$

B. $505 + \sqrt{30}$

C. $5\sqrt{30}$

D. $5 + \sqrt{30}$

Answer: C



Watch Video Solution

354. If the variance of the distribution

x_i	4	8	11	17	20	24	32
f_i	3	5	9	5	4	3	1

is 45.8, then the variance of the distribution.

x_i	10	18	24	36	42	50	66
f_i	3	5	9	5	4	3	1

A. 93.6

B. $\sqrt{93.9}$

C. 183.2

D. $\sqrt{183.2}$

Answer: C



Watch Video Solution

355. A die is formed so that the probability of getting a number i when it is rolled is proportional to i . ($i = 1, 2, 3, 4, 5, 6$). The probability of getting an odd number on the die when it is rolled is

A. $\frac{1}{2}$

B. $\frac{4}{7}$

C. $\frac{2}{7}$

D. $\frac{3}{7}$

Answer: D



Watch Video Solution

356. A problem is given to 3 students A, B and C whose chances of solving it are $\frac{1}{2}$, $\frac{1}{3}$ and $\frac{1}{4}$ respectively. Then, the probability of the problem being solved by exactly one of them, if all the three try independently, is

A. $\frac{3}{4}$

B. $\frac{11}{24}$

C. $\frac{23}{24}$

D. $\frac{1}{4}$

Answer: B



Watch Video Solution

357. In a certain recruitment test with multiple choice questions, there are four options to answer each question, out of which only one is correct. An intelligent student knows 90% correct answers while a weak student known only 20% correct answers. If an intelligent student gets

the correct answer for a question, then the probability that he was guessing it, is

A. $\frac{1}{37}$

B. $\frac{1}{10}$

C. $\frac{9}{37}$

D. $\frac{1}{2}$

Answer: A



Watch Video Solution

358. A random variable X has the following distribution

Values of $X(x)$	0	1	2	3	4	5	6	7
$P(X = x)$	0	k	2k	2k	3k	k^2	$2k^2$	$7k^2 + k$
Then $P(0 < X < 6) =$								

A. $\frac{9}{10}$

B. $\left(\frac{9}{10}\right)^2$

C. $\left(\frac{3}{10}\right)$

D. $\frac{1}{10}$

Answer: B



Watch Video Solution

359. An unbiased coin is tossed n times. If the probability of getting atleast one head is greater than 0.8, then the least value of n is

A. 2

B. 3

C. 4

D. 5

Answer: B



Watch Video Solution

360. If a point P moves such that the sum of the distance from P to the point A(1, -1) and B(-1, 1) is always 4, then the equation for the locus of P is

A. $16x^2 - 64x + 7y^2 = 48$

B. $3x^2 + 2xy + 3y^2 = 8$

C. $6x + 4y = 3$

D. $x^2 + y^2 - 8x + 6y = 0$

Answer: B



View Text Solution

361. The transformed equation of $3x^2 - 6xy + 8y^2 = 8$ when the axes are rotated about the origin through an angle $\frac{\pi}{4}$ in the positive direction, is

A. $5x^2 + 10xy + 17y^2 + 16 = 0$

B. $5x^2 + 10xy + 17y^2 - 16 = 0$

C. $5x^2 - 10xy + 17y^2 - 16 = 0$

D. $5x^2 - 10xy + 17y^2 + 16 = 0$

Answer: B



Watch Video Solution

362. A variable line passing through a fixed point (α, β) intersects the coordinate axes at A and B. If O is the origin, then the locus of the centroid of the $\triangle OAB$ is

A. $\beta x + \alpha y - 2\alpha\beta = 0$

B. $\beta x + \alpha y - 3xy = 0$

C. $\alpha x + \beta y - (\alpha^2 + \beta^2) = 0$

D. $\beta x + \alpha y + 3xy = 0$

Answer: B

[Watch Video Solution](#)

363. If $m = 1$ is the slope of a line L , then the product of the slopes of non-parallel lines which are inclined at an angle of 60° with L is

A. 1

B. -1

C. $\sqrt{3}$

D. $-\frac{1}{2}$

Answer: A

[Watch Video Solution](#)

364. The area (in sq units) of the quadrilateral formed by the lines $2x + 3y + 6 = 0$, $2x - 3y + 6 = 0$, $2x + 3y - 6 = 0$ and $2x - 3y - 6 = 0$ is

A. 12

B. 36

C. 6

D. 18

Answer: A



Watch Video Solution

365. If the straight lines $2x+3y-1=0$, $x+2y-1=0$, and $ax+by-1=0$ form a triangle with the origin as orthocenter, then (a,b) is given by

A. $(-8, 8)$

B. $(0, 7)$

C. $(6, 4)$

D. $(-3, 3)$

Answer: A



Watch Video Solution

366. The product of the perpendicular distances from $(1, -1)$ to the pair of lines $x^2 - 4xy + y^2 = 0$, is

A. 1

B. $\frac{2}{3}$

C. $\frac{3}{2}$

D. 2

Answer: C



Watch Video Solution

367. If a circle touches the lines $3x - 4y - 10 = 0$ and $3x - 4y + 30 = 0$ and its centre lies on the line $x + 2y = 0$, then the equation of the circle is

A. $x^2 + y^2 + 4x - 2y - 11 = 0$

B. $x^2 + y^2 + 2x - 4y - 11 = 0$

C. $x^2 + y^2 - 4x + 2y - 11 = 0$

D. $x^2 + y^2 + 2x - y - 11 = 0$

Answer: A



Watch Video Solution

368. The line $4x+4y-11=0$ intersects the circle $x^2 + y^2 - 6x - 4y + 4 = 0$ at A and B. The point of intersection of the tangents at A,B is

A. $(-1, 2)$

B. $(-1, -2)$

C. $(2, 1)$

D. $(-2, -1)$

Answer: D



Watch Video Solution

369. The equation of the circle which passes through the point (3, 2) bisects the circumference of the circle $x^2 + y^2 = 15$ and cuts the circle $x^2 + y^2 + 4x + 6y + 3 = 0$ orthogonally is

A. $x^2 + y^2 + 6x + 8y - 43 = 0$

B. $x^2 + y^2 + 6x - 8y - 15 = 0$

C. $x^2 + y^2 - 6x + 8y - 11 = 0$

D. $x^2 + y^2 - 6x - 8y + 21 = 0$

Answer: B



Watch Video Solution

370. $x^2 + y^2 + 2x + 4y - 20 = 0$ and $x^2 + y^2 + 6x - 8y + 10 = 0$ are the given circles. Which one of the following is correct?

A. They intersect orthogonally and will have two common tangents.

The length of their common chord is $\frac{5\sqrt{3}}{\sqrt{2}}$

B. They intersect at right angles and will have two common tangents.

The length of their common chord is 2

C. They do not intersect orthogonally and will have three common tangents. The length of their direct common tangent is 5

D. They touch each other internally and will have only one common tangent

Answer: A



View Text Solution

371. Find the equation of the circle passing through the points of intersections of circles $x^2 + y^2 + 6x + 4y - 12 = 0$, $x^2 + y^2 - 4x - 6y - 12 = 0$, and having radius $\sqrt{13}$.

A. $x^2 + y^2 + 6x + 8y + 12 = 0$

B. $x^2 + y^2 + 8x + 6y - 12 = 0$

C. $x^2 + y^2 + 6x + 8y - 12 = 0$

D. $x^2 + y^2 - 6x - 8y - 12 = 0$

Answer: C



Watch Video Solution

372. If a parabola passess through the points $(-2, 1)$, $(1, 2)$ and $(-1, 3)$ having horizontal axis, then the length of the latus rectum of that parabola is

A. 5

B. $\frac{5}{2}$

C. $\frac{2}{5}$

D. $\frac{1}{5}$

Answer: C



View Text Solution

373. The equation of one of the common tangents of the circle

$x^2 + y^2 - 6y + 4 = 0$ and the parabola $y^2 = x$ is

A. $2x - y + 1 = 0$

B. $2x - y = 1$

C. $4x - y + 1 = 0$

D. $x - 2y + 1 = 0$

Answer: D



View Text Solution

374. The equation of the ellipse having a vertex at $(6, 1)$, a focus at $(4, 1)$

and the eccentricity $\frac{3}{5}$ is

A. $\frac{(x - 1)^2}{16} + \frac{(y - 1)^2}{25} = 1$

B. $\frac{(x - 1)^2}{25} + \frac{(y - 1)^2}{16} = 1$

C. $\frac{(x + 1)^2}{25} + \frac{(y + 1)^2}{16} = 1$

D. $\frac{(x+1)^2}{16} + \frac{(y+1)^2}{25} = 1$

Answer: B



View Text Solution

375. If the tangent at the point $\left(4\cos 2\theta, \frac{16}{\sqrt{11}}\sin 2\theta\right)$ on the ellipse $16x^2 + 11y^2 = 256$ touches the circle $x^2 + y^2 - 2x = 15$, then $\theta =$

A. $\pm \frac{\pi}{3}$

B. $\pm \frac{\pi}{6}$

C. $\pm \frac{\pi}{4}$

D. $\pm \frac{\pi}{8}$

Answer: B



Watch Video Solution

376. A tangent to the curve $9b^2x^2 - 4a^2y^2 = 36a^2b^2$ makes intercepts of unit length on each of the coordinate axes, then the point (a, b) lies on

A. $x^2 - y^2 = 1$

B. $x^2 + y^2 = 1$

C. $4x^2 - 9y^2 = 1$

D. $4x^2 + 9y^2 = 1$

Answer: C



Watch Video Solution

377. The harmonic conjugate of P (-9, 12, -15) with respect to the line segment AB, where A = (1, -2, 3) and B = (-4, 5, -6) is

A. $\left(-\frac{2}{3}, \frac{1}{3}, 0\right)$

B. (6, -9, 12)

C. $\left(-\frac{7}{3}, \frac{8}{3}, -3\right)$

D. $\left(\frac{7}{3}, -\frac{8}{3}, \frac{9}{3}\right)$

Answer: C



Watch Video Solution

378. If the direction ratios of the lines L_1 and L_2 are 2, -1, 1 and 3, -3, 4 respectively, then the direction cosines of a line that is perpendicular to both L_1 and L_2 are

A. $\pm \frac{2}{\sqrt{6}}, \pm \frac{1}{\sqrt{6}}, \pm \frac{1}{\sqrt{6}}$

B. $\pm \frac{1}{\sqrt{35}}, \pm \frac{5}{\sqrt{35}}, \pm \frac{3}{\sqrt{35}}$

C. $\pm \frac{3}{\sqrt{34}}, \pm \frac{3}{\sqrt{34}}, \pm \frac{4}{\sqrt{34}}$

D. $\pm \frac{1}{\sqrt{14}}, \pm \frac{2}{\sqrt{14}}, \pm \frac{3}{\sqrt{14}}$

Answer: B



View Text Solution

379. If the equation of the plane bisecting the line segment joining the points P (3, 2, 4) and Q (-1, 0, -2) and perpendicular to PQ is $ax + by + cz + d = 0$, then $ac + bd$

A. A 0

B. B 12

C. C 6

D. D 1

Answer: A



Watch Video Solution

380.
$$\cos \left[\lim_{x \rightarrow \infty} \frac{2\pi|x| + \pi x}{|x| - 3x} + \lim_{x \rightarrow 0} \frac{\cos\left(\frac{\pi}{2} \cos^2 x\right)}{x^2} \right]$$

A. 1

B. -1

C. 0

D. $\frac{1}{\sqrt{2}}$

Answer: B



Watch Video Solution

381. $\lim_{n \rightarrow \infty} \frac{[6^2 + 12^2 + 18^2 + \dots + (6n)^2]^2}{[5 + 10 + 15 + \dots + 5n][2^3 + 4^3 + 6^3 + \dots + 8n^3]}$

A. $\frac{4}{5}$

B. $\frac{144}{5}$

C. $\frac{4}{25}$

D. $\frac{144}{25}$

Answer: B



View Text Solution

382. If $[x]$ denotes the greatest integer not exceeding the number x , then

$f(x)$ defined by

$$f(x) = \begin{cases} [x] & \text{if } x < 2 \\ [x] - 1 & \text{if } x \geq 2 \end{cases} \text{ is continuous in the interval.}$$

A. $[1, 2] \cup (2, 3)$

B. $[1, 3)$

C. $(1, 3)$

D. \mathbb{R}

Answer: B



Watch Video Solution

383. $f(x) = \left(\frac{a+x}{1+x} \right)^{a+1+2x}$ show that

$$f'(0) = a^{a+1} \left(2 \log a + \frac{1-a^2}{a} \right)$$

A. a^{a+1}

B. $a^{a+1} \left\{ \frac{1-a^2}{a} + 2 \log a \right\}$

C. $2\log a$

D. $a^{a+1} \left\{ \frac{(1+a)^2}{a-2\log a} \right\}$

Answer: B



Watch Video Solution

384. If $y = \log_2(\log_2 x)$, then $\frac{dy}{dx} =$

A. $\frac{\log_2 e}{2x \log_e x}$

B. $\frac{1}{x \log_e x \log_e 2}$

C. $\frac{1}{\log_e (2x)^x}$

D. $\frac{1}{\log_e e \log_e x}$

Answer: B



Watch Video Solution

385. If $y = x \log\left(\frac{x}{2-3x}\right)$ for $0 < x < \frac{2}{3}$, then $\frac{d^2y}{dx^2}$ at $x = \frac{1}{2}$ is

- A. 4
- B. 16
- C. 32
- D. 2

Answer: C



View Text Solution

386. An approximate value of $\sqrt[4]{18}$ is

- A. 2.0512
- B. 2.0425
- C. 2.0625

D. 2.0834

Answer: C



Watch Video Solution

387. The sum of the maximum and the minimum values of $3x^4 - 2x^3 - 6x^2 + 6x + 4$, in $(0, 2)$ is

A. 28

B. $\frac{167}{16}$

C. $\frac{134}{15}$

D. $\frac{87}{16}$

Answer: B



View Text Solution

388. The constant c of Lagrange's mean value theorem for

$$f(x) = \cos x - \sin 2x \text{ in } \left[-\frac{\pi}{2}, \frac{\pi}{2}\right] \text{ is}$$

A. 0

B. $\sin^{-1}\left(\frac{1 \pm \sqrt{33}}{8}\right)$

C. $\cos^{-1}\left(\frac{1 \pm \sqrt{33}}{8}\right)$

D. $\pm \frac{\pi}{4}$

Answer: B



View Text Solution

389. Air is discharging from a large spherical balloon at the rate of 4 cubic meters per minute. Then, the rate at which the surface area is shrinking when the radius of the balloon is 8 m, is

A. $2m^2 / \text{min}$

B. $1m^2 / \text{min}$

C. $4m^2 / \text{min}$

D. $8m^2 / \text{min}$

Answer: B



Watch Video Solution

390. Evaluate: $\int \frac{x}{x^2 + 3x + 2} dx$

A. $\frac{2}{9} \log \left| \frac{x-1}{x+2} \right| + c$

B. $\frac{2}{9} \log \left| \frac{x+2}{x-1} \right| + c$

C. $\frac{1}{3} \frac{1}{x-1} + \frac{2}{9} \log \left| \frac{x-1}{x+2} \right| + c$

D. $-\frac{1}{3} \frac{1}{(x-1)} + \frac{2}{9} \log \left| \frac{x-1}{x+2} \right| + c$

Answer: D



Watch Video Solution

391. $\int \frac{\sqrt{x^2 + 1} [\log(x^2 + 1) - 2\log x]}{x^4} dx$ is equal to

- A. $\frac{1}{9} \left(1 + \frac{1}{x^2}\right)^{\frac{3}{2}} \left[2 - 3\log\left(1 + \frac{1}{x^2}\right)\right] + c$
- B. $\frac{1}{3} \left(1 + \frac{1}{x^2}\right)^{\frac{1}{2}} \left[6 - \log\left(1 + \frac{1}{x^2}\right)^2\right] + c$
- C. $\frac{1}{9} \left(1 + \frac{1}{x^2}\right) \left[3 - 2\log\left(1 + \frac{1}{x^2}\right)^{\frac{1}{2}}\right] + c$
- D. $\frac{1}{3} \left(1 + \frac{1}{x^2}\right)^{\frac{3}{2}} \left[3 - \log\left(1 + \frac{1}{x^2}\right)\right] + c$

Answer: A



Watch Video Solution

392. Evaluate the following integrals

$$\int \frac{dx}{\sin x + \sin 2x}$$

- A. $\frac{1}{6} \log(1 - \cos x) + \frac{1}{2} \log(1 + \cos x) + \frac{2}{3} \log|1 + 2 \cos x| + c$
- B. $\frac{1}{6} \log(1 - \cos x) - \frac{1}{2} \log(1 + \cos x) - \frac{2}{3} \log|1 + 2 \cos x| + c$
- C. $\frac{1}{6} \log(1 - \cos x) + \frac{1}{2} \log(1 + \cos x) - \frac{2}{3} \log|1 + 2 \cos x| + c$

D. $\frac{1}{6} \log[(1 - \cos x)(1 + \cos x)|1 + 2 \cos x|] + c$

Answer: C



Watch Video Solution

393. $\int (\sec^4 x + \tan^4 x) dx =$

A. $\frac{2}{3} \tan^3 x - \frac{2}{3} \tan x + x + c$

B. $\frac{1}{3} \sec^2 x \tan x + \frac{5}{3} \tan x + \frac{\tan^3 x}{3} + x + c$

C. $\frac{2}{3} \tan^3 x + x + c$

D. $\frac{1}{3} \sec^2 x \tan x - \frac{5}{3} \tan x + \frac{\tan^3 x}{3} + x + c$

Answer: C



View Text Solution

394.

$$\lim_{n \rightarrow \infty} \frac{1}{n} \left\{ \sin^5 \left(\frac{\pi}{6n} \right) + \sin^5 \left(\frac{2\pi}{6n} \right) + \sin^5 \left(\frac{3\pi}{6n} \right) + \dots + \sin^5 \left(\frac{\pi}{2} \right) \right\} =$$

A. $\frac{8}{15\pi}$

B. $\frac{8}{5\pi}$

C. $\frac{32}{5\pi}$

D. $\frac{16}{5\pi}$

Answer: D



View Text Solution

395. $\int_0^{\frac{\pi}{2}} \log(\sin 2x) dx$

A. $\pi \log 2$

B. $-\pi \log 2$

C. $\frac{\pi}{2} \log 2$

D. $-\frac{\pi}{2}\log 2$

Answer: D



Watch Video Solution

396. The area (in sq units) between the curve $y^2 = 8x$ and its latus rectum is

A. $\frac{32}{3}$

B. $\frac{64}{3}$

C. $\frac{16}{3}$

D. $\frac{8\sqrt{2}}{3}$

Answer: A



Watch Video Solution

397. The order and the degree of the differential equation

$$y = px + \sqrt{a^2 p^2 + b^2}, \text{ (where } p = \frac{dy}{dx} \text{) are respectively.}$$

A. 2, 1

B. 1, 2

C. 1, 1

D. 2, 2

Answer: B



Watch Video Solution

398. The solution of the differential equation $\frac{dx}{dy} + 2yx = 2y$ which passes through the point (2, 0) is

A. $(x - 1) = 2e^{y^2}$

B. $(x - 1) = 2e^{-y^2}$

C. $(x - 1) = e^{y^2}$

D. $(x - 1) = e^{-y^2}$

Answer: D



Watch Video Solution