

# MATHS

# **BOOKS - CENGAGE MATHS (ENGLISH)**

## **SCALER TRIPLE PRODUCTS**



**1.** Number of integral value(s) of  $\lambda$  for which vectors  $x^2\hat{i} - \hat{j} + x\hat{k}, (\lambda - 1)\hat{i} - 2\lambda\hat{j} - \hat{k}$  and  $\hat{i} - \hat{j} + \hat{k}$ , in the order form right-handed system  $\forall x \in \mathsf{R}$ , is

(a) 0 (b) 2 (c) 4 (d) 6

A. 0

B. 2

C. 4

D. 6

Answer: A

**Watch Video Solution** 

2. Let 
$$\overrightarrow{a}, \overrightarrow{b}, \overrightarrow{c}$$
 be three linearly independent vectors, then  

$$\left[\overrightarrow{a}+2\overrightarrow{b}-\overrightarrow{c}2\overrightarrow{a}+\overrightarrow{b}+\overrightarrow{c}4\overrightarrow{a}-\overrightarrow{b}+5\overrightarrow{c}\right]$$

$$\left[\overrightarrow{a}\overrightarrow{b}\overrightarrow{c}\right]$$

A. 0

B. 1

C. 2

D. - 1

### Answer: A



**3.** If 
$$\overrightarrow{a}, \overrightarrow{b}$$
 are two unit vectors such that  $\overrightarrow{a} + \left(\overrightarrow{a} \times \overrightarrow{b}\right) = \overrightarrow{c}$ , where  $\left|\overrightarrow{c}\right| = 2$ , then value of  $\left[\overrightarrow{a}, \overrightarrow{b}, \overrightarrow{c}\right]$  is

A. 0

 $\mathsf{B.}\pm 1$ 

C.-3

D. 3

### Answer: D

**Watch Video Solution** 

**4.** Let  $\overrightarrow{a}$  and  $\overrightarrow{b}$  be unit vectors that are perpendicular to each other I. then  $\left[\overrightarrow{a} + \left(\overrightarrow{a} \times \overrightarrow{b}\right)\overrightarrow{b} + \left(\overrightarrow{a} \times \overrightarrow{b}\right)\overrightarrow{a} \times \overrightarrow{b}\right]$  will

always be equal to

A. 1

B. zero

C. -1

D. 3

#### Answer: A



5. If V is the volume of the parallelepiped having three coterminous edges as  $\overrightarrow{a}$ ,  $\overrightarrow{b}$  and  $\overrightarrow{c}$ , then the volume of the parallelepiped having three coterminous edges as

$$\vec{\alpha} = \left(\vec{a} \cdot \vec{a}\right) \vec{a} + \left(\vec{a} \cdot \vec{b}\right) \vec{b} + \left(\vec{a} \cdot \vec{c}\right) \vec{c},$$
  

$$\vec{\beta} = \left(\vec{b} \cdot \vec{a}\right) \vec{a} + \left(\vec{b} \cdot \vec{b}\right) + \left(\vec{b} \cdot \vec{c}\right) \vec{c}$$
  
and 
$$\vec{\lambda} = \left(\vec{c} \cdot \vec{a}\right) \vec{a} + \left(\vec{c} \cdot \vec{b}\right) \vec{b} + \left(\vec{c} \cdot \vec{c}\right) \vec{c}$$
 is

A. 3V

B. 4V

 $\mathsf{C}.V^2$ 

D.  $V^3$ 

#### Answer: D

Watch Video Solution

**6.** A parallelepiped is formed by planes drawn parallel to coordinate axes through the points A=(1,2,3) and B=(9,8,5). The volume of that parallelepiped is equal to (in cubic units)

A. 192

B.48

C. 32

D. 96

Answer: D

Watch Video Solution

7. If  $\overrightarrow{a}, \overrightarrow{b}$  and  $\overrightarrow{c}$  are any three vectors forming a linearly independent system, then  $\forall \theta \in R$  $\overrightarrow{p} = \overrightarrow{a} \cos \theta + \overrightarrow{b} \sin \theta + \overrightarrow{c} (\cos 2\theta)$  $\overrightarrow{q} = \overrightarrow{a} \cos \left(\frac{2\pi}{3} + \theta\right) + \overrightarrow{b} \sin \left(\frac{2\pi}{3} + \theta\right) + \overrightarrow{c} (\cos 2) \left(\frac{2\pi}{3} + \theta\right)$ 

and

$$\overrightarrow{r} = \overrightarrow{a}\cos\left(\theta - \frac{2\pi}{3}
ight) + \overrightarrow{b}\sin\left(\theta - \frac{2\pi}{3}
ight) + \overrightarrow{c}\cos 2\left(\theta - \frac{2\pi}{3}
ight)$$
  
then  $\left[\overrightarrow{p}\overrightarrow{q}\overrightarrow{r}
ight]$ 

A. 
$$\begin{bmatrix} \overrightarrow{a} & \overrightarrow{b} & \overrightarrow{c} \end{bmatrix} \cos \theta$$
  
B.  $\begin{bmatrix} \overrightarrow{a} & \overrightarrow{b} & \overrightarrow{c} \end{bmatrix} \cos 2\theta$   
C.  $\begin{bmatrix} \overrightarrow{a} & \overrightarrow{b} & \overrightarrow{c} \end{bmatrix} \cos 3\theta$ 

D. None of these

#### Answer: D



8. Let 
$$\overrightarrow{r} = \left(\overrightarrow{a} \times \overrightarrow{b}\right) \sin x + \left(\overrightarrow{b} \times \overrightarrow{c}\right) \cos y + \left(\overrightarrow{c} \times \overrightarrow{a}\right)$$
,  
where  $\overrightarrow{a}, \overrightarrow{b}$  and  $\overrightarrow{c}$  are non-zero non-coplanar vectors, If  $\overrightarrow{r}$  is  
orthogonal to  $3\overrightarrow{a} + 5\overrightarrow{b} + 2\overrightarrow{c}$ , then the value of  
 $\sec^2 y + \csc^2 x + \sec y \csc x$  is

### A. 3

B. 4

C. 5

D. 6

Answer: A

Watch Video Solution

**9.** In a regular tetrahedron, let  $\boldsymbol{\theta}$  be angle betweenn any edge

and a face not containing the edge. The value of  $\cos^2 heta$  is

A. 1/6

B. 1/9

 $\mathsf{C.}\,1/3$ 

D. None of these

Answer: C



**10.** DABC be a tetrahedron such that AD is perpendicular to the base ABC and  $\angle ABC = 30^\circ$ . The volume of tetrahedron is 18. if value of AB + BC + AD is minimum, then the length of AC is

A.  $6\sqrt{2-\sqrt{3}}$ B.  $3(\sqrt{6}-\sqrt{2})$ C.  $6\sqrt{2+\sqrt{3}}$ D.  $3(\sqrt{6}+\sqrt{2})$ 

Watch Video Solution

#### Answer: A

11. If  $lpha \left( \overrightarrow{a} \times \overrightarrow{b} \right) + eta \left( \overrightarrow{b} \times \overrightarrow{c} \right) + \lambda \left( \overrightarrow{c} \times \overrightarrow{a} \right) = 0$ , then

A.  $\overrightarrow{a}, \overrightarrow{b}, \overrightarrow{c}$  are coplanar is all  $lpha, eta, \lambda 
eq 0$ 

B.  $\overrightarrow{a}, \overrightarrow{b}, \rightarrow$  are coplanar if any one of  $lpha, eta, \lambda \neq 0$ 

C.  $\overrightarrow{a}, \overrightarrow{b}, \overrightarrow{c}$  are non-coplanar for any  $lpha, eta, \lambda 
eq 0$ 

D. None of these

Answer: A::B

**Watch Video Solution**