



MATHS

BOOKS - DEEPTI MATHS (TELUGU ENGLISH)

APPLICATIONS OF DIFFERENTIATION

SOLVED EXAMPLES

1. The apporoximate value of $\frac{1}{\sqrt[4]{16.16}}$ is

A. 0.1999

B. 4.0008

C. 0.49875

D. 0.4983

Answer: C



Watch Video Solution

2. A circular plate expands when heated from a radius of 5 cm to 5.06 cm. The percentage increase in area is

A. 2.4

B. 0.72

C. 0.4

D. 0.6

Answer: A



Watch Video Solution

3. If the radius of a sphere is raised from 10 cm to 10.02 cm when heated, then the approximate change in the volume is

A. 8π cubic cm

B. 80π cubic cm

C. 0.06π cubic cm

D. 16π cubic cm

Answer: A



[Watch Video Solution](#)

4. The equation of the normal to the curve $y = x^3 - 2x^2 + 4$ at $x = 2$ is

A. $x + 4y = 18$

B. $x - 4y = 18$

C. $x + 4y + 18 = 0$

D. $x - 4y + 18 = 0$

Answer: A



[Watch Video Solution](#)

5. The point on the curve $y = x^3 + 5$, the tangent at which is parallel to the line $12x - y = 7$ is

A. $(1,0),(-1,-4)$

B. $(0,-1),(-2,3)$

C. $(2,13),(-2,-3)$

D. $(1,2),(1,-2)$

Answer: C



Watch Video Solution

6. The equation of the normal to the curve $y = 2x^3 + 6x^2 - 9$ where the curve crosses the y-axis is

A. $x = 9$

B. $x = 1$

C. $y + 9 = 0$

D. $y - 9 = 0$

Answer: C



Watch Video Solution

7. If θ is the angle between the curves $y = x^2$, $x = y^2$ at $(1,1)$,
then $\tan \theta =$

A. 3

B. $3/4$

C. $3/5$

D. $5/14$

Answer: B



[Watch Video Solution](#)

8. The length of the subnormal to the curve $y = x^3$ at $(1,1)$ is

A. 3

B. 16

C. 24

D. 8

Answer: A



Watch Video Solution

9. A stone projected vertically upward moves according to the law $s = 100t - 16t^2$. The maximum height reached is

A. $520/3$ unit

B. $625/4$ unit

C. 653 unit

D. 560 unit

Answer: B



Watch Video Solution

10. A particle moves according to law $s = t^3 - 3t^2 + 3t + 12$.

The velocity when the acceleration is zero is

A. $10\text{unit}/\text{sec}$

B. 0

C. $8/3 \text{ unit}/\text{sec}$

D. $= 3\text{unit}/\text{sec}$

Answer: B



Watch Video Solution

11. A person of height 2mt starts from a lamp post of height 5 mt and walks away at the constant rate of 6 km per hour. The rate at which his shadow increases is

- A. 2 kmph
- B. 6.4 kmph
- C. $8/3$ kmph
- D. 4 kmph

Answer: D



[Watch Video Solution](#)

12. The function $f(x) = 9x^2 - 15x - x^3 + 10$ is increasing in

- A. $(2/2, \infty)$

B. (1,5)

C. $(\frac{2}{3}, 5)$

D. $(\frac{2}{3}, \frac{3}{2})$

Answer: B



[Watch Video Solution](#)

13. The function $f(x) = 10 - 3x^3 + x$ is decreasing in

A. $(-\frac{5}{4}, \infty)$

B. $(\frac{2}{3}, \frac{3}{2})$

C. $(-\frac{1}{3}, \frac{1}{3})$

D. $\mathbb{R} - [-\frac{1}{3}, \frac{1}{3}]$

Answer: D



[Watch Video Solution](#)

14. The stationary point of $3x^4 - 4x^3 + 1$ is

- A. (1,0)
- B. (1,29)
- C. (1,1)
- D. none

Answer: A



[Watch Video Solution](#)

15. The minimum value of $x^3 - 6x^2 + 9x + 1$ is

A. 1

B. 2

C. 8

D. 4

Answer: A



Watch Video Solution

16. Maximum point of $f(x) = \cos ec x$ in $(-\pi, 0)$ is

A. $x = -\pi/2$

B. $x = -\pi/3$

C. $x = -\pi/4$

D. none

Answer: A



Watch Video Solution

17. If the line $ax + by + c = 0$ is a normal to the curve $xy = 1$ then

A. $a > 0, b < 0$

B. $a > 0, b > 0$

C. $a > b = 0$

D. $a < 0, b < 0$

Answer: A



Watch Video Solution

18. A particle is moving on a straight line so that its distance s from a fixed point at any time t is proportional to t^n . If v be the velocity and 'a' the acceleration at any time, then $\frac{nas}{n-1}$ equals

A. v

B. v^2

C. v^3

D. $2v$

Answer: B



[Watch Video Solution](#)

19. A body whose mass is 3kgs, performs rectilinear motion according to the formula $s = 1 + t + t^2$ (s is in cm, and t is in seconds). Then the kinetic energy of the body when $t = 5$ is

- A. $181 \cdot 15$ ergs
- B. $181 \cdot 5 \times 10^2$ ergs
- C. $181 \cdot 5 \times 10^3$ ergs
- D. $181 \cdot 5 \times 10^4$ ergs

Answer: C



[Watch Video Solution](#)

20. If $-4 \leq x \leq 4$ then the critical points of $f(x) = x^2 - 6(|x|) + 4$ are

A. 3,-2

B. 6,-6

C. 0,1

D. 3,-3,0

Answer: D



[Watch Video Solution](#)

EXERCISE 1A (APPROXIMATIONS AND ERRORS)

1. If $f(x) = 2x^2 + 3x - 5$, $x = 3$, $\delta x = 0.02$, then $\delta f =$

A. 0.3008

B. 0.3

C. 0.308

D. 0.8

Answer: A



[Watch Video Solution](#)

2. If $f(x) = 1/x$, $x = 1$, $\delta x = 0.02$, then $\delta f =$

A. 0.02

B. -0.02

C. 0.0196

D. -0.0196

Answer: D



[Watch Video Solution](#)

3. If $f(x) = \log x$, $x = 2$, $\delta x = 0.02$, then $df =$

A. $\log(1.01)$

B. 0.01

C. $= -\text{LOG}(1.01)$

D. -0.01

Answer: B



[Watch Video Solution](#)

4. If $f(x) = 1/x$, $x = 2$, $\delta x = 0.2$, then $df =$

A. 0.02

B. -0.02

C. `0.05

D. -0.05

Answer: D



[Watch Video Solution](#)

5. If $f(x) = 1/x^2$, $x = 2$, $\delta = -0.01$, then $df =$

A. 0.002519

B. 0.002915

C. 0.0025

D. 0.0019

Answer: C



Watch Video Solution

6. The approximate change in y , when

$$y = x^2 + 2x, x = 3, \delta = 0.01 \text{ is}$$

A. 3.6

B. 2

C. 0.08

D. 0.3

Answer: C



Watch Video Solution

7. The approximate change in y , when

$$y = 1/x^2, x = 1, \delta x = -0.01 \text{ is}$$

- A. 0.02
- B. -0.02
- C. 0.05
- D. -0.05

Answer: A



[Watch Video Solution](#)

8. The approximate value of $(2.001)^4$ is

- A. 27.54

B. 16.032

C. 2.9907

D. 5.0133

Answer: B



Watch Video Solution

9. The approximate value of $(3 \cdot 02)^5$ is

A. 128.75

B. 16.32

C. 251.1

D. 210.38

Answer: C



[Watch Video Solution](#)

10. The approximate value of $(1.0002)^{3000}$ is

A. 1.2

B. 1.4

C. 1.6

D. 1.8

Answer: C



[Watch Video Solution](#)

11. The value of $(127)^{1/3}$ to 4 decimal places is

A. 5.0267

B. 5.4267

C. 5.5267

D. 5.0001

Answer: A



Watch Video Solution

12. The approximate value of $\sqrt[4]{80}$ is

A. 27.54

B. 16.032

C. 2.9907

D. 5.0133

Answer: C



View Text Solution

13. The approximate value of $\frac{1}{\sqrt{25 \cdot 25}}$ is

A. 0.1999

B. 4.0008

C. 0.49875

D. 0.4983

Answer: A



View Text Solution

14. The approximate value of $\frac{1}{\sqrt[3]{8 \cdot 08}}$ is

A. 0.1999

B. 4.0008

C. 0.49875

D. 0.4983

Answer: D



[Watch Video Solution](#)

15. The approximate value of $\log(2.01)$, given that $\log 2 = 0.6934$ is

A. 0.6984

B. 0.49974

C. 1.6834

D. 1.6683

Answer: A



Watch Video Solution

16. The approximate value of $\sin 30^\circ 1'$, given that $1^\circ = 0.01745$ radian is

A. 1.00832

B. 0.50025

C. 0.00362

D. 1.00058

Answer: B



[View Text Solution](#)

17. The approximate value of $\cos 60^\circ$, given that $1^\circ = 0.01745$ radian is

A. 1.0349

B. 0.7193

C. 0.4849

D. 1.00058

Answer: C



[View Text Solution](#)

18. If $1^\circ = \alpha$ radians, then find the approximate value of $\cos 60^\circ 1'$.

A. $\frac{1}{2} + \frac{\alpha\sqrt{3}}{120}$

B. $\frac{1}{2} - \frac{\alpha}{120}$

C. $\frac{1}{2} - \frac{\alpha\sqrt{3}}{120}$

D. $\frac{1}{2} + \frac{\alpha}{120}$

Answer: C



[Watch Video Solution](#)

19. The approximate value of $\tan 45^\circ 1'$, given that $1^\circ = 0.01745$ radian is

A. 1.0349

B. 0.7193

C. 0.4849

D. 1.00058

Answer: D



[View Text Solution](#)

20. How many times of relative error in l is the relative error in

T , when $T = 2\pi\sqrt{l/g}$

A. 2

B. 3

C. 4

D. 5

Answer: A



Watch Video Solution

21. If an error of 0.02 cm is made while measuring the radius 10 cm of a circle, then the error in the area is

A. 0.02π sq.cm

B. 4.4π sq.cm

C. 0.4π sq.cm

D. 0.6π sq.cm

Answer: C



Watch Video Solution

22. IF an error of 0.01 cm is made while measuring the radius 2 cm of a circle, then the relative error in the circumference is

- A. 0.004
- B. 0.4
- C. 0.005
- D. 0.5

Answer: C



Watch Video Solution

23. IF an error of 0.01 cm is made while measuring the radius 5 cm of a circle, then the percentage error in the circumference is

A. 0.2

B. 0.02

C. 0.002

D. 0.0002

Answer: A



Watch Video Solution

24. If there is an error of 0.02 sq.cm is made in the area of a circle while measuring the radius 5 cm, then the percentage error in the circumference of the circle is

A. $\frac{0.4}{\pi}$

B. $\frac{0.04}{\pi}$

C. $\frac{4}{\pi}$

D. $\frac{0.004}{\pi}$

Answer: B



Watch Video Solution

25. A circular plate expands when heated from a radius of 5 cm to 5.06 cm. The approximate increase in area is

A. 2.4π sq.cm

B. 0.72π sq.cm

C. 0.4π sq.cm

D. 0.6π sq.cm

Answer: D



[Watch Video Solution](#)

26. The diameter x of a circle is found by measurement to be 5 cm with maximum error is 0.05 cm. The approximate maximum error in the area is

A. 0.125π sq.cm

B. 0.72π sq.cm

C. 0.05π sq.cm

D. 0.04π sq.cm

Answer: A



[Watch Video Solution](#)

27. The diameter x of a circle is found by measurement to be 5 cm with maximum error of 0.05 cm. The relative error in the area is

- A. 2
- B. 0.02
- C. 0.002
- D. 0.0002

Answer: B



[Watch Video Solution](#)

28. The circumference of a circle is measured as 56 cm with error 0.02 cm. The percentage error in its area is

A. $1/7$

B. $1/28$

C. $1/14$

D. $1/56$

Answer: C



Watch Video Solution

29. In measuring the circumference of a circle , there is an error of 0.05 cm. If with this error the circumference of the circle is measured in c cm, then the error in area is

A. $\frac{0.025c}{\pi}$ sq.cm

B. $\frac{0.25c}{\pi}$ sq.cm

C. $\frac{0.0025c}{\pi}$ sq.cm

D. none

Answer: A



Watch Video Solution

30. In measuring the circumference of a circle , there is an error of 0.05 cm. If with this error the circumference of the circle is measured in c cm, then the percentage error in area is

A. $\frac{0.1}{c}$

B. $\frac{0.01}{c}$

C. $\frac{0.001}{c}$

D. $\frac{10}{c}$

Answer: D



Watch Video Solution

31. The circumference of a circle is measured as 14 cm with an error of 0.01 cm. The approximate percentage error in the area of the circle is

A. $3/10$

B. $1/8$

C. $1/7$

D. $3/2$

Answer: C



Watch Video Solution

32. In measuring the area of a circle in 25π sq.cm, there is an error of 0.02π sq.cm. The percentage error in its circumference is

A. 0.04

B. 0.02

C. 0.01

D. 0.05

Answer: A



[Watch Video Solution](#)

33. In measuring the vertical angle of the sector of a circle of radius 30 cms, an error of 1° is made. The error in the area of

the sector is

A. 2.5π sq.cms.

B. 25π sq.cms.

C. 3π sq.cms.

D. 30π sq.cms.

Answer: A



[View Text Solution](#)

34. If an error of 0.02 cm is made while measuring the radius 10 cm of a sphere, then the error in the volume is

A. 8π cubic cm

B. 80π cubic cm

C. 0.06π cubic cm

D. 16π cubic cm

Answer: A



Watch Video Solution

35. If there is a possible error of 0.02 cm in the measurement of the diameter of a sphere then the possible percentage error in its volume when the radius 10 cm is

A. 0.1

B. 0.2

C. 0.3

D. 0.4

Answer: C



[View Text Solution](#)

36. There is an error of ± 0.04 cm in the measurement of the diameter of a sphere. When the radius is 10 cm, the percentage error in the volume of the sphere is :

A. ± 1.2

B. ± 1.0

C. ± 0.8

D. ± 0.6

Answer: D



[Watch Video Solution](#)

37. If there is an error of $\frac{1}{10}\%$ in the measurement of the radius of a sphere, then the percentage error in the calculation of the volume of the sphere is

A. $\frac{3}{10}$

B. $\frac{1}{8}$

C. $\frac{1}{7}$

D. $\frac{3}{2}$

Answer: A



[View Text Solution](#)

38. If the radius of a sphere is raised from 10 cm to 10.02 cm when heated, then the percentage increase in volume is

A. 0.2

B. 0.6

C. 0.05

D. $\pi / 10$

Answer: B



[View Text Solution](#)

39. The radius of a sphere is 3 cm. If an error of 0.03 cm is made in measuring the radius of the sphere, then the percentage error in surface area is

A. 0.125π sq.cm

B. 0.72π sq.cm

C. $.0.05\pi$ sq.cm

D. 0.04π sq.cm

Answer: B



[Watch Video Solution](#)

40. The radius of a sphere is 3 cm. If an error of 0.03 cm is made in measuring the radius of the sphere, then the percentage error in surface area is

A. 2

B. 0.02

C. 0.002

D. 0.0002

Answer: A



Watch Video Solution

41. If there is an error of 0.01 cm in the diameter of a sphere when its radius 5 cm, then the percentage error in its surface area is

A. 0.2

B. 0.6

C. 0.05

D. $\pi / 10$

Answer: A



View Text Solution

42. If there is a possible error of 0.01 cm in the measurement of side of a cube, the possible error in its surface area when the side is 10 cm is

A. 1.2 sq.cm

B. 1.4 sq.cm

C. 2.4 sq.cm

D. 3.6 sq.cm

Answer: A



[View Text Solution](#)

43. If there is an error of 0.02 cm in the measurement of the side as 10 cm of a cube, then error in the surface area is

A. 1.2 sq.cm

B. 1.4 sq.cm

C. 2.4 sq.cm

D. 3.6 sq.cm

Answer: C



[View Text Solution](#)

44. If there is an error of 0.05 cm is made while measuring the side 10 cm of a cube, then the error in the volume is

A. 10 cubic cm

B. 12 cubic cm

C. 15 cubic cm

D. 20 cubic cm

Answer: C

 [View Text Solution](#)

45. If there is an error of 0.05 cm in the measurement of the side as 2 cm of a cube, then relative error in the volume is

A. 0.075

B. 0.0075

C. 7.5

D. 0.75

Answer: A

 [View Text Solution](#)

46. The percentage error in measuring the side of a cube is 0.5, Then the percentage error in its volume is

A. $1/2$

B. 1

C. $3/2$

D. 2

Answer: C



[View Text Solution](#)

47. In a cube the percentage increase in the side is 1. The percentage increase in volume of cube is

A. 2

B. $1/2$

C. $1/3$

D. 3

Answer: D



Watch Video Solution

48. The approximate percentage reduction in the volume of a cube of ice if each side of ice cube is reduced by 0.7 percentage due to melting is

A. 2.1

B. 2.5

C. 3.2

D. 3.3

Answer: A



[View Text Solution](#)

49. If the side of a cube is 10.01 cm, the approximate volume of the cube is

A. 103 cubic cm

B. 1003 cubic cm

C. 110 cubic cm

D. 1010 cubic cm

Answer: B



[Watch Video Solution](#)

50. If there is an error 0.01 cm in the measurement of the radius 10 cm of a cylinder of fixed height 20 cm then error in the volume is

- A. 4π cubic cm
- B. 2.5π cubic cm
- C. 0.06 cubic cm
- D. 0.6 cubic cm

Answer: A



[Watch Video Solution](#)

51. If there is an error 0.01 cm in the measurement of the radius 10 cm of a cylinder of fixed height 20 cm then percentage error in the volume is

- A. 0.2
- B. 0.02
- C. 0.002
- D. 0.0002

Answer: A



[Watch Video Solution](#)

52. The radius and height of a cylinder are measured as 5 cm and 10 cm respectively and there is an error of 0.02 cm in the

both measurements. The approximate error in the volume is

- A. 4π cubic cm
- B. 2.5π cubic cm
- C. 0.06π cubic cm
- D. 0.6π cubic cm

Answer: B



[View Text Solution](#)

53. The radius and height of a cylinder are measured as 5 cm and 10 cm respectively and there is an error of 0.02 cm in both the measurements. The percentage error in volume is

- A. 1

B. 0.01

C. 0.001

D. -1

Answer: A



[Watch Video Solution](#)

54. The radius and height of a cone are measured as 6 cm and 12 cm respectively and there is an error of 0.06 cm in both the measurements. The approximate error in the volume of the cone is

A. $410 \pi \sqrt{2}$ sq.cm

B. 60π sq.cm

C. 3.6π sq.cm

D. $320 \pi \sqrt{2}$ sq.cm

Answer: C

 [Watch Video Solution](#)

55. If errors of 1% each are made in the base radius and height of a cylinder, then the percentage error in its volume, is

A. 1

B. 2

C. 3

D. none

Answer: C

 [Watch Video Solution](#)

56. The semi-vertical angle of a cone is 45° . If the height of the cone is 20.025, then its approximate lateral surface area, is

A. $410 \pi \sqrt{2}$ sq.cm

B. 60π sq.cm

C. 3.6π sq.cm

D. 9045π sq.cm

Answer: A



Watch Video Solution

57. The semi vertical angle of a cone is 45° . The height of the cone is 30.05 cm. The approximately its volume is

A. $410 \pi \sqrt{2}$ sq.cm

B. 60π sq.cm

C. 3.6π sq.cm

D. 9045π sq.cm

Answer: D



[Watch Video Solution](#)

58. If there is an error of 0.05 cm, while measuring the side of an equilateral triangle as 5 cm, then the percnetage error in area is

A. $\frac{2}{3}$

B. $\frac{1}{3}$

C. 2

D. 1

Answer: C



Watch Video Solution

59. The angle A of $\triangle ABC$ is found by measurement to be 63° and the area is calculated by the formula $\frac{1}{2}bc \sin A$. The percentage error in the calculated value of the area due to an error of 15 minutes in the measured value of A is

A. $\frac{5\pi}{54} \cot 47^\circ$

B. $\frac{\pi bc}{1440} \cos 63^\circ$

C. $\frac{5\pi}{36} \cot 63^\circ$

D. $\frac{\pi bc}{1120} \cos 42^\circ$

Answer: C



View Text Solution

60. In $\triangle ABC$ the sides b, c are given. If there is an error δA in increasing angle A then $\delta =$

A. $\frac{\Delta}{2a} \cdot \delta A$

B. $\frac{2\Delta}{a} \delta A$

C. $bc \sin A \cdot \delta A$

D. none

Answer: B



[View Text Solution](#)

61. If the area of $\triangle ABC$ is calculated from the measurements of b , c , A and k is the error in A , then the percentage error in area is

- A. $95 k \cot A$
- B. $100 k \cot A$
- C. $110 k \cot A$
- D. $111 k \cot A$

Answer: B



[Watch Video Solution](#)

62. If in a triangle the side a and the angle A remain constant, while other elements are changed slightly then

A. $\delta b \sin B + \delta c \sin C = 0$

B. $\delta b \cos B + \delta c \cos C = 0$

C. $\delta b \sec B + \delta c \sec C = 0$

D. $\delta b \cos cB + \delta c \cos ecC = 0$

Answer: C



[View Text Solution](#)

63. If the length of a simple pendulum is decreased by 3%, then the percentage error in its period T is

A. 2

B. 2.5

C. 1.8

D. 1.5

Answer: D



Watch Video Solution

64. If there is an error of 2% in measuring its length, of a simple pendulum then the percentage error in the period will be

A. 1

B. -1

C. 2

D. 4

Answer: A

 [Watch Video Solution](#)

65. Given $PV=C$ (constant). The percentage of increase in V corresponding to an increase 1% in the value of P is

A. 0

B. 1

C. -1

D. 2

Answer: C

 [View Text Solution](#)

66. The pressure P and the volume v of a gas are connected by the relation $pv^{1/4} = a$ constant. The percentage increase in the pressure corresponding to a diminution of $\frac{1}{2}\%$ in the volume is

A. $1/2$

B. $1/4$

C. 4

D. $1/8$

Answer: D



[Watch Video Solution](#)

67. The focal length of a mirror is given by $\frac{2}{f} = \frac{1}{v} = -\frac{1}{u}$. In

finding the values of u and v , the errors are equal and equal to

' p '. Then the relative error in f is

A. $\frac{p}{2} \left(\frac{1}{u} + \frac{1}{v} \right)$

B. $p \left(\frac{1}{u} + \frac{1}{v} \right)$

C. $\frac{p}{2} \left(\frac{1}{u} - \frac{1}{v} \right)$

D. $p \left(\frac{1}{u} - \frac{1}{v} \right)$

Answer: B



[View Text Solution](#)

EXERCISE 1B (TANGENT AND NORMAL)

1. The gradient of the curve $y = x^3 - 3x^2 - 2x + 7$ at (1,3) is

A. 3

B. -4

C. -5

D. 7

Answer: C



[Watch Video Solution](#)

2. The slope of the tangent to the curve $y = 6 + x - x^2$ at (2,4) is

A. 10

B. $1/2$

C. -3

D. $1/\sqrt{2}$

Answer: C



Watch Video Solution

3. The slope of the tangent to the curve $y = \sin x$ at $x = \pi/4$ is

A. 0

B. 1

C. -1

D. $1/\sqrt{2}$

Answer: D



Watch Video Solution

4. The slope of the tangent to the curve $x^2 + y^2 = 4$ at $(\sqrt{2}, \sqrt{2})$ is

A. 0

B. 1

C. -1

D. $1/\sqrt{2}$

Answer: C



Watch Video Solution

5. The slope of the tangent to the curve $x = at^3$, $y = at^4$ at $t=1$ is

A. $-1/2$

B. $-2/3$

C. $1/4$

D. $4/3$

Answer: D



Watch Video Solution

6. Slope of the normal to the curve $y = \cos 2x$ "at" $x = \pi/6$ is

A. $-1/2$

B. $1/\sqrt{3}$

C. $3/4$

D. $4/3$

Answer: B



[Watch Video Solution](#)

7. Slope of the normal curve $xy = 12$ at $(3, 4)$ is

A. $-1/2$

B. $1/\sqrt{3}$

C. $3/4$

D. $4/3$

Answer: C



[Watch Video Solution](#)

8. Slope of the normal to the curve $x^3 + y^3 = 6xy$ at $(3, 3)$ is

A. 0

B. 1

C. -1

D. $1/\sqrt{2}$

Answer: B



[Watch Video Solution](#)

9. The slope of the normal to the curve

$x = a(\theta - \sin \theta)$, $y = a(1 - \cos \theta)$ at $\theta = \pi/2$ is

A. 0

B. 1

C. -1

D. $1/\sqrt{2}$

Answer: C



Watch Video Solution

10. Slope of the normal to the curve

$x = a(t + \sin t), y = a(t - \sin t)$ is

A. $\frac{\cot^2 t}{2}$

B. $-\frac{\cot^2 t}{2}$

C. $\frac{\tan^2 t}{2}$

D. $-\frac{\tan^2 t}{2}$

Answer: B

 [Watch Video Solution](#)

11. If normal of the curve is parallel to x axis then

A. $\frac{dy}{dx} = 0$

B. $\frac{dy}{dx} = 1$

C. $\frac{dx}{dy} = 0$

D. $\frac{dx}{dy} = 1$

Answer: C

 [Watch Video Solution](#)

12. If the slope of the tangent to the curve $xy + ax = by$ at (1,1) is 2, then (a,b)=

A. (0, 1)

B. (1, 2)

C. (- 1, 2)

D. (1, - 2)

Answer: B



[Watch Video Solution](#)

13. The equation of the tangent to the curve $y = x^3 + 1$ at (1,2) is

A. $3x - y = 1$

B. $x - 3y + 1 = 0$

C. $3x + y = 1$

D. $x + 3y + 1 = 0$

Answer: A



[Watch Video Solution](#)

14. The equation of the tangent to the curve $6y = 7 - x^3$ at $(1,1)$ is

A. $2x + y = 3$

B. $x + 2y = 3$

C. $x + y = 1$

D. $x + y + 2 = 0$

Answer: B



Watch Video Solution

15. The equation of the tangent to the curve $y(x + 1) = 4$ at the point $(2, 4/3)$ is

A. $4x + 9y + 20 = 0$

B. $4x + 9y - 20 = 0$

C. $9x + 4y + 20 = 0$

D. $9x - 4y + 20 = 0$

Answer: B



Watch Video Solution

16. The equation of the tangent to the curve

$$2x^2 - xy + 3y^2 = 18 \text{ at } (3, 1) \text{ is}$$

A. $11x + 3y - 36 = 0$

B. $11x - 3y + 36 = 0$

C. $3x + 11y - 2 = 0$

D. $3x - 11y + 2 = 0$

Answer: A



Watch Video Solution

17. The equation of the tangent to the curve $y = x^3 - 2x + 7$

at $(1, 6)$ is

A. $y = x + 5$

B. $x + y = 7$

C. $2x + y = 8$

D. $x + 2y = 13$

Answer: A



Watch Video Solution

18. The equation of the tangent to the curve $y = x^3 - 2x + 7$ at (x_1, y_1) is

A. $\frac{xx_1}{a} + \frac{yy_1}{b} = 1$

B. $\frac{xx_1}{a} - \frac{yy_1}{b} = 1$

C. $\frac{xx_1}{a^2} + \frac{yy_1}{b^2} = 1$

D. $\frac{xx_1}{a^2} - \frac{yy_1}{b^2} = 1$

Answer: C



View Text Solution

19. The equation of the tangent to the curve $y = \frac{6x}{x^2 - 1}$ at $(2, 4)$ is

A. $10x + 3y - 32 = 0$

B. $10x - 3y + 32 = 0$

C. $3x + 10y - 34 = 0$

D. $3x - 10y + 34 = 0$

Answer: A



Watch Video Solution

20. The equation of the tangent to the curve $y = \frac{8}{4 + x^2}$ at $x = 2$ is

A. $2x + y + 3 = 0$

B. $x - 2y + 4 = 0$

C. $2x - y - 3 = 0$

D. $x + 2y - 4 = 0$

Answer: D



Watch Video Solution

21. Equation of the tangent line at $x = a$ to the curve $y = a \log \sec \frac{x}{a}$ is

A. $(y - a \log \sec a) \tan 1 = x - a$

B. $(x - a)\tan 1 = y - a \log \sec 1$

C. $(x - a)\cos 1 = (y - a \log \sec 1)\sin$

D. none

Answer: B



Watch Video Solution

22. Equation of the tangent line at $y = a/4$ to the curve $y(x^2 + a^2) = ax^2$ is

A. $8y = -3\sqrt{3x} + a$

B. $8y = 3\sqrt{3x} + a$

C. $8y = 3\sqrt{3x} - a$

D. none

Answer: C



Watch Video Solution

23. Equation of the tangent to the curve $y^2 = 4ax$ at $(at^2, 2at)$ is

A. $x + yt - at^2 = 0$

B. $xt - y = 2at + at^3$

C. $xt + y = 2at + at^3$

D. $x - yt + at^2 = 0$

Answer: D



Watch Video Solution

24. The equation of the tangent to the curve $y^2 = \frac{x^3}{2a - x}$ at (a, a) is

A. $y + 2x = a$

B. $2x - y = a$

C. $x + 2y = 3a$

D. $2y - x = 3a$

Answer: B



Watch Video Solution

25. The equation of the tangent to the curve $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$ at $(a \sec \theta, b \tan \theta)$ is

A. $\frac{ax}{\sec \theta} + \frac{by}{\tan \theta} = a^2 + b^2$

$$\text{B. } \frac{ax}{\cos \theta} - \frac{by}{\sin \theta} = a^2 - b^2$$

$$\text{C. } \frac{x}{y} \cos \theta + \frac{y}{b} \sin \theta = 1$$

$$\text{D. } \frac{x}{y} \tan \theta - \frac{y}{b} \sec \theta = 1$$

Answer: D



View Text Solution

26. The equation of the tangent to the curve

$$\left(\frac{x}{a}\right)^4 + \left(\frac{y}{b}\right)^4 = 2 \text{ at } (a, b) \text{ is}$$

$$\text{A. } \frac{x}{a} + \frac{y}{b} = 2$$

$$\text{B. } \frac{x}{a} - \frac{y}{b} = 2$$

$$\text{C. } ax + by = a^2 + b^2$$

$$\text{D. } ax - by = a^2 - b^2$$

Answer: A



Watch Video Solution

27. The tangent line at $\left(\frac{a}{\sqrt{8}}, \frac{a}{\sqrt{8}}\right)$ to the curve

$x^{2/3} + y^{2/3} = a^{2/3}$ is parallel to the line

A. $x = -y$

B. $x = y$

C. $x = 0$

D. $y = 0$

Answer: A



View Text Solution

28. The equation of the tangent to the curve

$$\left(\frac{x}{a}\right)^{2/3} + \left(\frac{y}{b}\right)^{2/3} = 1 \text{ at } (a \cos^3 \theta, b \sin^3 \theta) \text{ is}$$

A. $ax \cos \theta + by \sin \theta = a^2 \cos^4 \theta + b^2 \sin^4 \theta$

B. $ax \cos \theta - by \sin \theta = a^2 \cos^4 \theta - b^2 \sin^4 \theta$

C. $\frac{x}{a \cos \theta} + \frac{y}{b \sin \theta} = 1$

D. $\frac{x}{a \cos \theta} - \frac{y}{b \sin \theta} = 1$

Answer: C



Watch Video Solution

29. The equation of the normal to the curve

$$x = a(\theta - \sin \theta), y = a(1 - \cos \theta) \text{ at } (\theta = \pi/2) \text{ is}$$

A. $2x - 2y + 4a - \pi a = 0$

B. $2x - 2y - \pi a = 0$

C. $2x + 2y + \pi a = 0$

D. $2x + 2a - 4a - \pi a = 0$

Answer: A



Watch Video Solution

30. Equation of the tangent at $\theta = \pi/4$ to the curve

$x = a \cos 2\theta, y = 2\sqrt{2}a \sin \theta$ is

A. $x + y = a$

B. $x + y = 2a$

C. $y = x + 2a$

D. none

Answer: B



[View Text Solution](#)

31. The equation of the tangent to the curve $\sqrt{x} + \sqrt{y} = 5$ at (9,4) is

A. $2x - 3y + 30 = 0$

B. $3x - 2y - 19 = 0$

C. $2x + 3y - 30 = 0$

D. $3x + 2y + 19 = 0$

Answer: C



[Watch Video Solution](#)

32. The equation of the normal to the curve $y = 3x^2 + 4x - 6$ at $(1, 1)$ is

A. $x + 10y - 11 = 0$

B. $x - 10y + 12 = 0$

C. $x + y - 9 = 0$

D. none

Answer: A



[Watch Video Solution](#)

33. The equation of the normal to the curve $y = x + \frac{1}{x}$ at $(1, 2)$ is

A. $x = 1$

B. $x = 2$

C. $x = 3$

D. $x = 4$

Answer: A



Watch Video Solution

34. The equation of the normal to the curve $2y = 3 - x^2$ at $(1, 1)$ is

A. $x + y = 0$

B. $x - y = 0$

C. $x + y = 2$

D. $x - y = 2$

Answer: B



Watch Video Solution

35. The equation of the normal to the curve $x^2 = 4y$ at $(2, 1)$ is

A. $x + y + 3 = 0$

B. $x + y - 3 = 0$

C. $x - y + 3 = 0$

D. $x - y - 3 = 0$

Answer: B



Watch Video Solution

36. The equation of the normal to the curve $y^2 = x^3$ at $x = 8$ is

A. $x \pm 3\sqrt{2}y = 104$

B. $x \pm 2\sqrt{3}y = 104$

C. $x \pm 5\sqrt{2}y = 104$

D. $x \pm 2\sqrt{5}y = 104$

Answer: A



Watch Video Solution

37. The equation of the normal to the curve $y^4 = ax^3$ at (a, a) is

A. $x + 2y = 3a$

B. $3x - 4y + a = 0$

C. $4x + 3y = 7a$

D. $4x - 3y = 0$

Answer: C



Watch Video Solution

38. The equation of the normal to the curve $3y^2 = 4x + 1$ at (1,2) is

A. $3x + y + 5 = 0$

B. $3x + y - 5 = 0$

C. $3x - y + 5 = 0$

D. $3x - y - 5 = 0$

Answer: B



Watch Video Solution

39. the equation of the normal to the curve $y = \frac{6x}{x^2 - 1}$ at $(2, 4)$ is

A. $10x + 3y - 32 = 0$

B. $10x - 3y + 32 = 0$

C. $3x + 10y - 34 = 0$

D. $3x - 10y + 34 = 0$

Answer: D



Watch Video Solution

40. The equation of the normal to the curve $y = \frac{8}{4 + x^2}$ at $x = 2$ is

A. $2x + y + 3 = 0$

B. $x - 2y + 4 = 0$

C. $2x - y - 3 = 0$

D. $x + 2y - 4 = 0$

Answer: C



Watch Video Solution

41. The equation of the normal to the curve $y^2 = 4ax$ at $(at^2, 2at)$ is

A. $x + yt - at^2 = 0$

B. $xt - y = 2at + at^3$

C. $xt + y = 2at + at^3$

D. $x - yt + at^2 = 0$

Answer: C



Watch Video Solution

42. The equation of the normal to the curve $2x^2 - xy + 3y^2 = 18$ at $(3, 1)$ is

A. $11x + 3y - 36 = 0$

B. $11x - 3y + 36 = 0$

C. $3x + 11y - 2 = 0$

D. $3x - 11y + 2 = 0$

Answer: D



Watch Video Solution

43. The equation of the normal to the curve $y^2 = \frac{x^3}{2a - x}$ at (a, a) is

A. $y + 2x = a$

B. $2x - y = a$

C. $x + 2y = 3a$

D. $2y - x = 3a$

Answer: C



Watch Video Solution

44. The equation of the normal to the curve $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$ at $(a \cos \theta, b \sin \theta)$ is

A. $\frac{ax}{\sec \theta} + \frac{by}{\tan \theta} = a^2 + b^2$

B. $\frac{ax}{\cos \theta} - \frac{by}{\sin \theta} = a^2 - b^2$

C. $\frac{x}{a} \cos \theta + \frac{y}{b} \sin \theta = 1$

D. $\frac{x}{a} \sec \theta - \frac{y}{b} \tan \theta = 1$

Answer: B



Watch Video Solution

45. The equation of the normal to the curve $\left(\frac{x}{a}\right)^4 + \left(\frac{y}{b}\right)^4 = 2$ at (a, b) is

A. $\frac{x}{a} + \frac{y}{b} = 2$

$$B. \frac{x}{a} - \frac{y}{b} = 2$$

$$C. ax + by = a^2 + b^2$$

$$D. ax - by = a^2 - b^2$$

Answer: D



Watch Video Solution

46. The equation of the normal to the curve

$$\left(\frac{x}{a}\right)^{2/3} + \left(\frac{y}{b}\right)^{2/3} = 1 \text{ at } (a \cos^3 \theta, b \sin^3 \theta) \text{ is}$$

$$A. ax \cos \theta + by \sin \theta = a^2 \cos^4 \theta + b^2 \sin^4 \theta$$

$$B. ax \cos \theta - by \sin \theta = a^2 \cos^4 \theta - b^2 \sin^4 \theta$$

$$C. \frac{x}{a \cos \theta} + \frac{y}{b \sin \theta} = 1$$

$$D. \frac{x}{a \cos \theta} - \frac{y}{b \sin \theta} = 1$$

Answer: B



Watch Video Solution

47. The equation of the normal to the curve $x = a(\theta - \sin \theta)$, $y = a(1 - \cos \theta)$ at $(\theta = \pi/2)$ is

A. $2x - 2y + 4a - \pi a = 0$

B. $2x + 2y - \pi a = 0$

C. $2x + 2y + \pi a = 0$

D. $2x + 2y - 4a - \pi a = 0$

Answer: B



Watch Video Solution

48. The normal to the curve $x = a(1 + \cos \theta)$, $y = a \sin \theta$ at ' θ ' always passes through the fixed point

A. $(a, 0)$

B. (a, a)

C. $(0, 0)$

D. $(0, a)$

Answer: A



Watch Video Solution

49. The normal to the curve $x = a(\cos \theta + \theta \sin \theta)$, $y = a(\sin \theta - \theta \cos \theta)$ at any point ' θ ' is such that

- A. it passes through the origin
- B. it makes angle $\pi/2 + \theta$ with the x-axis
- C. it passes through $(a\pi/2, -a)$
- D. it is at a constant distance from the origin

Answer: D



Watch Video Solution

50. If the equation of the tangent at $(2,3)$ on the curve

$y = ax^2 + b$ is $y = 4x - 5$ then $a =$

- A. 0
- B. 1
- C. -1

D. 2

Answer: B

 [Watch Video Solution](#)

51. If $y = 4x - 5$ is a tangent to the curve $y^2 = px^3 + q$ at $(2, 3)$, then

A. $p = 2, q = -7$

B. $p = -2, q = 7$

C. $p = -2, q = -7$

D. $p = 2, q = 7$

 [Watch Video Solution](#)

52. The angle made by the tangent at any point of the curve

$x = a(t + \sin t \cos t)$, $y = a(1 + \sin t)^2$ with x-axis is

A. $\frac{\pi}{2}$

B. $\frac{\pi}{4}$

C. $\frac{\pi + t}{2}$

D. $\frac{\pi + 2t}{4}$

Answer: D



[View Text Solution](#)

53. The angle which the tangent at (2,1) to the curve

$x^3 + y^3 = 9$ with the x-axis is

A. $\tan^{-1} 4$

B. $\tan^{-1} 2$

C. $\pi/2$

D. $\pi/4$

Answer: A



Watch Video Solution

54. If the tangent to the curve $xy + ax + by = 0$ at $(1,1)$ is inclined at an angle $\tan^{-1} 2$ with x-axis, then

A. $a = 1, b = 2$

B. $a = 1, b = -2$

C. $a = -1, b = 2$

D. $a = -1, b = -2$

Answer: B



Watch Video Solution

55. If the line $ax + by + c = 0$ is a normal to the curve $xy = 1$ then

A. $a > 0, b > 0$

B. $a > 0, b < 0$

C. $a < 0, b < 0$

D. none

Answer: B



Watch Video Solution

56. The coordinates of the point P on the curve $x = a(\theta + \sin \theta)$, $y = a(1 - \cos \theta)$ where the tangent is inclined at angle $\frac{\pi}{4}$ to the x-axis, are

A. $\left(a\left(\frac{\pi}{2} - 1\right), a\right)$

B. $\left(a\left(\frac{\pi}{2} + 1\right), a\right)$

C. $\left(a\frac{\pi}{2}, a\right)$

D. (a, a)

Answer: B



Watch Video Solution

57. Points at which the tangents to the hyperbola $y = \frac{x - 1}{x + 1}$ are parallel to the line $y = 2x + 1$ are

A. $(0, 1), (2, -3)$

B. $(0, -1), (-2, 3)$

C. $(0, 2), (3, -5)$

D. none

Answer: A



Watch Video Solution

58. The point on the curve $y = x^2 + 5$, the tangent at which is perpendicular to the line $x + 2y = 2$ is

A. $(1, 6)$

B. $(1, -6)$

C. $(-1, 6)$

D. $(-1, -6)$

Answer: A

 [Watch Video Solution](#)

59. The curve $y - e^{xy} + x = 0$ has a vertical tangent at the point

A. $(1, 1)$

B. at no point

C. $(0, 1)$

D. $(1, 0)$

Answer: A

 [Watch Video Solution](#)

60. The point on the curve $y = x^3 + x - 2$, the tangent at which is parallel to the line $y = 4x - 1$ is

- A. $(1, 0), (-1, -4)$
- B. $(0, -1), (-2, 3)$
- C. $(2, 13), (-2, -3)$
- D. $(1, 2), (1, -2)$

Answer: A



[Watch Video Solution](#)

61. The line $x/a + y/b = 1$ is a tangent to curve $y = be^{-x/a}$ at the point

A. $(0, 0)$

B. $(0, a)$

C. $(0, b)$

D. $(b, 0)$

Answer: A



[View Text Solution](#)

62. The points on the curve $y = x^3$, the tangent at which are inclined at an angle of 60° to x-axis are

A. $\left(3^{-1/4}, 3^{-3/4}\right), \left(-3^{-1/4}, -3^{-3/4}\right)$

B. $\left(3^{-1/2}, 3^{-2/5}\right), \left(-3^{1/3}, -3^{-2/3}\right)$

C. $\left(2^{1/4}, 2^{-2/5}\right), \left(-3^{1/2}, -3^{-1/2}\right)$

D. none

Answer: D

 [Watch Video Solution](#)

63. The point on the curve $x^2 + y^2 - 2x - 3 = 0$ at which the tangent is parallel to x-axis is

A. $(1, 0), (-1, -4)$

B. $(0, -1), (-2, 3)$

C. $(2, 13), (-2, -3)$

D. $(1, 2), (1, -2)$

Answer: D

 [Watch Video Solution](#)

64. The point on the curve $x^4 - 4x^3 + 4x^2 + 1$ at which the tangent is parallel to x-axis is

- A. $(0, 1), (1, 2), (2, 1)$
- B. $(0 - 1), (-1, 2), (2, 1)$
- C. $(0, 1), (1, -2), (2, -1)$
- D. $(0, -1), (-1, 2), (-2, 1)$

Answer: A



[Watch Video Solution](#)

65. The points on the curve $y = x^2 + \sqrt{1 - x^2}$ at which the tangent is perpendicular to x-axis are

A. $(1, 1)$ only

B. $(\pm 1, 1)$

C. $(1, \pm 1)$

D. $(-1, 1)$ only

Answer: B



[Watch Video Solution](#)

66. The points on the curve $2a^2y = -x^3 - 3ax^2$ at which the tangent is perpendicular to y-axis is

A. $(0, 0), (2a, 2a)$

B. $(0, 0), (-2a, 2a)$

C. $(0, 0), (2a, -2a)$

D. $(0, 0), (-2a, -2a)$

Answer: D



Watch Video Solution

67. The equation of the tangent to the curve $4x^2 + 9y^2 = 40$ and having slope $-2/9$ is

A. $9x - 8y \pm 26 = 0$

B. $2x + 9y \pm 20 = 0$

C. $2x + 3y \pm 26 = 0$

D. $3x + y + 6 = 0$

Answer: B



Watch Video Solution

68. The equation of the tangent to the curve $9x^2 + 16y^2 = 52$ and which is parallel to the line $9x - 8y = 1$ is

A. $9x - 8y \pm 26 = 0$

B. $2x + 9y \pm 20 = 0$

C. $2x + 3y \pm 26 = 0$

D. $3x + y + 6 = 0$

Answer: C



[Watch Video Solution](#)

69. The equation of the tangent to the curve $x^2 + y^2 = 52$ and which is parallel to $2x + 3y = 6$ is

A. $9x - 8y \pm 26 = 0$

B. $2x + 9y \pm 20 = 0$

C. $2x + 3y \pm 26 = 0$

D. $3x + y + 6 = 0$

Answer: C



Watch Video Solution

70. The equation of the tangent to the curve $y = x + \frac{4}{x^2}$, that is parallel to the x-axis, is

A. $y = 0$

B. $y = 1$

C. $y = 2$

D. $y = 3$

Answer: D



Watch Video Solution

71. The equation of the tangent to the curve $y = x^3 + 3x^2 - 5$ and which is perpendicular to $2x - 6y + 1 = 0$ is

A. $9x - 8y \pm 26 = 0$

B. $2x + 9y \pm 20 = 0$

C. $2x + 3y \pm 26 = 0$

D. $3x + y + 6 = 0$

Answer: D



Watch Video Solution

72. The equation of the tangent to the curve $y^2 = 4x + 5$ and which is parallel to $y = 2x + 7$ is

A. $y = x + 2$

B. $x = y + 3$

C. $2x = y + 3$

D. $y = 2x + 3$

Answer: D



Watch Video Solution

73. The equation of the tangent to the curve $x^2 + 2y = 8$ and which is perpendicular to $x - 2y + 1 = 0$ is

A. $2x + y + 6 = 0$

B. $2x + y - 6 = 0$

C. $2x - y + 6 = 0$

D. $2x - y - 6 = 0$

Answer: B



[Watch Video Solution](#)

74. The points of contact of the tangents drawn from origin to the curve $3y = 3 + x^2$ is

A. $(3, 2)$

B. $(3, \sqrt{2})$

C. $(\pm \sqrt{3}, 2)$

D. $(3, \pm \sqrt{2})$

Answer: C



Watch Video Solution

75. The point of contact of the tangents drawn from origin to the curve $y = x^2 + 3x + 4$ is

A. $(2, 14), (2, 2)$

B. $(-2, 14), (2, 2)$

C. $(2, 14), (-2, 2)$

D. $(2, -14), (-2, 2)$

Answer: C



Watch Video Solution

76. The equation of the tangent to the curve $y(x - 2)(x - 3) - x + 7 = 0$ where the curve cuts x-axis is

A. $20x + y - 140 = 0$

B. $x - 20y - 7 = 0$

C. $20x - y + 140 = 0$

D. $x + 20y + 7 = 0$

Answer: B



Watch Video Solution

77. The equation of the tangent to the curve $y = be^{-x/a}$ where it crosses the y-axis is

A. $ax + by = 1$

B. $\frac{x}{a} + \frac{y}{b} = 1$

C. $\frac{x}{b} + \frac{y}{a} = 1$

D. $ax - by = 1$

Answer: B



Watch Video Solution

78. Equation of the tangent to the curve $y = 2x^3 - 6x^2 - 9$ at the point where the curve crosses the y-axis is

A. $y + 9 = 0$

B. $y - 9 = 0$

C. $2y + 1 = 0$

D. $2y - 1 = 0$

Answer: A



Watch Video Solution

79. The equation of the tangent to the curve $y = \frac{x + 9}{x + 5}$ so

that is passes through the origin is

A. $x + y = 0$

B. $x - y = 0$

C. $x + y = 1$

D. $x - y = 1$

Answer: A



Watch Video Solution

80. The equation of the normal to the curve $y = be^{-x/a}$ where it cuts y-axis is

A. $\frac{x}{a} + \frac{y}{b} = 1$

B. $\frac{x}{a} - \frac{y}{b} = 1$

C. $ax + by - b^2 = 0$

D. $ax - by + b^2 = 0$

Answer: D



Watch Video Solution

81. The equation of the normal to the curve

$y = 2x^3 + 6x^2 - 9$ where the curve crosses the y-axis is

A. $x = 0$

B. $x = 1$

C. $x = 2$

D. $x = 3$

Answer: A



Watch Video Solution

82. The distance between the origin and the normal to the

curve $y = e^{2x} + x^2$ at $x = 0$ is

A. 2

B. $\sqrt{5}$

C. $2\sqrt{5}$

D. $2/\sqrt{5}$

Answer: D



Watch Video Solution

83. The normal to the curve $x = a(\cos \theta + \theta \sin \theta)$, $y = a(\sin \theta - \theta \cos \theta)$ at any point ' θ ' is such that

A. it makes constant angle with positive x-axis

B. it passes through (0, 0)

C. it is at a constant distance from (0, 0)

D. none

Answer: C

 [Watch Video Solution](#)

84. If the normal to the curve $x^3 - y^2 = 0$ at $(m^2, -m^3)$ is $y = mx - 2m^3$, then the value of m^2 is

A. $1/3$

B. $1/6$

C. $2/3$

D. $2/3$

Answer: C

 [Watch Video Solution](#)

85. The portion of the tangent drawn at any point on $x^{2/3} + y^{2/3} = a^{2/3}$ ($a > 0$), except the points on the coordinate axes, included between the the coordinates axes is

A. a

B. $2a$

C. $a^{2/3}$

D. a^2

Answer: A



Watch Video Solution

86. The tangent at $\theta = \pi/4$ to the curve $x = a \cos^3 \theta, y = a \sin^3 \theta$ meets the x and y axis in A and B , then the length of AB is

A. a

B. $2a$

C. a^2

D. $a/2$

Answer: A



[Watch Video Solution](#)

87. Tangent at any point of the curve $\left(\frac{x}{a}\right)^{2/3} + \left(\frac{y}{b}\right)^{2/3} = 1$ makes intercepts x_1 and y_1 on the axes. Then

A. $\left(\frac{x_1}{a}\right)^{2/3} + \left(\frac{y_1}{b}\right)^{2/3} = 1$

B. $\frac{x_1^2}{a^2} + \frac{y_1^2}{b^2} = 1$

C. $\frac{x_1^3}{a^3} + \frac{y_1^3}{b^3} = 1$

D. none

Answer: B



Watch Video Solution

88. The sum of the intercepts on the coordinate axes of any tangent to $\sqrt{x} + \sqrt{y} = \sqrt{a}$ is

A. a

B. $3a$

C. $5a$

D. $9a$

Answer: A



Watch Video Solution

89. IF the tangent at any point P on the curve $x^m y^n = a^{m+n}$, $mn \neq 0$ meets the coordinate axes in A,B then show that $AP : BP$ is a constant.

A. $m : n$

B. $n : m$

C. $-m : n$

D. $-n : m$

Answer: B

 [Watch Video Solution](#)

90. The tangent at any point of the curve $x = at^3, y = at^4$ divides the abscissa of the point of contact in the ratio

A. 2:3

B. 3:2

C. 1:3

D. 3:1

Answer: C

 [Watch Video Solution](#)

91. If p and q are the lengths of the perpendiculars from the origin on the tangent and the normal to the curve $x^{2/3} + y^{2/3} = a^{2/3}$, then $4p^2 + q^2 =$

A. a

B. a^2

C. $2a^2$

D. $5a^2$

Answer: B



[Watch Video Solution](#)

92. If the tangent at any point on the curve $x^4 + y^4 = a^4$ cuts off intercepts p and q on the coordinate axes, then

$$p^{-4/3} + q^{-4/3} =$$

A. $a^{-4/3}$

B. $a^{-1/2}$

C. $a^{1/2}$

D. a

Answer: A



[Watch Video Solution](#)

93. The sum of the squares of the intercepts on the coordinate axes of any tangent to $x^{2/3} + y^{2/3} = a^{2/3}$ is

A. a^2

B. $aa^2/2$

C. $2a^2$

D. $3a^2$

Answer: A



Watch Video Solution

94. If the tangent at any point on the curve $x^{1/3} + y^{1/3} = a^{1/3}$ ($a > 0$) cuts off intercepts p and q on the coordinate axes then $\sqrt{p} + \sqrt{q} =$

A. \sqrt{a}

B. $\sqrt[3]{a}$

C. $2\sqrt{a}$

D. $2\sqrt[3]{a}$

Answer: A



[Watch Video Solution](#)

95. If the sum of the squares of the intercepts on the axes cut off by the tangent to the curve $x^{1/3} + y^{1/3} = a^{1/3}$ with $(a > 0)$ at $(a/8, a/8)$ is 2, then a has the value

A. 1

B. 2

C. 4

D. 8

Answer: C



[Watch Video Solution](#)

96. If the tangent at the point (at^2, at^3) on the curve $ay^2 = x^3$ meets the curve again at Q , then $q =$

A. $\left(\frac{at^2}{4}, \frac{-at^3}{8}\right)$

B. $\left(\frac{at}{4}, 8at\right)$

C. $\left(\frac{at}{2}, 2at^2\right)$

D. $\left(\frac{at}{2}, at^2\right)$

Answer: A



Watch Video Solution

97. If the tangent at $(1, 1)$ on $y^2 = x(2 - x)^2$ meets the curve again at P , then P is

A. (4, 4)

B. (- 1, 2)

C. (9/4, 3/8)

D. none

Answer: C



Watch Video Solution

98. If the tangent at pto the curve $xy = c^2$ meets the axes at A, B and pdivides AB in the ratio

A. 1:2

B. 1:1

C. 2:5

D. 3:5

Answer: B



Watch Video Solution

99. The area of the triangle formed by the tangent to the curve $xy = a^2$ at point on the curve, with the coordinate axes is

A. a^2

B. $2a^2$

C. $4a^2$

D. $8a^2$

Answer: B

 [Watch Video Solution](#)

100. The area of the triangle formed by the tangent to the curve $y = 8 / (4 + x^2)$ at $x = 2$ and the co-ordinates axes is

A. 2 sq.units

B. 4 sq.units

C. 8 sq.units

D. $7/2$ sq.units

Answer: B

 [Watch Video Solution](#)

101. If δ is the area of the triangle formed by the positive x axis and the normal and tangent to the circle $x^2 + y^2 = 4$ at $(1, \sqrt{3})$, then $\delta =$

A. $\frac{\sqrt{3}}{2}$

B. $\sqrt{3}$

C. $2\sqrt{3}$

D. 6

Answer: C



[Watch Video Solution](#)

102. If the tangent at $\theta = \pi/4$ to the curve $x = a \cos^3 \theta, y = a \sin^3 \theta$ meets the x and y axis in A and B,

then the area of δAOB is

A. a^2

B. $a^2 / 2$

C. $a^2 / 4$

D. $a^2 / 8$

Answer: C



[Watch Video Solution](#)

103. If the area of the triangle, included between the axes and any tangent to the curve $xy^n = a^{n+1}$ is constant, then the value of n is

A. -1

B. -2

C. 1

D. 2

Answer: C



[Watch Video Solution](#)

104. The area of the triangle formed by the normal to the curve $x = e^{\sin y}$ at $(1, 0)$ with the coordinate axes is

A. $1/4$

B. $1/2$

C. $3/4$

D. 1

Answer: B



Watch Video Solution

105. The area of the triangle formed by the tangent and the normal at the points (a, a) on the curve $y^2 = \frac{x^3}{2a - x}$ and the line $x = 2a$ is

A. $a^2/4$

B. $a^2/2$

C. $5a^2/4$

D. $9a^2/4$

Answer: C



Watch Video Solution

106. Area of the triangle formed by the tangent, normal to the curve $x^2/a^2 + y^2/b^2 = 1$ at the point $(a/\sqrt{2}, b/\sqrt{2})$ and the x axis is

A. $\frac{ab}{4} \sqrt{a^2 + b^2}$

B. $4ab$

C. $\frac{b}{4a} (a^2 + b^2)$

D. none

Answer: C



Watch Video Solution

107. Area of the triangle formed by the tangent, normal at $(1, 1)$ on the curve $\sqrt{x} + \sqrt{y} = 2$ and the x axis is

A. 1 sq.units

B. 2 sq.units $1/2$ sq.units

C. $1/2$ sq.units

D. 4 sq.units

Answer: A



[Watch Video Solution](#)

108. Area of the triangle formed by the tangent, normal at (a, a) on $y(2a - x) = x^2$ and the x axis is

A. $a^2/3$

B. $5a^2$

C. $5a^2/3$

D. none

Answer: C



Watch Video Solution

109. The point on the intersection of the tangents drawn to the curve $x^2y = 1 - y$ at the points where it is intersected by the curve $x^2y = 1 - y$ at the points where it is intersected by the curve $xy = 1 - y$ is

A. (0, 1)

B. (1, 1/2)

C. (0, -1)

D. (1/2, 1)

Answer: A



Watch Video Solution

110. The two curves $y = x^2 + 1$, $y = 3x^2 - 4x + 3$ at $(1, 2)$

- A. touch each other
- B. cut orthogonally
- C. cut at an angle of 45°
- D. none

Answer: A



Watch Video Solution

111. The two curves $y = x^{-3}$, $y = e^{3(1-x)}$ at (1,1)

- A. touch each other
- B. cut orthogonally
- C. cut at an angle of 45°
- D. none

Answer: A



[Watch Video Solution](#)

112. The two curves $y^2 = 4(x + 1)$, $y^2 = 36(9 - x)$ at (8, 6)

- A. touch each other
- B. cut orthogonally

C. cut at an angle of 45°

D. none

Answer: B



[Watch Video Solution](#)

113. The two curves $y = \frac{x + 3}{x^2 + 1}$, $y = \frac{x^2 - 7x + 11}{x - 1}$ at $(2, 1)$

A. touch each other

B. cut orthogonally

C. cut at an angle of 45°

D. none

Answer: C



[Watch Video Solution](#)

114. The two curves $2x^2 + y^2 = 20$, $x^2 - 4y^2 + 8 = 0$

- A. touch each other
- B. cut orthogonally
- C. cut at an angle of 45°
- D. none

Answer: B



Watch Video Solution

115. The two curves $x^2 + y^2 = 25$, $2x^2 - 9y + 18 = 0$

- A. touch each other

B. cut orthogonally

C. cut at an angle of 45°

D. none

Answer: B



[Watch Video Solution](#)

116. The curves $y = x^3 - 3x^2 - 8x - 4$, $y = 3x^2 + 7x + 4$ touch at the point $(-1, 0)$. The equation of the common tangent is

A. $x + y + 1 = 0$

B. $x + y - 1 = 0$

C. $x - y + 1 = 0$

D. $x - y - 1 = 0$

Answer: C



Watch Video Solution

117. The curves $y = x^2 - 1$, $y = 8x - x^2 - 9$ touch each other at the point (2, 3). The equation of the common normal is

A. $4x + y + 5 = 0$

B. $4x - y - 5 = 0$

C. $x + 4y - 14 = 0$

D. $x - 4y + 14 = 0$

Answer: C



Watch Video Solution

118. The angle between the curves $y^2 = 4x$ and $x^2 = 2y - 3$ at the point $(1, 2)$ is

A. 30°

B. 60°

C. 90°

D. 0°

Answer: D



Watch Video Solution

119. The angle between the curves $y^2 = 8x$, $x^2 = 4y - 12$ at $(2, 4)$ is

A. $\pi/2$

B. $\pi/4$

C. $\pi/6$

D. 0

Answer: D



[Watch Video Solution](#)

120. The angle between the curves $xy = 4$ and $x^2 - y^2 = 15$ at the point $(-4, -1)$ is

A. 60°

B. 90°

C. $\tan^{-1}(1/2)$

D. $\tan^{-1}(5/2)$

Answer: B



[Watch Video Solution](#)

121. The angle between the curves $y^2 = 4x + 4$ and $y^2 = 36(9 - x)$ is

A. 30°

B. 45°

C. 60°

D. 90°

Answer: D



Watch Video Solution

122. The angle between curves $y^2 = 4ax$, $ay = 2x^2$ is

A. $\tan^{-1}\left(\frac{3}{4}\right)$

B. $\tan^{-1}\left(\frac{3}{5}\right)$

C. $\tan^{-1}\left(\frac{4}{3}\right)$

D. $\tan^{-1}\left(\frac{5}{3}\right)$

Answer: B



Watch Video Solution

123. The angle between the curves $y^2 = x$, $x^2 = y$ at $(1, 1)$ is

A. $\frac{\pi}{2}$

B. $\frac{\pi}{4}$

C. $\tan^{-1} \frac{3}{4}$

D. $\tan^{-1} \frac{4}{3}$

Answer: C



[Watch Video Solution](#)

124. The curves $y = 3x^2$, $y^2 = 2x$ intersect at origin

A. orthogonally

B. at an angle of $\pi/4$

C. at an angle of $\pi / 3$

D. at an angle of $\pi / 6$

Answer: A



Watch Video Solution

125. The angle between the curves $y^2 = 4x$, $x^2 = 4y$ at $(0, 0)$ is

A. $\pi / 2$

B. $\pi / 3$

C. $\pi / 6$

D. 0

Answer: a



Watch Video Solution

126. The curves $y = x^3$, $6y = 7 - x^2$ intersect at $(1, 1)$ at an angle of

A. $\pi/2$

B. $\pi/4$

C. $\pi/3$

D. none

Answer: A



Watch Video Solution

127. The angle between the curves $y = x$, $y = 1/x$ at $(1, 1)$ is

A. $\pi/2$

B. $\pi/3$

C. $\pi/6$

D. 0

Answer: A



Watch Video Solution

128. The angle between the curves $x^2 = 2y$, $x^2 + y^2 = 8$ at (2, 2) is

A. $\tan^{-1}\left(\frac{1}{3}\right)$

B. $\tan^{-1}(3)$

C. $\tan^{-1}\left(\frac{1}{2}\right)$

D. $\tan^{-1}\left(\frac{2}{3}\right)$

Answer: B

 [Watch Video Solution](#)

129. The angle between the curves $x^2 = 4y$, $x^2 + y^2 = 5$ at $(-2, 1)$ is

A. 30°

B. 60°

C. $\tan^{-1}3$

D. none

Answer: B

 [Watch Video Solution](#)

130. The angle between the curves $x^2 = 4y$, $y^2 = 4x$ at $(4, 4)$ is

A. $\tan^{-1}\left(\frac{1}{2}\right)$

B. $\tan^{-1}\left(\frac{3}{4}\right)$

C. $\frac{\pi}{2}$

D. $\frac{\pi}{4}$

Answer: B



[Watch Video Solution](#)

131. The angle between the curves $x^2 = 4y$, $x^2 + y^2 = 5$ at $(-2, 1)$ is

A. 3

B. $3/4$

C. $3/5$

D. $5/14$

Answer: A



[Watch Video Solution](#)

132. If θ is the angle between the curves $y^2 = 4ax$, $ay = 2x^2$ at $(a, 2a)$, then $\tan \theta =$

A. 3

B. $3/4$

C. $3/5$

D. $5/14$

Answer: C

 [Watch Video Solution](#)

133. If θ is the angle between the curves $xy = 2$, $y^2 = 4x$ at $(1, 2)$, then $\tan \theta =$

A. 3

B. $3/4$

C. $3/5$

D. $5/14$

Answer: A

 [Watch Video Solution](#)

134. Find the angle between the curves $xy=2$ and $x^2 + 4y = 0$

A. 1

B. -1

C. 2

D. 3

Answer: D



Watch Video Solution

135. The angle between the curves $y = \sin x$ and $y = \cos x$ is

A. 2

B. $\sqrt{2}$

C. $1\sqrt{2}$

D. $2\sqrt{2}$

Answer: D



Watch Video Solution

136. The angle between the curves $y = \sin x$ and $y = \cos x$ is

A. $\tan^{-1}(2\sqrt{2})$

B. $\tan^{-1}(3\sqrt{2})$

C. $\tan^{-1}(3\sqrt{3})$

D. $\tan^{-1}(5\sqrt{2})$

Answer: A



Watch Video Solution

137. If θ is the angle of intersection of the curves $y^2 = x^3$ and $y = 2x^2 - 1$ at $(1, 1)$, then $|\tan \theta| =$

A. $5/14$

B. $5/12$

C. $25/12$

D. none

Answer: A



Watch Video Solution

138. The angle between the curves $xy = 2$ and $y^2 = 4x$ is

A. $\tan^{-1}\left(\frac{1}{3}\right)$

B. $\tan^{-1}3$

C. $\tan^{-1}\left(\frac{1}{2}\right)$

D. $\tan^{-1}\left(\frac{2}{3}\right)$

Answer: B



Watch Video Solution

139. The angle between the curves $x^2 + y^2 = 4$ and $x^2 = 3y$ is

A. $\tan^{-1}\left(\frac{6}{13}\right)$

B. $\tan^{-1}\left(\frac{3}{4}\right)$

C. $\tan^{-1}\left(\frac{5}{\sqrt{3}}\right)$

D. $\tan^{-1}\left(4\frac{\sqrt{2}}{7}\right)$

Answer: C



Watch Video Solution

140. The angle between the curves $y^2 = 4x$ and $x^2 = 4y$ is

A. $\tan^{-1}\left(\frac{6}{13}\right)$

B. $\tan^{-1}\left(\frac{3}{4}\right), \frac{\pi}{2}$

C. $\tan^{-1}\left(\frac{5}{\sqrt{3}}\right)$

D. $\tan^{-1}\left(4\frac{\sqrt{2}}{7}\right)$

Answer: B



Watch Video Solution

141. The angle between the curves $y = x^2$ and $y = 4 - x^2$ is

A. $\tan^{-1}\left(\frac{6}{13}\right)$

B. $\tan^{-1}\left(\frac{3}{4}\right)$

C. $\tan^{-1}\left(\frac{5}{\sqrt{3}}\right)$

D. $\tan^{-1}\left(4\frac{\sqrt{2}}{7}\right)$

Answer: D



Watch Video Solution

142. The angle between the curves $y^2 = 8x$, $x^2 = 4y - 12$ at $(2, 4)$ is

A. $\pi/2$

B. $\pi/4$

C. $\pi/6$

D. 0

Answer: D



[Watch Video Solution](#)

143. The condition that the two curves $x = y^2$, $xy = k$ cut orthogonally is

A. $2k^2 = 1$

B. $8k^2 = 1$

C. $8k^3 = 1$

D. $2k^3 = 1$

Answer: B



[Watch Video Solution](#)

144. The two curves $x = y^2$, $xy = a^3$ cut orthogonally at a point, then $a^2 =$

A. $1/3$

B. $1/2$

C. 2

D. 3

Answer: B



[Watch Video Solution](#)

145. The condition that the two curves $y^2 = 4ax$, $xy = c^2$ cut orthogonally is

A. $c^2 = 16a^2$

B. $c^2 = 32a^2$

C. $c^4 = 16a^4$

D. $c^4 = 32a^4$

Answer: D



Watch Video Solution

146. The curves $ax^2 + by^2 = 1$ and $Ax^2 + By^2 = 1$ intersect orthogonally, then

A. $\frac{1}{a} + \frac{1}{A} = \frac{1}{b} + \frac{1}{B}$

$$\text{B. } \frac{1}{a} - \frac{1}{A} = \frac{1}{b} - \frac{1}{B}$$

$$\text{C. } \frac{1}{a} + \frac{1}{A} = \frac{1}{b} - \frac{1}{B}$$

D. none

Answer: B



Watch Video Solution

147. If the curves $x^2 + py^2 = 1$ and $qx^2 + y^2 = 1$ are orthogonal to each other, then

$$\text{A. } p - q = 2$$

$$\text{B. } \frac{1}{p} - \frac{1}{q} = 2$$

$$\text{C. } \frac{1}{p} + \frac{1}{q} = -2$$

$$\text{D. } \frac{1}{p} + \frac{1}{q} = 2$$

Answer: D



Watch Video Solution

148. If the curves $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$ and $\frac{x^2}{25} + \frac{y^2}{16} = 1$ cut each other orthogonally, then $a^2 - b^2 =$

A. 400

B. 75

C. 41

D. 9

Answer: D



Watch Video Solution

149. If the curves $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$ and $\frac{x^2}{l^2} - \frac{y^2}{m^2} = 1$ cut each other orthogonally, then

A. $a^2 + b^2 = l^2 + m^2$

B. $a^2 - b^2 = l^2 - m^2$

C. $a^2 - b^2 = l^2 + m^2$

D. $a^2 + b^2 = l^2 - m^2$

Answer: C



Watch Video Solution

150. The curves $\frac{x^2}{a^2 + k_1} + \frac{y^2}{b^2 + k_1} = 1$ and $\frac{x^2}{a^2 + k_2} + \frac{y^2}{b^2 + k_2} = 1$ where $k_1 \neq k_2$ intersect at an angle

A. 0

B. $\pi/4$

C. $\pi/3$

D. $\pi/2$

Answer: D



Watch Video Solution

151. If the curves $y^2 = 6x$, $9x^2 + by^2 = 16$, cut each other at right angles then the value of b is

A. 2

B. 4

C. $9/2$

D. none

Answer: C



Watch Video Solution

152. Angle between the tangents to the curve $y = x^2 - 5x + 6$ at the points $(2, 0)$ and $(3, 0)$ is

A. $\pi/6$

B. $\pi/4$

C. $\pi/3$

D. $\pi/2$

Answer: D



Watch Video Solution

153. The length of the tangent of the curve $y = x^3 + 1$ at $(1, 2)$ is

A. $\sqrt{10}$

B. $2\sqrt{10}$

C. $2\sqrt{10}/3$

D. 6

Answer: C



Watch Video Solution

154. The length of the tangent of the curve $y^2 = \frac{x^3}{2a - x}$ at (a, a) is

A. $\sqrt{5}|a|$

B. $2|a|$

C. $\sqrt{5}\frac{|a|}{2}$

D. $\frac{|a|}{2}$

Answer: C



Watch Video Solution

155. The length of the tangent of the curve

$2x^2 + 3xy - 2y^2 = 8$ at $(2, 3)$ is

A. $\frac{\sqrt{325}}{2}$

B. $3\frac{\sqrt{325}}{17}$

C. $\frac{17}{2}$

D. $\frac{18}{17}$

Answer: B



Watch Video Solution

156. The length of the tangent of the curve

$x = a \cos^3 \theta, y = a \sin^3 \theta (a > 0)$ is

A. $a \sin^2 \theta$

B. $a \sin^2 \theta |\tan \theta|$

C. $a \sin^2 \theta |\cos \theta|$

D. $a \sin^4 |\sec \theta|$

Answer: A



Watch Video Solution

157. The length of the normal to the curve $y = x^2 + 1$ at $(1, 2)$ is

- A. $\sqrt{5}$
- B. $2\sqrt{5}$
- C. 1
- D. 4

Answer: B



[Watch Video Solution](#)

158. The length of the normal to the curve $y = c \cos\left(\frac{hx}{c}\right)$ at any point is

- A. y/c

B. y^2 / c

C. $2y / c$

D. $2y^2 / c$

Answer: B



Watch Video Solution

159. The length of the normal to the curve $y^2 = \frac{x^3}{2a - x}$ at (a, a) is

A. $\sqrt{5}|a|$

B. $2|a|$

C. $\sqrt{5} \frac{|a|}{2}$

D. $\frac{|a|}{2}$

Answer: A



Watch Video Solution

160. The length of the normal of the curve $2x^2 + 3xy - 2y^2 = 8$ at $(2, 3)$ is

A. $\frac{\sqrt{325}}{2}$

B. $3\frac{\sqrt{325}}{17}$

C. $\frac{17}{2}$

D. $\frac{18}{17}$

Answer: A



Watch Video Solution

161. The length of the normal to the curves

$$x = a \cos^3 \theta, y = a \sin^3 \theta (a > 0) \text{ is}$$

A. $a \sin^2 \theta$

B. $a \sin^2 \theta |\tan \theta|$

C. $a \sin^2 \theta |\cos \theta|$

D. $a \sin^4 |\sec \theta|$

Answer: B



Watch Video Solution

162. The length of the normal at pon the curve

$$x = a(t + \sin t), y = a(1 - \cos t) \text{ is}$$

A. $a \sin t$

B. $2a \sin^3 \frac{t}{2} \sec \frac{t}{2}$

C. $2a \sin \frac{t}{2} \tan \frac{t}{2}$

D. $2a \sin \frac{t}{2}$

Answer: C



Watch Video Solution

163. The length of the normal to the curve

$y = a \left(\frac{e^{-x/a} + e^{x/a}}{2} \right)$ at any point varies as the

A. abscissa of the point

B. ordinate of the point

C. square of the avscissa of the point

D. square of the ordinate of the point

Answer: D



Watch Video Solution

164. The length of subtangent to $y = be^{x/a}$ at any point is

A. a

B. $2a$

C. a^2

D. $a/2$

Answer: A



Watch Video Solution

165. The length of the subtangent at any point (x_1, y_1) on the curve $y = 5^x$ is

A. 5^{x_1}

B. $y_1 \cdot 5^{x_1}$

C. $\log_e 5$

D. $\frac{1}{\log_e 5}$

Answer: D



[Watch Video Solution](#)

166. The length of subtangent to $\sqrt{x} + \sqrt{y} = 3$ at $(4, 1)$ is

A. 2

B. $\sqrt{2}$

C. $1/\sqrt{2}$

D. $2\sqrt{2}$

Answer: A



Watch Video Solution

167. The length of subtangent to $x^2y^2 = a^4$ at (a, a) is

A. a

B. $2a$

C. a^2

D. $a/2$

Answer: A



Watch Video Solution

168. The length of subtangent to $x^2 + xy + y^2 = 7$ at $(1, -3)$ is

A. 5

B. $1/5$

C. $3/5$

D. 15

Answer: D



Watch Video Solution

169. The length of subtangent to $y = x \sin x$ at $x = \pi/2$ is

A. $\pi/2$

B. $\pi/4$

C. $\pi/6$

D. 0

Answer: A



Watch Video Solution

170. The length of the subtangent to the curve $y^2 = \frac{x^2}{2a - x}$

at (a, a) is

A. $\sqrt{5}|a|$

B. $2|a|$

C. $\sqrt{5}\frac{|a|}{2}$

D. $\frac{|a|}{2}$

Answer: D

 [Watch Video Solution](#)

171. The length of the subtangent of the curve $2x^2 + 3xy - 2y^2 = 8$ at $(2, 3)$ is

A. $\frac{\sqrt{325}}{2}$

B. $3\frac{\sqrt{325}}{17}$

C. $\frac{17}{2}$

D. $\frac{18}{17}$

Answer: D

 [Watch Video Solution](#)

172. The length of subtangent to

$x = a(\theta + \sin \theta)$, $y = a(1 - \cos \theta)$ at θ is

A. $a|\sin \theta|$

B. $a|\cos \theta|$

C. $a|\tan \theta|$

D. $a|\cot \theta|$

Answer: A



[Watch Video Solution](#)

173. The length of the subtangent of the curve

$x = a \cos^3 \theta$, $y = a \sin^3 \theta$ ($a > 0$) is

A. $a \sin^2 \theta$

B. $a \sin^2 \theta |\tan \theta|$

C. $a \sin^2 \theta |\cos \theta|$

D. $a \sin^4 |\sec \theta|$

Answer: C



Watch Video Solution

174. For the parabola $y^2 = 4ax$, the ratio of the subtangent to the abscissa is

A. 1 : 1

B. 2 : 1

C. $x : y$

D. $x^2 : y$

Answer: B

 [Watch Video Solution](#)

175. The subtangent at $x = \pi/2$ on the curve $y = x \sin x$ is

A. 1

B. $\pi/2$

C. 0

D. π

Answer: B

 [Watch Video Solution](#)

176. The subtangent to the curve $x^m y^n = a^{m+n}$ at any point (x, y) is

A. $-\frac{mx}{y}$

B. $-\frac{ny}{m}$

C. $-\frac{mx}{n}$

D. $\frac{nx}{m}$

Answer: D



[Watch Video Solution](#)

177. The length of the subtangent (if exists) at any point θ on the ellipse $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$ is

A. $a|\sin \theta|\sec^2 \theta$

B. $a \sin \theta|\sec \theta|$

C. $a|\sin \theta \cos \theta|$

D. $a \sin^2 \theta|\sec \theta|$

Answer: D



Watch Video Solution

178. The length of the subtangent to the curve $x^2y^2 = a^4$ at any point $(-a, a)$ is

A. $3a$

B. $2a$

C. a

D. $4a$

Answer: C



[Watch Video Solution](#)

179. The length of the subtangent at $(2, 2)$ to the curve

$$x^5 = 2y^4 \text{ is}$$

A. $5/2$

B. $8/5$

C. $2/5$

D. $5/8$

Answer: B



[Watch Video Solution](#)

180. The curve $y^2 = (x + a)^3$, the square of the subtangent is

..... Subnormal

A. equal to

B. varies as

C. double the

D. square of the

Answer: B



[Watch Video Solution](#)

181. The length of subnormal at $(-1, 4)$ on $y = 4x^2$ is

A. 4

B. 16

C. 32

D. 8

Answer: C



Watch Video Solution

182. The length of the subnormal to the curve $y^2 = x^3$ at $(4, 8)$

is

A. 24

B. $8/3$

C. $3/8$

D. none

Answer: A



Watch Video Solution

183. The length of the subnormal to the curve $y^2 = 2px$ is

A. p

B. $2p$

C. $3p$

D. $4p$

Answer: A



Watch Video Solution

184. The length of the subnormal to the curve $x^2 = 4ay$ at $(4a, 4a)$ is

A. $2a$

B. $4a$

C. $6a$

D. $8a$

Answer: D



[Watch Video Solution](#)

185. The length of subnormal of the curve $2x^2 + 3xy - 2y^2 = 8$ at $(2, 3)$ is

A. $\frac{\sqrt{325}}{2}$

B. $3\frac{\sqrt{325}}{17}$

C. $\frac{17}{2}$

D. $\frac{18}{17}$

Answer: C



Watch Video Solution

186. The length of subnormal to the curve $y = b \sin \frac{x}{a}$ at any point is

A. $a \tan \frac{x}{a}$

B. $\frac{b}{2a} \sin \frac{x}{a}$

C. $\frac{b}{2a} \sin \frac{2x}{a}$

D. $\frac{b^2}{2a} \sin \frac{2x}{a}$

Answer: D



Watch Video Solution

187. The length of the subnormal of the curves

$x = a \cos^3 \theta$ and $y = a \sin^3 \theta$ ($a > 0$) is

A. $a \sin^2 \theta$

B. $a \sin^2 \theta |\tan \theta|$

C. $a \sin^2 \theta |\cos \theta|$

D. $a \sin^4 \theta |\sec \theta|$

Answer: D



Watch Video Solution

188. The length of subnormal of the curves

$y = \frac{a}{2} (e^{x/a} + e^{-x/a})$ at any point is

A. $a \cosh^2 \frac{x}{a}$

B. $\frac{a}{2} \sinh \frac{2x}{a}$

C. $a \cosh \frac{x}{a}$

D. $a \sinh \frac{2x}{a}$

Answer: B



Watch Video Solution

189. The subnormal of the curve $y = a^x$ at any point on the curve varies directly as

A. cube of the ordinate (y^3)

B. square of the ordinate (y^2)

C. ordinate (y)

D. none

Answer: B



[Watch Video Solution](#)

190. The subnormal to the curve $xy = c^2$ at any point varies directly as

A. cube of the ordinate (y^3)

B. square of the ordinate (y^2)

C. ordinate (y)

D. none

Answer: A



Watch Video Solution

191. The subtangent at any point of the curve $x^m y^n = a^{m+n}$ varies as its

- A. abscissa of the point
- B. ordinate of the point
- C. square of abscissa
- D. square of ordinate

Answer: A



Watch Video Solution

192. For the curve $y^2 = (x + a)^3$, the square of the subtangent is Subnormal

A. equal to

B. varies as

C. double the

D. square of the

Answer: B



[Watch Video Solution](#)

193. The subtangent, ordinate and subnormal to the parabola $y^2 = 4ax$ at a point (different from the origin) are in

A. A.P.

B. G.P.

C. H.P.

D. none

Answer: B



Watch Video Solution

194. If the relation between subnormal SN and subtangent ST at any point on the curve $by^2 = (x + a)^3$ is $p(SN) = q(ST)^2$, then $p/q =$

A. $8/27$

B. $27/8$

C. $8\frac{b}{27}$

D. $27b/8$

Answer: C



Watch Video Solution

195. If the subnormal of the curve $xy^n = a^{n+1}$ is constant, then the value of n is

A. 1

B. -1

C. 5

D. -2

Answer: D



Watch Video Solution

196. If the subnormal at any point $y = a^{1-n}x^n$ is of constant length, then the value of n is

A. 1

B. $1/2$

C. 2

D. -2

Answer: B



[Watch Video Solution](#)

197. If at any point on the curve $y = f(x)$, the length of the subnormal is constant, then the curve will be a

A. circle

B. ellipse

C. parabola

D. straight line

Answer: C



Watch Video Solution

198. The length of the subtangent to the curve $y = ae^{x/b}$ at any point is

A. constant

B. equal to subnormal

C. equal to the square of the subnormal

D. none

Answer: A



[Watch Video Solution](#)

199. the length of the subnormal to the curve $y^2 = 4ax$ at any point is

A. constant

B. equal to subnormal

C. equal to the square of the subnormal

D. none

Answer: A



[Watch Video Solution](#)

200. Length of the subtangent at any point on $y^n = a^{n-1}x$ is

- A. proportional to abscissa
- B. proportional to ordinate
- C. length of the subnormal
- D. none

Answer: B



Watch Video Solution

201. Length of the subnormal at any point on $y^n = a^{n-1}x$ is constant when $n =$

A. 1

B. -1

C. -2

D. 2

Answer: D



Watch Video Solution

202. The length of the tangent to the curve

$$x = a\left(\cos t + \log \tan \frac{t}{2}\right), y = a \sin t \text{ at any point is}$$

A. constant

B. equal to subnormal

C. equal to the square of the subnormal

D. none

Answer: A

 [Watch Video Solution](#)

203. The sum of the lengths of the subtangent and subnormal at $\theta = \pi/3$ on the cycloid $x = a(\theta - \sin \theta)$, $y = a(1 - \cos \theta)$ is

A. $2a$

B. $2\sqrt{a}$

C. $2a/\sqrt{3}$

D. $a/\sqrt{3}$

Answer: C

 [Watch Video Solution](#)

204. If the length of the subtangent = 9 and the length of the subnormal = 4 at a point (x, y) on $y = f(x)$ then ordinate of the point =

A. 36

B. $9/4$

C. $4/9$

D. ± 6

Answer: D

 [Watch Video Solution](#)

205. The sum of the lengths of tangent and subtangent at a point of $y = a \log(x^2 - a^2)$, ($a > 0$) is proportional to

A. $|x|$

B. $|y|$

C. $|xy|$

D. $|x/y|$

Answer: C



Watch Video Solution

EXERCISE 1C (RATE OF CHANGE)

1. The distance 's' described by a particle in t seconds is given by $s = ae^t + be^{-t}$. The velocity at any time t is

A. $ae^t + be^{-t}$

B. $ae^t - be^{-t}$

C. $-ae^{-t} - be^t$

D. $ae^{-t} + be^t$

Answer: B



[Watch Video Solution](#)

2. The distance described by a particle in t seconds is given by $s = ae^t + be^{-t}$. The acceleration is

A. $ae^t + be^{-t}$

B. $ae^t - be^{-t}$

C. $-ae^{-t} - be^t$

D. $ae^{-t} + be^t$

Answer: A



Watch Video Solution

3. If the distance s travelled by a particle in time t is given by

$s = t^2 - 2t + 5$, then its acceleration is

A. 0

B. 1

C. 2

D. 3

Answer: C



Watch Video Solution

4. The distance travelled by a particle in time t is given by $s = t^3 - 2t^2 - 3t + 5$. The velocity of the particle when $t = 2$ sec is

A. 1 unit/sec

B. 2 unit/sec

C. $1/2$ unit/sec

D. 3 unit/sec

Answer: A



Watch Video Solution

5. A particle is projected vertically upward. Its height ' h ' at time ' t ' has the relation $h = 60t - 16t^2$. The velocity at which it hits the ground is

A. 60

B. 30

C. 90

D. 180

Answer: A



[Watch Video Solution](#)

6. A stone projected vertically upward moves according to the law $s = 100t - 16t^2$. The acceleration at $t = 2$ sec is

A. $-32\text{unit} / \text{sec}^2$

B. $32\text{unit} / \text{sec}^2$

C. $16\text{unit} / \text{sec}^2$

D. $8\text{unit} / \text{sec}^2$

Answer: A



Watch Video Solution

7. A particle moves along a line according to the law

$s = t^4 - 5t^2 + 8$. The initial velocity is

A. 1

B. 5

C. 4

D. 0

Answer: D



Watch Video Solution

8. The distance moved by the particle in time t is given by $s = t^3 - 12t^2 + 6t + 8$. At the instant when its acceleration is zero, the velocity is

A. 42

B. -42

C. 48

D. -48

Answer: B



Watch Video Solution

9. A particle moves along a line according to the law $s = 4t^3 - 3t^2 + 2$. At what time will the acceleration be equal to 42 unit/sec^2 ?

A. 1 sec

B. 2 sec

C. 4 sec

D. 8 sec

Answer: B



Watch Video Solution

10. The distance from a fixed point O of a particle moving in a straight line from O is given by $s = 16 + 48t - t^3$. The direction of motion of the particle after $t = 4$ sec is

- A. towards O
- B. away from O
- C. rest
- D. none

Answer: A



[Watch Video Solution](#)

11. the distance s covered in time t by a particle moving along a straight line is given by $s = \sqrt{1+t}$. Its acceleration is

proportional to the ... of its velocity at the instant

A. square

B. cube

C. double

D. none

Answer: B



[Watch Video Solution](#)

12. The velocity v and the distance s travelled by a particle has the relation $2 + 3v^2 = s^2$. Then acceleration is

A. s

B. $s/2$

C. $s/3$

D. v

Answer: C



Watch Video Solution

13. The velocity v of a point moving along a line when it is at a distance x from the origin is given by $a + bv^2 = x^2$.

Acceleration of the point at t is

A. x

B. x/b

C. x/bv

D. none

Answer: B



[Watch Video Solution](#)

14. The distance travelled by a particle in time t is given by $s = 10t - 7t^3$. The maximum velocity is

- A. 10 unit/sec
- B. 0
- C. $8/3$ unit/sec
- D. -3 unit/sec

Answer: A



[Watch Video Solution](#)

15. For particle moving in a straight line it is observed that the distance x at time t is given by $x = 6t - \frac{1}{2}t^2$. The maximum velocity during the motion is

A. 3

B. 6

C. 9

D. 12

Answer: B



[Watch Video Solution](#)

16. A particle is moving along a straight line according to the law $s = 16 + 48t - t^3$. The distance travelled by the particle before coming to rest at an instant is

A. 100 unit

B. 120 unit

C. 144 unit

D. 136 unit

Answer: C



Watch Video Solution

17. A particle moving according to the law $s = 6t - \frac{1}{2}t^3$. At

what time its velocity vanishes ?

A. 1 sec

B. 2 sec

C. 4 sec

D. 8 sec

Answer: B



Watch Video Solution

18. A particle is moving along a line according to the law $s = t^3 - 3t^2 + 5$. The acceleration of the particle at the instant where the velocity is zero is

A. 2 unit/sec²

B. 4 unit/sec²

C. 6 unit/sec²

D. 8 unit/sec²

Answer: C



Watch Video Solution

19. A particle is moving in a straight line with the relation between the time and the distance in such a way that $s = t^3 - 9t^2 + 24t - 18$. The value of its velocity when the acceleration is zero is

A. 10 unit/sec

B. 0

C. $8/3$ unit/sec

D. -3 unit/sec

Answer: D



Watch Video Solution

20. The distance s feet travelled by a particle in time t seconds is given by $s = t^3 - 6t^2 - 4t - 8$. Its acceleration vanishes at time $t =$

A. 2

B. 3

C. 4

D. 1

Answer: A



[Watch Video Solution](#)

21. A stone is thrown vertically up and the height s reached in time t is given by $s = 80t - 16t^2$. The stone reaches the maximum height in time $t =$

A. 2

B. 2.5

C. 3

D. 3.5

Answer: B



Watch Video Solution

22. A stone thrown upwards, has its equation of motion

$s = 490t - 4.9t^2$. The the maximum height reached by it is

A. 24500

B. 12500

C. 12250

D. 25400

Answer: C



Watch Video Solution

23. A stone is projected vertically upwards with an initial velocity 112 ft/sec and moves such that $s = 112t - 16t^2$ where s is the distance from the starting point and t is the time. The greatest height reached by the stone is

A. 100 ft

B. 134 ft

C. 178 ft

D. 196 ft

Answer: D



Watch Video Solution

24. A stone projected vertically upward moves according to the law $s = 48t - 16t^2$. The time taken by the stone to reach the point of projection is

A. 1 sec

B. 2 sec

C. 3 sec

D. 6 sec

Answer: C



Watch Video Solution

25. A stone is thrown vertically up and the height s reached in time t is given by $s = 80t - 16t^2$. The stone reaches the maximum height in time $t =$

A. 2

B. 2.5

C. 3

D. 3.5

Answer: B



[Watch Video Solution](#)

26. The displacement of a body of mass 100kg in a retilinear motion is given by the formula $s = 2t^2 + 3t + 1$. The K.E. of

the body 5 sec after the start is

A. 56000

B. 26450

C. 20000

D. none

Answer: B



[Watch Video Solution](#)

27. A car starts from rest and attains the speed of 10 km/hr respectively at the end of the first and second minute, If the car moves on a straight road, the distance travelled in 2 minute is

A. $1/3$ km

B. $1/4$ km

C. 15 km

D. 20 km

Answer: A



[Watch Video Solution](#)

28. A particle moves on a line according to the law $s = at^2 + bt + c$. If the displacement after one second is 16 cm, the velocity after 2 second is 24 cm/sec and the acceleration is 8 cm/sec^2 , then $(a, b, c) =$

A. (4, 8, 4)

B. (4, 4, 8)

C. (8, 4, 4)

D. (8, 8, 4)

Answer: A

 [Watch Video Solution](#)

29. The point P is moving with uniform velocity v along a straight line AB . O is a point on a perpendicular to AB at A and distance l from it. The angular velocity of P about O is

A. $\frac{lv^2}{OP}$

B. $\frac{l^2v}{OP}$

C. $\frac{lv}{OP^2}$

D. $\frac{lv^2}{OP^2}$

Answer: C



Watch Video Solution

30. If the rate of decrease of $\frac{x^2}{2} - 2x + 5$ is twice the decrease of x , then $x =$

A. 2

B. 3

C. 4

D. 1

Answer: C



Watch Video Solution

31. If the rate of change in $y = 2x^3 + 3x^2 - 30x + 7$ is 6 times the rate of change in x , then $x =$

A. 1, 5

B. 1, -5

C. 2, 3

D. 2, -3

Answer: D



[Watch Video Solution](#)

32. At the point (2, 5) on the curve $y = x^3 - 2x + 1$ the gradient of the curve is increasing

A. 6 times

B. 12 times

C. 30 times

D. 10 times as fast as x

Answer: B



Watch Video Solution

33. The point on the parabola $x^2 = 8y$ for which the abscissa and ordinate changes at the same rate

A. (1, 1)

B. (1, 2)

C. (4, 2)

D. (2, 3)

Answer: C



Watch Video Solution

34. The point on the parabola $y^2 = 4x$ for which the abscissa and ordinate changes at the same rate

- A. (1, 1)
- B. (1, 2)
- C. (4, 2)
- D. (2, 3)

Answer: B



Watch Video Solution

35. A point on the parabola $y^2 = 18x$ at which the ordinate increases at twice the rate of the abscissa is

A. (2, 4)

B. $(9/8, 9/2)$

C. $(-9/8, 9/2)$

D. (2, -4)

Answer: B



[Watch Video Solution](#)

36. A particle moves along the curve $y = x^2 + 2x$. Then the point on the curve such that x and y coordinates of the particle change with the same rate

A. (1, 3)

B. $(1/2, 5/2)$

C. $(-1/2, -3/4)$

D. (-1, -1)

Answer: C



Watch Video Solution

37. The point on the circle $x^2 + y^2 = 2$ at which the abscissa and ordinate increase at the same rate is

A. (1, -1)

B. (1, 1)

C. (-1, -1)

D. none

Answer: A



Watch Video Solution

38. At what value of an angle the rates of change in sine and tangent of the same angle are equal

A. $2n\pi$

B. $n\pi$

C. $n\pi / 2$

D. none

Answer: A



Watch Video Solution

39. A point is moving on $y = 4 - 2x^2$. The x -coordinates of the point is decreasing at the rate of 5 units per second. The rate at which y coordinates of the point is changing when the point is $(1, 2)$ is

- A. 5 unit/sec
- B. 10 unit/sec
- C. 15 unit/sec
- D. 20 unit/sec

Answer: D



[Watch Video Solution](#)

40. If the rate of change in the radius of a circle is 0.02 cm/sec, then the rate of change in the area of the circle when the radius is 5 cm is

A. π sq.cm/sec

B. 0.05 sq.cm/sec

C. 0.2π sq.cm/sec

D. 3 sq.cm/sec

Answer: C



[Watch Video Solution](#)

41. The radius of a circular plate is increasing at the rate of 0.01 cm/sec when the radius is 12 cm. Then the rate at which

the area increases is

A. 0.24π sq.cm/sec

B. 60π sq.cm/sec

C. 24π sq.cm/sec

D. 1.2π sq.cm/sec

Answer: A



[Watch Video Solution](#)

42. The radius of the circular disc increases at a uniform rate of 0.025 cm per sec. The rate at which the area of the disc increases, when the radius is 15 cm is

A. 0.75π sq.cm/sec

B. 30 sq.cm/sec

C. $30\pi \text{ sq.cm/sec}$

D. $0.4\pi \text{ sq.cm/sec}$

Answer: A



[Watch Video Solution](#)

43. If the rate of change in the radius of a circle is 0.5 sq.cm/sec , then the rate of change in the perimeter of the circle is

A. $\pi \text{ cm/sec}$

B. 0.05 cm/sec

C. $0.2\pi \text{ cm/sec}$

D. 3 cm/sec

Answer: A



Watch Video Solution

44. If the rate of change in the area of a circle is π sq.cm/sec, then the rate of change in the radius of the circle when the radius is 10 cm is

A. π cm/sec

B. 0.05 cm/sec

C. 0.2π cm/sec

D. 3 cm/sec

Answer: B

 [Watch Video Solution](#)

45. When a circular oil drop expands on water, its area increases at the uniform rate of 40 sq.cm per minute. The rate of increase in the radius when the radius 5 cm is

A. $\frac{4}{\pi}$ cm/m

B. $\frac{1}{200}$ cm/m

C. 8 cm/m

D. 4 cm/m

Answer: A

 [Watch Video Solution](#)

46. A stone is dropped into a quiet pond and waves move in circles outward from the place where it strikes, at a speed of 30 cm per second. At the instant when the radius of the wave ring is 50 m, the rate of increases in the area of the wave ring is

A. 0.75π sq.cm/sec

B. 30 sq.cm/sec

C. 30π sq.m/sec

D. 0.4π sq.cm/sec

Answer: C



Watch Video Solution

47. A stone is dropped into a quiet pond and waves move in circles outward from the place where it strikes, at a speed of 30 cm per second. At the instant when the radius of the wave ring is 50 m, the rate of increases in the circumference of the wave ring is

- A. 0.6π m/sec
- B. 6π m/sec
- C. 0.6π cm/sec
- D. 6π cm/sec

Answer: A



[Watch Video Solution](#)

48. If the rate of change of area of a circle is equal to the rate of change of its diameter then its radius =

A. $2/\pi$

B. $1/\pi$

C. $\pi/2$

D. π

Answer: B



[Watch Video Solution](#)

49. The side of a square increases at the rate of 1 cm per second. The rate at which perimeter increases is

A. 2 cm/sec

B. 4 cm/sec

C. 5 cm/sec

D. 3 cm/sec

Answer: B



Watch Video Solution

50. If the rate of change of the side of a square is 0.05 cm/sec, then the rate of change in the area of the square when the side is 10cm is

A. 0.5 sq.cm/sec

B. 1 sq.cm/sec

C. 5 sq.cm/sec

D. 10 sq.cm/sec

Answer: B



Watch Video Solution

51. the side of a square is equal to the diameter of a circle.
If the side and radius change at the same rate then the ratio of
the change of their areas is

A. $1 : \pi$

B. $\pi : 1$

C. $2 : \pi$

D. $1 : 2$

Answer: C

 [Watch Video Solution](#)

52. Two parallel sides of a rectangle are being lengthened at the rate of 2 cm/sec while the other two sides are shortened in such a way that the area of the rectangle is 50 sq.cm. The rate of change of the perimeter when the length of an increasing side 5 cm is

- A. 2 cm/sec
- B. 6 cm/sec
- C. -2 cm/sec
- D. -4 cm/sec

Answer: D

 [Watch Video Solution](#)

53. A rectangular vessel is of 2 mt long, 0.5 mt breadth and 1 mt deep. If water flows in at the rate of 900 cubic cm per sec, then the rate of increase of water level when 25 cm deep is

- A. 0.09 cm/sec
- B. 0.1 cm/sec
- C. 0.01 cm/sec
- D. 0/5 c/sec

Answer: A



[Watch Video Solution](#)

54. Each side of an equilateral triangle expands at the rate of 2 cm/sec. The rate of increase of its area when each side is 10 cm

is (in cm^2/sec)

A. $10\sqrt{2}$

B. $10\sqrt{3}$

C. 10

D. 5

Answer: B



[Watch Video Solution](#)

55. The side of an equilateral triangle increases at the uniform rate of 0.05 cm/sec. The rate of increase in the area of the triangle when the side is 20 cm is

A. 3 sq.cm/sec

B. $\frac{\sqrt{3}}{2}$ sq.cm/sec

C. 1.2 sq.cm/sec

D. 4π sq.cm/sec

Answer: B



[Watch Video Solution](#)

56. At a given instant the legs of a right angled triangle are 8 inch and 6 inch respectively. The first leg decreases at 1 inch per minute and second increases at 2 inch per minute. The rate of increasing of the area after 2 minute is

A. 1 sq.inch/min

B. 2 sq.inch/min

C. 3 sq.inch/min

D. 4 sq.inch/min

Answer: A

 [Watch Video Solution](#)

57. A variable triangle ABC is inscribed in a circle of diameter x units. At a particular instant the rate of change of side 'a' is $x/2$ time the rate of change of the opposite angle A then $A =$

A. $\frac{\pi}{4}$

B. $\frac{\pi}{3}$

C. $\frac{\pi}{2}$

D. π

Answer: B

 [Watch Video Solution](#)

58. A spherical balloon is filled with 4500π cubic meters of helium gas. If a leak in the balloon causes the gas to escape at the ratio of 72π cubic meters per minute, then the rate (in meters per minute) at which the radius of the balloon decreases 49 minutes after the leakage began is

A. $2/9$

B. $9/2$

C. $9/7$

D. $7/9$

Answer: B

 [Watch Video Solution](#)

59. Gas is leaking out of a spherical balloon at the rate of 1800 cubic cm per sec. When the radius of the balloon is 720 cm, the rate at which the surface area is shrinking is

- A. 5 sq.cm/sec
- B. 6 sq.cm/sec
- C. 10 sq.cm/sec
- D. 15 sq.cm/sec

Answer: C



[Watch Video Solution](#)

60. The volume of a metal hollow sphere is constant. If the outer radius is increasing at the rate of $\frac{1}{4}$ cm per sec. The rate

at which the inner radius is increasing when the radii are 8 cm and 4 cm respectively is

- A. 4 cm/sec
- B. 3 cm/sec
- C. 2 cm/sec
- D. 1 cm/sec

Answer: A



[Watch Video Solution](#)

61. The side of a cube is equal to the diameter of a sphere. If the side and radius increase at the same rate then the ratio of the increase of their surfaces is

A. $\pi : 6$

B. $2\pi : 3$

C. $3 : 2\pi$

D. $3 : \pi$

Answer: D



[Watch Video Solution](#)

62. The side of a cube is equal to the radius of a sphere. If the side and the radius increase at the same rate, then the relation between the rates of change of surface areas of the cube and sphere respectively is

A. $<$

B. $>$

C. =

D. none

Answer: A



[Watch Video Solution](#)

63. If the rates of increase of side of a cube and radius of a sphere are equal and rates of increase of their volumes are in the ratio $2 : 1$, then the ratio of the squares of side and radius =

A. $2\pi : 3$

B. $3\pi : 2$

C. $2 : 3\pi$

D. $8\pi : 3$

Answer: D



[Watch Video Solution](#)

64. The base radius of a cylindrical vessel full of oil is 30 cm. Oil is drawn at the rate of 27000 cubic cm per minute. The rate at which the level of the oil is falling in the vessel is

A. $30 / \pi$ cm/sec

B. π cm/sec

C. $\pi / 30$ cm/sec

D. 30 cm/sec

Answer: A



[Watch Video Solution](#)

65. Oil is being filled in a cylindrical tank of diameter 12 mt. The rate of increase in the height of oil corresponding to the rate of increase 1800π cubic cm minute in its volume is

A. $1/200$ cm/m

B. 200 cm/m

C. -200 cm/m

D. $-1/200$ cm/m

Answer: A



[Watch Video Solution](#)

66. Water is flowing into cylindrical tank of radius 7 ft in the rate of 22c.ft per sec. How fast is the water level increasing ?

A. 1 ft/sec

B. $\frac{1}{7}$ ft/sec

C. $\frac{2}{7}$ ft/sec

D. $\frac{7}{2}$ ft/sec

Answer: B



[Watch Video Solution](#)

67. The diameter and altitude of a right circular cylinder are found at a certain instant to be 20 cm and 40 cm respectively. If the diameter is increasing at the rate of 2 cm/sec then the rate of change in the altitude will keep the volume constant is

A. 2 cm/sec

B. 4 cm/sec

C. 6 cm/sec

D. -8 cm/sec

Answer: D



Watch Video Solution

68. The diameter and altitude of a right circular cylinder are found at a certain instant to be 20 cm and 40 cm respectively. If the diameter is increasing at the rate of 2 cm/sec then the rate of change in the altitude will keep the volume constant is

A. 2 cm/sec

B. 4 cm/sec

C. 6 cm/sec

D. -8 cm/sec

Answer: D



[Watch Video Solution](#)

69. An inverted cone has a depth of 10 cm and a base of radius 5 cm. Water is poured in to it at the rate of 1.5 cubic cm per second. The rate at which water is rising when the depth is 4 cm is

A. 0.5 cm/sec

B. $5/\pi$ cm/sec

C. $3/8\pi$ cm/sec

D. $8/3\pi$ cm/sec

Answer: C



[Watch Video Solution](#)

70. The radius of the base and depth of a conical funnel are 20 cm and 40 cm respectively. Water flows from the funnel at the rate 2.25 cm/sec. The rate at which the water level decreases when altitude is 30 cm is

A. $\frac{5}{8\pi}$ cm/sec

B. $\frac{1}{100\pi}$ cm/sec

C. $\frac{5}{12\pi}$ cm/sec

D. $\frac{1}{120\pi}$ cm/sec

Answer: B



Watch Video Solution

71. Water is being poured in to the inverted conical vessel at the rate of 1.5 cubic meter per minute. Its depth is always equal to twice its radius. The level of water is rising at the rate of $\frac{3}{8\pi}$ meter per minute when its depth is

- A. 1 mt
- B. 2 mt
- C. 3 mt
- D. 4 mt

Answer: D



Watch Video Solution

72. Sand is being poured on the ground from the orifice of an elevated pipe and forms a pile which has always the shape of a right circular cone whose height is equal to the radius of the base. If the sand is falling at the rate of 6 cubic ft per sec, the rate at which the height of the pile is rising when the height is 5 ft is

A. $5/8\pi$ cm/sec

B. $\frac{1}{100}\pi$ cm/sec

C. $5/12\pi$ cm/sec

D. $\frac{1}{120}\pi$ cm/sec

Answer: A



Watch Video Solution

73. Sand is being poured on the ground from the orifice of an elevated pipe and forms a pile which has always the shape of a right circular cone whose height is equal to the radius of the base. If the sand is falling at the rate of 6 cubic ft per sec, the rate at which the height of the pile is rising when the height is 5 ft is

A. $9/4\pi$ ft/sec

B. $3/8\pi$ ft/sec

C. $7/23\pi$ ft/sec

D. $6/25\pi$ ft/sec

Answer: D



Watch Video Solution

74. Water flows into a conical vessel at the rate of 5 cubic cm per second. If the semivertical angle of the vessel is 30° , then the rate of increase of water level when the water level in the vessel is 6 cm is

A. $\frac{5}{8\pi}$ cm/sec

B. $\frac{1}{100\pi}$ cm/sec

C. $\frac{5}{12\pi}$ cm/sec

D. $\frac{1}{120\pi}$ cm/sec

Answer: C



Watch Video Solution

75. A conical vessel of height 10 ft and semivertical angle 30° is full of water. It empties in such a way that the height of water in the vessel is decreasing at a constant rate of 1 inch per minute. The rate of which the volume of water in the vessel is decreasing when its height is 6 ft is

A. π c.ft/sec

B. 2π c.ft/sec

C. $1/2\pi$ c.ft/sec

D. 3π c.ft/sec

Answer: A



Watch Video Solution

76. The radius of the base of a cone is increasing at the rate of 3 cm/min and altitude is decreasing at the rate of 4 cm/min. The rate of change of lateral surface when the radius is 7 cm and altitude is 24 is

- A. 63π sq.cm/min
- B. 84π sq.cm/min
- C. 72π sq.cm/min
- D. 96π sq.cm/min

Answer: D



[Watch Video Solution](#)

77. The slant height of a cone is fixed as 7 cm. If the rate of increase in its height is 0.3 cm/sec, then the rate of increase of volume when height is 4 cm is

A. $\pi / 2$ cc/sec

B. π cc/sec

C. $\pi / 5$ cc/sec

D. $\pi / 10$ cc/sec

Answer: D



[Watch Video Solution](#)

78. If the semivertical angle of a cone is 45° then the rate of change of volume of the cone is

- A. curved area times the rate of change of r
- B. base area times the rate of change of l
- C. base area times the rate of change of r
- D. none

Answer: C



[Watch Video Solution](#)

EXERCISE 1D (MEAN VALUE THEOREMS)

1. The constant c of Rolle's theorem for the function $f(x) = 2x^3 + x^2 - 4x - 2$ in $[-\sqrt{2}, \sqrt{2}]$ is

- A. 0

B. 1

C. $\frac{1}{2}$

D. $\frac{2}{3}$

Answer: D



Watch Video Solution

2. The constant c of Rolle's theorem for the function

$f(x) = (x - a)(x - b)$ in $[a, b]$ is

A. \sqrt{ab}

B. $\frac{a + b}{2}$

C. $\frac{a - b}{2}$

D. $\frac{b - a}{2}$

Answer: B



View Text Solution

3. The constant c of Rolle's theorem for the function

$$f(x) = \log \frac{x^2 + ab}{(a+b)x} \text{ in } [a,b] \text{ where } 0 \notin [a, b] \text{ is}$$

A. \sqrt{ab}

B. $\frac{a+b}{2}$

C. $\frac{a-b}{2}$

D. $\frac{b-a}{2}$

Answer: A



Watch Video Solution

4. The constant c of Rolle's theorem for the function $f(x) = (x - a)^m(x - b)^n$ in $[a,b]$ where m,n are positive integers, is

A. $\frac{a + b}{2}$

B. $\frac{ma + nb}{m + n}$

C. $\frac{mb + na}{m + n}$

D. none

Answer: C



Watch Video Solution

5. The constant ' c ' of Rolle's theorem for the function $f(x) = \sin x$ in $[0, 2\pi]$ is

A. $\pi/6$

B. $\pi/3$

C. $\pi/4$

D. $\pi/2$

Answer: D



Watch Video Solution

6. If $a+b+c=0$ then the quadratic equation $3ax^2 + 2bx + c = 0$

has at least one root in

A. (0,1)

B. (1,3)

C. (2,3)

D. (-1,0)

Answer: A



[Watch Video Solution](#)

7. If $2a+3b+6c=0$, then at least one root of the equation $ax^2 + bx + c = 0$ lies in the interval

A. (0,1)

B. (1,3)

C. (2,3)

D. (1,2)

Answer: A



[Watch Video Solution](#)

8. Rolle's theorem can not applicable for

A. $f(x) = x^3 - 6x^2 + 11x - 6$ in $[1, 3]$

B. $f(x) = \sin x$ in $[0, \pi]$

C. $f(x) = 1 - (x - 1)^{2/3}$ in $[0, 2]$

D. $f(x) = x^2 - 3x + 2$ in $[1, 2]$

Answer: C



Watch Video Solution

9. Rolle's theorem can not applicable for

A. $f(x) = \sqrt{1 - x^2}$ in $[-1, 1]$

B. $f(x) = |x|$ in $[-1, 1]$

C. $f(x) = x^2 - 1$ in $[-1, 1]$

D. $f(x) = x^3 + x^2 - x - 1$ in $[-1, 1]$

Answer: B



[Watch Video Solution](#)

10. The constant c of the Lagrange's mean value theorem for the function $f(x) = 1 + x^2$ on $[1,2]$ is

A. $5/4$

B. $3/2$

C. $7/4$

D. $9/8$

Answer: B



[Watch Video Solution](#)

11. The constant c of Lagrange's theorem for $f(x) = x^3 - 4x^2 + 4x$ "in" $[0,2]$ is

A. 1

B. $1/2$

C. $2/3$

D. $3/2$

Answer: C



[Watch Video Solution](#)

12. The constant c of Lagrange's theorem for

$f(x) = x(x - 1)(x - 2)$ in $[0, 1/2]$ is

A. $\frac{1}{4}$

B. $\frac{6 + \sqrt{21}}{6}$

C. $\frac{6 - \sqrt{21}}{6}$

D. $\frac{\sqrt{21} - 6}{6}$

Answer: C



Watch Video Solution

13. The constant c of Lagrange's theorem for

$f(x) = (x - 1)(x - 2)(x - 3)$ in $[0, 4]$ is

A. $1 \pm \frac{2}{\sqrt{3}}$

B. $2 \pm \frac{2}{\sqrt{3}}$

C. $3 \pm \frac{2}{\sqrt{3}}$

D. $4 \pm \frac{2}{\sqrt{3}}$

Answer: B



Watch Video Solution

14. The constant c of Lagrange's theorem for

$$f(x) = \frac{x}{x-1} \text{ in } [2, 4] \text{ is}$$

A. 1

B. $\sqrt{3}$

C. $\sqrt{3} + 1$

D. $\sqrt{3} + 2$

Answer: C



Watch Video Solution

15. The constant c of Lagrange's mean value theorem for $f(x) = 2\sin x + \sin 2x$ in $[0, \pi]$ is

A. $\pi/6$

B. $\pi/4$

C. $\pi/3$

D. $\pi/2$

Answer: C



Watch Video Solution

16. A value of c for which the conclusion of Mean value Theorem holds for the function $f(x) = \log_e x$ on the interval $[1,3]$ is

A. $2 \log_3 e$

B. $\frac{1}{2} \log_e 3$

C. $\log_3 e$

D. $\log_e 3$

Answer: A



Watch Video Solution

17. The value of c in the Lagrange's mean - value theorem for $f(x) = \sqrt{x - 2}$ in the interval $[2,6]$ is

A. $\frac{5}{2}$

B. 3

C. 4

D. $\frac{9}{2}$

Answer: B



Watch Video Solution

18. The constant c of Lagrange's theorem for

$f(x) = lx^2 + mx + n$ ($l \neq 0$) in $[a, b]$ is

A. $\frac{a + b}{2}$

B. $\frac{b - a}{2}$

C. $\frac{a - b}{2}$

D. $\frac{a + b}{3}$

Answer: A

 [Watch Video Solution](#)

19. The constant c of Lagrange's theorem for $f(x) = lx^2 + mx + n$ in $[a, b]$ is

A. $\frac{1}{2}$

B. $\frac{1}{3}$

C. $\frac{1}{4}$

D. $\frac{1}{6}$

Answer: A

 [Watch Video Solution](#)

20. The constant θ of Lagrange's theorem for

$$f(x) = x^2 - 2x + 3 \text{ in } [1, 3/2] \text{ is}$$

A. $\frac{1}{2}$

B. $\frac{1}{3}$

C. $\frac{1}{4}$

D. $\frac{1}{6}$

Answer: A



[View Text Solution](#)

21. Lagrange's theorem can not be applicable for

A. $f(x) = x^2$ in $[1, 2]$

B. $f(x) = x^3$ in $[-1, 1]$

C. $f(x) = x$ in $[-1, 1]$

D. $f(x) = \frac{1}{x}$ in $[-1, 1]$

Answer: D



Watch Video Solution

22. Lagrange's theorem can not be applicable for

A. $f(x) = \sqrt{x^2} - 4$ in $[2, 4]$

B. $f(x) = |x|$ in $[-1, 2]$

C. $f(x) = x - \frac{1}{x}$ in $[1, 3]$

D. $f(x) = \log x$ in $[1, e]$

Answer: B



Watch Video Solution

23. If $f(x)$ satisfies Lagrange's mean value theorem in $[a,b]$ then there exists $c \in (a, b)$ such that

A. $f'(c) = 0$

B. $f'(c) = f(b) - f(a)$

C. the tangent at $x=c$ to the curve $y=f(x)$ is parallel to the chord joining $x=a, x=b$

D. the tangent at $x=c$ to the curve $y=f(x)$ is perpendicular to the chord joining $x=a, x=b$

Answer: C



Watch Video Solution

24. The constant c of Cauchy's mean value theorem for $f(x) = x^2, g(x) = x^3$ in $[1, 2]$ is

- A. $5/3$
- B. $5/4$
- C. $15/7$
- D. $14/9$

Answer: D



Watch Video Solution

25. The constant c of Cauchy's mean value theorem for

$f(x) = \sqrt{x}$, $g(x) = 1/\sqrt{x}$ in $[a, b]$ where $0 < a < b$ is

A. \sqrt{ab}

B. $\frac{a+b}{2}$

C. $\frac{2ab}{a+b}$

D. $\frac{1}{a} + \frac{1}{b}$

Answer: A



[Watch Video Solution](#)

26. The constant c of Cauchy's mean value theorem for the

functions $f(x) = \sqrt{x}$, $g(x) = 1/\sqrt{x}$ in $[1, 2]$ is

A. $\sqrt{2}$

B. $\sqrt{3}$

C. $3/2$

D. $5/4$

Answer: A



Watch Video Solution

27. The constant c of Cauchy's mean value theorem for $f(x) = e^x, g(x) = e^{-x}$ in $[a, b]$ is

A. \sqrt{ab}

B. $\frac{a+b}{2}$

C. $\frac{2ab}{a+b}$

D. $\frac{1}{a} + \frac{1}{b}$

Answer: B



Watch Video Solution

28. If f and g are differentiable functions in $[0,1]$ satisfying $f(0)=2=g(1),g(0)=0$ and $f(1) =6$, then for some $c \in (0, 1)$

A. $f'(c) = g'(c)$

B. $f'(c) = 2g'(c)$

C. $2f'(c) = g'(c)$

D. $2f'(c) - 3g'(c)$

Answer: B



Watch Video Solution

EXERCISE 1E (MAXIMA AND MINIMA)

1. If $x > 0$, then $f(x) = x^3 + 3x$ is

A. decreasing

B. increasing

C. oscillating

D. none

Answer: B



[Watch Video Solution](#)

2. The function $f(x) = 3x^2 - 4x$ is increasing in

A. $(2/3, \infty)$

B. $(\frac{2}{3}, 4)$

C. $(\frac{2}{3}, 2)$

D. $(\frac{2}{3}, \frac{3}{2})$

Answer: A



Watch Video Solution

3. The function $f(x) = 10 - x^3 + 3x$ is increasing in

A. $(-1/3, 1/3)$

B. $(\frac{2}{3}, 0)$

C. $(-1, 1)$

D. $(\frac{2}{3}, \frac{3}{2})$

Answer: C



Watch Video Solution

4. The function $f(x) = 3 + 12x - 9x^2 + 2x^3$ is increasing in

A. $(2/3, \infty)$

B. $(1, 5)$

C. $(2/3, 4)$

D. $(-\infty, 1) \cup (2, \infty)$

Answer: D



Watch Video Solution

5. The function $f(x)\sqrt{9-x^2}$ is increasing in

A. $(-3, 0)$

B. $(0, 4)$

C. $(-4, 0)$

D. R

Answer: A



Watch Video Solution

6. The function xe^x is increasing in

A. $(-3, 0)$

B. $(0, 4)$

C. $x > -1$

D. $(1/e, \infty)$

Answer: C



Watch Video Solution

7. If $f(x) = \log(1+x) - \frac{2x}{2+x}$ is increasing, then.....

A. $0 < x < \infty$

B. $-\infty < x < 0$

C. $-\infty < x < \infty$

D. $1 < x < 2$

Answer: A



Watch Video Solution

8. The function $f(x) = \cot^{-1} x + x$ increasing in the interval

A. $(1, \infty)$

B. $(0, \infty)$

C. $(-\infty, \infty)$

D. $(0, \infty)$

Answer: C



[Watch Video Solution](#)

9. The function $\log(\log x)$ is increasing in

A. $(1, \infty)$

B. $(0, \infty)$

C. ∞

D. R

Answer: A



[Watch Video Solution](#)

10. The function $\sinh(\sin x)$ increasing in

A. $(1, \infty)$

B. $(0, \infty)$

C. ∞

D. $\left(2n\pi - \frac{\pi}{2}, 2n\pi + \frac{\pi}{2}\right) (n \in \mathbb{Z})$

Answer: D



[Watch Video Solution](#)

11. The function $\sin(\tanh x)$ increases in

A. 0

B. \mathbb{R}

C. 1

D. ∞

Answer: B



[Watch Video Solution](#)

12. The function $\tan^{-1}(\sin x)$ increasing in

A. $\left(2n\pi - \frac{\pi}{2}, 2n\pi + \frac{\pi}{2}\right) (n \in \mathbb{Z})$

B. $\left(2n\pi + \frac{\pi}{2}, 2n\pi + \frac{3\pi}{2}\right) (n \in \mathbb{Z})$

C. $\left(2n\pi + \frac{\pi}{4}, 2n\pi + \frac{\pi}{2}\right) (n \in \mathbb{Z})$

D. $\left(2n\pi - \frac{\pi}{2}, 2n\pi + \frac{3\pi}{2}\right) (n \in \mathbb{Z})$

Answer: A



Watch Video Solution

13. The function $f(x) = \tan^{-1}(\sin x + \cos x)$ is an increasing function in

A. $(\pi/4, \pi/2)$

B. $(-\pi/2, \pi/4)$

C. $(0, \pi/2)$

D. $(-\pi/2, \pi/2)$

Answer: B



Watch Video Solution

14. $f(x) = \frac{x^2}{x+2}$ is increasing in

A. $(-2, 0)$

B. $(-4, -2)$

C. $(-4, 0)$

D. $(0, \infty)$

Answer: D



Watch Video Solution

15. The interval of increasing for $y = x - 2 \sin x$, $[0, 2\pi]$ is

- A. $(0, \pi)$
- B. $(\pi/3, \pi)$
- C. $(\pi/2, \pi)$
- D. $(0, \pi/3)$

Answer: B



[Watch Video Solution](#)

16. The interval in which $f(x) = 2x^2 - \log x$ increasing

- A. $(-1/2, 0)$
- B. $(0, 1/2)$

C. $(-\infty, -1/2)$

D. $(1/2, \infty)$

Answer: D



Watch Video Solution

17. The function $\frac{\log x}{x}$ is increasing in

A. $(1, 2e)$

B. $(0, e)$

C. $(2, 2e)$

D. $(1/e, 2e)$

Answer: B



Watch Video Solution

18. The values of x for which $x^3 - 6x^2 - 36x + 7$ increases are

A. \mathbb{R}

B. ϕ

C. $(-2, 6)$

D. $(-\infty, -2) \cup (6, \infty)$

Answer: D



[Watch Video Solution](#)

19. The values of x for which $2x^3 - 3x^2 - 36x + 10$ has extreme values are

A. \mathbb{R}

B. $(-2, 3)$

C. $(\frac{2}{3}, \infty)$

D. $(-\infty, -2) \cup (6, \infty)$

Answer: B



Watch Video Solution

20. The value of x for which $x^2 + \frac{250}{x}$ has extreme values are

A. \mathbb{R}

B. $(-2, 3)$

C. $(-2, 6)$

D. 5

Answer: D



Watch Video Solution

21. The values of x for which $(x-1)(x-2)(x-3)$ has extreme values are

A. 1,2

B. e

C. $2 \pm 1/\sqrt{3}$

D. 1

Answer: C



Watch Video Solution

22. $f(x) = \sin x$ is increasing in

A. $(\pi/2, \pi)$

B. $(\pi, 3\pi/2)$

C. $(3\pi/2, 2\pi)$

D. none

Answer: C



Watch Video Solution

23. The function $x \cdot \log \left(\frac{1+x}{x} \right)$ ($x > 0$) is increasing in

A. $(1, \infty)$

B. $(0, \infty)$

C. $(2, 2e)$

D. $(1/e, 2e)$

Answer: B



Watch Video Solution

24. The function $f(x) = x^3 - 9x^2 + 15x + 25$ is decreasing in

A. ϕ

B. \mathbb{R}

C. $(1, 5)$

D. $(-\infty, 1) \cup (5, \infty)$

Answer: C



Watch Video Solution

25. The function $f(x) \sqrt{25 - 4x^2}$ is decreasing in

A. $(-3,0)$

B. $(0,5/2)$

C. $(-5/2,0)$

D. R

Answer: B



Watch Video Solution

26. The function $f(x) = \frac{\log x}{x}$ decreases in

A. $(-\infty, e)$

B. (e, ∞)

C. $(0, e)$

D. none

Answer: B



Watch Video Solution

27. The function x^x is decreases in

A. $(0, 1/e)$

B. $(0, 4)$

C. $(-4, 0)$

D. $(1/e, \infty)$

Answer: A



Watch Video Solution

28. The function $f(x) = x^3(x - 2)^2$ decreases in

A. $(0, \infty)$

B. ∞

C. $(6/5, 2)$

D. \mathbb{R}

Answer: C



Watch Video Solution

29. The function $\cos x$ decreases in

A.

$$\left(2n\pi + \frac{\pi}{2}, 2n\pi + \pi\right) \cup \left(2n\pi + \frac{3\pi}{2}, (2n + 2)\pi\right) (n \in \mathbb{Z})$$

B. $\left(2n\pi, 2n\pi + \frac{\pi}{2}\right) \cup \left(2n\pi + \pi, 2n\pi + \frac{3\pi}{2}\right) (n \in \mathbb{Z})$

C. $\left(2n\pi, 2n\pi + \frac{\pi}{2}\right) \cup \left(2n\pi + \pi, 2n\pi + \frac{3\pi}{2}\right) (n \in \mathbb{Z})$

D. $\left(2n\pi + \frac{\pi}{2}, 2n\pi + \frac{3\pi}{2}\right) (n \in \mathbb{Z})$

Answer: B



Watch Video Solution

30. The set of all values of a for which the function

$$f(x) = \left(\frac{\sqrt{1+4}}{1-a} - 1\right)x^5 - 3x + \log 5 \text{ decreases for all real } x$$

is

A. $(-\infty, \infty)$

B. $\left[-4, \frac{3 - \sqrt{21}}{2}\right] \cup (1, \infty)$

C. $(1, \infty)$

D. $\left[-3, \frac{3 - \sqrt{27}}{2}\right] \cup (2, \infty)$

Answer: B



View Text Solution

31. $f(x) = \sec x$ is decreasing in `

A. $(-\pi/2, 0)$

B. $(0, \pi/2)$

C. $(\pi/2, \pi)$

D. $(-\pi/2, \pi/2)$

Answer: A



Watch Video Solution

32. $f(x) = \cos^{-1} x$ is decreasing in

A. $(-1,0)$

B. $(0, \pi/2)$

C. $(-1,1)$

D. none

Answer: C



Watch Video Solution

33. $f(x) = \frac{x}{a} + \frac{a}{x}$ ($a > 0$) is decreasing in

A. $-a \leq x \leq a$

B. $0 < x < a$

C. $-a < x < a$

D. $(-a, 0), \cup (0, a)$

Answer: D



Watch Video Solution

34. The value of $f(0)$ so that $f(x) = \frac{\sin x}{x}$ is continuous at $x=0$ is

A. increasing in $(0, \pi/2)$

B. decreasing in $(0, \pi/2)$

C. stationary at $x = \pi/2$

D. none

Answer: B



[Watch Video Solution](#)

35. The function $\frac{1n(1+x)}{x}$ in $(0, \infty)$ is

A. increasing

B. decreasing

C. not decreasing

D. not increasing

Answer: B



[Watch Video Solution](#)

36. The function $\frac{1}{x} \ln(1+x)$ in $(0, \infty)$ is

- A. increasing
- B. not decreasing
- C. decreasing
- D. not increasing

Answer: C



[Watch Video Solution](#)

37. The increasing function in $(0, \pi/4)$ is

A. $\cos x + \sin x$

B. $\cos x - \sin x$

C. $\frac{\sin x}{x}$

D. $\frac{x}{\sin x}$

Answer: A



[Watch Video Solution](#)

38. In the interval $(\pi/2, \pi)$

A. $f(x) = \cot x$ is increasing

B. $f(x) = \cos x$ is decreasing

C. $f(x) \tan x$ is decreasing

D. none

Answer: B



[Watch Video Solution](#)

39. In the interval $(0, \infty)$

A. $f(x)=|x|$ is increasing

B. $f(x) = e^x$ is decreasing

C. $f(x) \cos x$ is increasing

D. none

Answer: A



[Watch Video Solution](#)

40. In the interval $(-3,3)$ the function $f(x) = \frac{x}{3} + \frac{3}{x}$, $x \neq 0$ is

A. increasing

B. decreasing

C. neither increasing nor decreasing

D. partly increasing and partly decreasing

Answer: B



[Watch Video Solution](#)

41. If $y = x^3 - ax^2 + 12x + 5$ is increasing for all values of x , then a lies between

A. $-12, 12$

B. $-11, 11$

C. $-6, 6$

D. $-10, 10$

Answer: C



Watch Video Solution

42. The set of all x for which $\sin x \leq x$ is

A. $(0, \infty)$

B. $(-1, \infty)$

C. $(-1, 0)$

D. $(0, \infty)$

Answer: D



Watch Video Solution

43. $\tan x > x$ when x lies in

A. $(0, \pi/2)$

B. $(\pi/2, \pi)$

C. $(-\pi/2, 0)$

D. none

Answer: A



Watch Video Solution

44. The larger of $\sin x + \tan x, 2x$ in $0 < x < \pi/2$ is

- A. $\sin x + \tan x$
- B. $2x$
- C. cannot be determined
- D. none

Answer: A

 [Watch Video Solution](#)

45. The set of all x for which $\sin x \leq x$ is

- A. $(-\infty, 1)$
- B. $(0, \infty)$

C. $[-1, 1]$

D. $(-\infty, \infty)$

Answer: B

 [Watch Video Solution](#)

46. If $x < 0$ then $f(x) = x^2 - x$ is

A. increasing

B. decreasing

C. none

D. none

Answer: B

 [Watch Video Solution](#)

47. For real values of x , the function $x \sin x$ is

- A. decreasing
- B. increasing
- C. not decreasing
- D. not increasing

Answer: C



Watch Video Solution

48. $f(x) = x - 1/x$ is

- A. increasing in \mathbb{R}

B. decreasing in R^+

C. increasing in $R - \{0\}$

D. none

Answer: C



Watch Video Solution

49. If $f(x) = \sin x - bx + c$ decreasing along the entire number scale then

A. $b \geq 1$

B. $b > 1$

C. $b \leq 1$

D. $b < 1$

Answer: B



Watch Video Solution

50. If the function $f(x) = 2x^2 - kx + 5$ is increasing on $[1,2]$ then k lies in the interval

A. $(-\infty, 4)$

B. $(4, \infty)$

C. $(-\infty, 8)$

D. $(8, \infty)$

Answer: A



Watch Video Solution

51. If $f(x) = x^3 + ax^2 + bx + 5\sin^2 x$ is an increasing function on \mathbb{R} , then

A. $a^2 - 3b - 15 > 0$

B. $a^2 - 2b + 15 > 0$

C. $a^2 - 3b + 15 < 0$

D. $a > 0$ and $b > 0$

Answer: C



Watch Video Solution

52. If $f(x) = kx^3 - 9x^2 + 9x + 3$ is increasing on \mathbb{R} , then

A. $k < 3$

B. $k > 3$

C. $k \leq 3$

D. none

Answer: B



Watch Video Solution

53. The values of 'a' for which the function $(a + 2)x^3 - 3ax^2 + 9ax - 1$ decreases monotonically throughout for all real x are

A. $a < -2$

B. $a > -2$

C. $-3 < a < 0$

D. $-\infty < a \leq -3$

Answer: D



Watch Video Solution

54. The value of a such that $x^3 - ax^2 + 48x + 1$ increasing on \mathbb{R} is

A. $|a| \leq 12$

B. $a \leq 12$

C. 12

D. $|a|$

Answer: A



Watch Video Solution

55. The values of a, b such that $x^3 + 3ax^2 + 3a^2x + b$ is increasing on $\mathbb{R} - \{-a\}$ are

A. 1,2

B. a, b are any real numbers

C. $(-1, 2)$

D. ± 1

Answer: B



Watch Video Solution

56. The function $f(x) = x^3 + ax^2 + bx + c$, $a^2 \leq 3b$ has

A. positive real numbers $\in a^2 \leq 3b$

B. real numbers $\in a^2 \leq 3b$

C. negative real numbers $\in a^2 \leq 3b$

D. none

Answer: B



[Watch Video Solution](#)

57. The function $f(x) = x^3 + ax^2 + bx + c$, $a^2 \leq 3b$ has

A. one maximum value

B. one minimum value

C. no extreme value

D. one maximum and one minimum value

Answer: B



[Watch Video Solution](#)

58. The condition for $f(x) = x^3 + px^2 + qx + r'$, $x \in R$ to have no extreme value, is

A. $p^2 < 3q$

B. $2p^2 < q$

C. $p^2 < \frac{1}{4}q$

D. $p^2 > 3q$

Answer: A



[Watch Video Solution](#)

59. The condition that $f(x) = ax^3 + bx^2 + cx + d$ has no extreme value is

A. $b^2 = 4ac$

B. $b^2 = 3ac$

C. $b^2 < 3ac$

D. $b^2 > 3ac$

Answer: C



Watch Video Solution

60. The condition $f(x) = \frac{x}{\log x}$ has minimum value at $x =$

A. $3/2$

B. e

C. $-e$

D. 1

Answer: B



[Watch Video Solution](#)

61. The function $f(x) = xe^{-x}$ ($x \in \mathbb{R}$) attains a maximum value at $x = \dots$

A. 2

B. $\frac{1}{e}$

C. 1

D. 3

Answer: C



[Watch Video Solution](#)

62. $f(x) = \sin x(1 + \cos x)$ has maximum value at $x =$

A. 0

B. π

C. $\frac{\pi}{3}$

D. 1

Answer: C



Watch Video Solution

63. $f(x) = \sin^m x \cos^n x$ has maximum value at $x =$

A. $\tan^{-1} \sqrt{m/n}$

B. m/n

C. mn

D. 1

Answer: A



Watch Video Solution

64. The function $y = 2x^3 - 3x^2 - 12x + 8$ has minimum at $x =$

A. -1

B. 2

C. $-1/2$

D. $3/2$

Answer: B



Watch Video Solution

65. The function $y = x^4 - 6x^2 + 8x + 11$ has a minimum at $x =$

A. 1

B. -2

C. 3

D. 4

Answer: B



[Watch Video Solution](#)

66. The function $x(x-1)(x-2)$ attains its maximum value when x is

A. 1

B. $1 + \frac{a}{\sqrt{3}}$

C. $1 - \frac{1}{\sqrt{3}}$

D. $1 \pm \sqrt{3}$

Answer: C



Watch Video Solution

67. The function $f(x) = \frac{x}{2} + \frac{2}{x}$ has a local minimum at

A. $x=0$

B. $x=1$

C. $x=2$

D. $x=-2$

Answer: C



Watch Video Solution

68. The function $f(x) = x^5 - 5x^4 + 5x^3 - 1$ has

- A. one minimum and two maxima
- B. two minima and one maximum
- C. two minima and two maxima
- D. one minimum and one maximum

Answer: D



Watch Video Solution

69. Let $f(x) = a_0 + a_1x^2 + a_2x^4 + \dots + a_nx^{2n}$ be a polynomial in $x \in R$ with $0 < a_0 < a_1 < \dots < a_n$ then

$f(x)$ has

- A. neither a maximum nor a minimum
- B. only one maximum
- C. only one minimum
- D. none

Answer: C



[Watch Video Solution](#)

70. $f(x) = (\sin^{-1} x)^2 + (\cos^{-1} x)^2$ is stationary at

A. $x = 1/\sqrt{2}$

B. $x = \pi/4$

C. $x=1$

D. $x=0$

Answer: A

 [Watch Video Solution](#)

71. $f(x) = |x|$ has

A. minimum at $x=0$

B. maximum at $x=0$

C. neither max nor min at $x=0$

D. none

Answer: A

 [Watch Video Solution](#)

72. The function which has neither maximum nor minimum at $x=0$ is

A. $f(x) = x^2$

B. $f(x) = \cos x$

C. $f(x) = x^3 - 8$

D. $f(x) = \cosh x$

Answer: C



[Watch Video Solution](#)

73. The function $f(x) = \tan x$ has

A. no max points

B. no min points

C. neither max nor min points

D. none

Answer: C



Watch Video Solution

74. $f(x) = \tanh^{-1} x$ is

A. increasing in $(-1,1)$

B. decreasing in $(-1,1)$

C. max at $x=0$

D. min at $x=0$

Answer: A



Watch Video Solution

75. The least value of $(x-a)(x-b)$ occurs at $x=$

A. G.M of a,b

B. A.M of a,b

C. H.M of a,b

D. $a+b$

Answer: B



Watch Video Solution

76. Maximum value of $\frac{x}{(x+a)(x+b)}$ occurs when $x=$

A. A.M of a,b

B. G.M of a,b

C. H.M of a,b

D. none

Answer: B



[Watch Video Solution](#)

77. In the interval $[0,1]$ the function $x^{25}(1-x)^{75}$ takes a maximum value at

A. 0

B. $1/4$

C. $1/2$

D. 1/3

Answer: B

 [Watch Video Solution](#)

78. The stationary point of $f(x) = 2x^3 - 9x^2 + 12x - 3$ is

A. (1, 5), (5, 1)

B. (1, 2), (2, 1)

C. (5, 25)

D. (5, 75)

Answer: B

 [Watch Video Solution](#)

79. The stationary point of $x^2 + \frac{16}{x}$ is

A. (2,12)

B. (1,2)

C. (1,12)

D. (1,1)

Answer: A



[Watch Video Solution](#)

80. The stationary point of x^x is

A. $(e^{-1}, e^{-1/e})$

B. $(e, 1/e)$

C. (1,12)

D. (1,1)

Answer: A



[Watch Video Solution](#)

81. The stationary points of $2x^3 - 9x^2 - 24x + 16$ are

A. (- 1, 29), (4, - 96)

B. (1,29)

C. (1,1)

D. none

Answer: A



[Watch Video Solution](#)

82. The stationary value of $f(x) = \frac{\log x}{x}$ is

A. 0

B. 1

C. e

D. $1/e$

Answer: D



[Watch Video Solution](#)

83. The stationary value of $(x - 2)^{2/3}(2x - 4)$ is

A. 0

B. 2

C. 3

D. none

Answer: A



[Watch Video Solution](#)

84. The stationary value of $8x^2 - x^4 - 4$ is

A. 1,2,1

B. -4, 12, 12

C. 3,6,8

D. none

Answer: B



Watch Video Solution

85. The stationary values of $f(x) = x(\log x)^2$ are

A. $-1, 4/e$

B. $1, e^{-2}$

C. $1, 4e^2$

D. none

Answer: D



Watch Video Solution

86. The turning values of $x^3 - 3px + q$ ($p > 0$) are

A. $q + 2p\sqrt{p}, q - 2p\sqrt{p}$

B. $q + p, q - p$

C. $2p, 3p$

D. none

Answer: A



Watch Video Solution

87. If $x=-1$ and $x=2$ are extreme points of

$f(x) = \alpha \log|x| + \beta x^2 + x$ then

A. $\alpha = 2, \beta = -\frac{1}{2}$

B. $\alpha = 2, \beta = \frac{1}{2}$

C. $\alpha = -6, \beta = \frac{1}{2}$

$$D. \alpha = -6, \beta = -\frac{1}{2}$$

Answer: A



Watch Video Solution

88. The value of 'a' for which the function $f(x) = a \sin x + \frac{1}{3} \sin 3x$ has an extremum at $x = \pi/3$ is

A. 1

B. -1

C. 0

D. 2

Answer: D



Watch Video Solution

89. The set of all values of a for which the function

$$f(x) = (a^2 - 3a + 2)(\cos^2 x/4 - \sin^2 x/4) + (a - 1)x + \sin x$$

does not possess critical points is

A. $[1, \infty]$

B. $(0, 1) \cup (1, 4)$

C. $(-2, 4)$

D. $(1, 3) \cup (3, 5)$

Answer: B



[View Text Solution](#)

90. The minimum value of $(x - \alpha)(x - \beta)$ is

A. 0

B. $\alpha\beta$

C. $\frac{1}{4}(\alpha - \beta)^2$

D. $-\frac{1}{4}(\alpha - \beta)^2$

Answer: D



Watch Video Solution

91. The minimum values of $x^3 - 9x^2 + 24x - 12$ is

A. 1

B. 2

C. -8

D. 4

Answer: D



Watch Video Solution

92. If $x > 0$ the minimum value of x^x is

A. e^{-1}

B. $e^{1/e}$

C. $e^{-1/e}$

D. e

Answer: C



Watch Video Solution

93. The maximum value of $\frac{x}{1+x^2}$ is

- A. $1/2$
- B. 2
- C. $-1/2$
- D. none

Answer: C



[Watch Video Solution](#)

94. The absolute maximum of $y = x^3 - 3x + 2$ in $0 \leq x \leq 2$ is

- A. 4
- B. 6

C. 2

D. 0

Answer: A



Watch Video Solution

95. The maximum value of $f(x) = 2x^3 - 21x^2 + 36x + 20$, in the interval $0 \leq x \leq 2$ is

A. 37

B. 44

C. 32

D. 30

Answer: A



[Watch Video Solution](#)

96. If m, n are positive integers, maximum value of $x^m(a - x)^n$ in $(0, a)$ is

A. $m^m(a - m)^n$

B. $m^m n^n$

C. $\frac{m^m n^n a^{m+n}}{(m+n)^{m+n}}$

D. none

Answer: C



[View Text Solution](#)

97. If n is positive integer then greatest value of $x(a - x)^n$ on $(0, a)$ is

A. 0

B. $(2a)^{n+1}$

C. a^n

D. $\frac{a(an)^n}{(n+1)^{n+1}}$

Answer: D



[View Text Solution](#)

98. If $A > 0$, $B > 0$ and $A + B = \pi/3$, then the maximum value of $\tan A \tan B$ is

A. $1/\sqrt{3}$

B. $1/3$

C. 3

D. $\sqrt{3}$

Answer: B



Watch Video Solution

99. If $x > 0$, the maximum value of $\frac{\log x}{x}$ is

A. e

B. $2e$

C. $1/2e$

D. $1/e$

Answer: D



Watch Video Solution

100. The maximum value of x^{-x} is

A. e^e

B. e^{-e}

C. $e^{-1/e}$

D. $e^{1/e}$

Answer: D



Watch Video Solution

101. The maximum value of $x^3 - 3x$ in $[0,2]$ is

A. -2

B. 0

C. 2

D. 1

Answer: C



[Watch Video Solution](#)

102. The maximum value of $(x-1)(x-2)(x-3)$ is

A. $\frac{2}{3\sqrt{3}}$

B. $\frac{2}{\sqrt{3}}$

C. $\frac{2}{3}$

D. $\frac{1}{\sqrt{3}}$

Answer: A



Watch Video Solution

103. The maximum value of $x^4 + 3x^3 - 2x^2 - 9x + 6$ is

A. 11

B. $\frac{3}{8}$

C. 3

D. 12

Answer: A



Watch Video Solution

104. The maximum value of $\frac{x}{1+x^2}$ is

- A. $1/2$
- B. 2
- C. $-e$
- D. none

Answer: A



[Watch Video Solution](#)

105. Find the least and the greatest value of $2 \sin x + \sin 2x$ over $[0, 2\pi]$.

A. $3\sqrt{3}$

B. 3

C. $\frac{3\sqrt{3}}{2}$

D. 2

Answer: C



Watch Video Solution

106. Maximum value of $y = \sec x$ in $(\pi/2, 3\pi/2)$ is

A. $-\sqrt{2}$

B. $\sqrt{2}$

C. 1

D. -1

Answer: D



Watch Video Solution

107. The maximum value of $(\sin x)^{\sin x}$ is

A. $7/3$

B. 7

C. $\pi/2$

D. 1

Answer: D



Watch Video Solution

108. The maximum value of $y = \sin^3 x \cos x$ at $\pi/3$ is

A. $3\sqrt{3}$

B. 3

C. $\frac{3\sqrt{3}}{16}$

D. 16

Answer: C



[Watch Video Solution](#)

109. The maximum value of $a \sin x + b \cos x$ is

A. $\frac{\tan^{-1} a}{b}$

B. $\frac{a}{b}$

C. $\sqrt{a^2 + b^2}$

D. none

Answer: C



Watch Video Solution

110. The greatest value of $f(x) = 2x^2 + 2/x^2$ for $-2 \leq x < 0$, $0 < x \leq 2$ and $f(0) = 1$ is

A. $17/2$

B. 1

C. 0

D. none

Answer: A



Watch Video Solution

111. The least value of $y = \frac{a^2}{x} + \frac{b^2}{1-x}$ on $(0,1)$ is

A. 0

B. 1

C. $(a+b)$

D. $(a + b)^2$

Answer: D



View Text Solution

112. If $y = \sum_{i=1}^n (x - x_i)^2$, x_i are constants, then y has minimum value at $x =$

A. n

B. $\sum x_i$

C. $\frac{\sum x_i}{n}$

D. none

Answer: C



Watch Video Solution

113. x and y are two + ve numbers suchs that $xy=1$ Then the minimum value of $x+y$ is

A. 2

B. $\sqrt{2}$

C. 3

D. $\sqrt{3}$

Answer: A



Watch Video Solution

114. If $x, y = 12$ then the minimum value of $x^2 + y^2$ is

A. 72

B. 144

C. 48

D. 36

Answer: A



Watch Video Solution

115. The greatest value of $\sin^3 x + \cos^3 x$ in $\left[0, \frac{\pi}{2}\right]$ is

- A. 1
- B. -1
- C. 2
- D. -2

Answer: A



[Watch Video Solution](#)

116. The greatest value of xe^{-x} is

- A. $1/e$
- B. -1

C. 2

D. -2

Answer: A



[View Text Solution](#)

117. The absolute minimum of $y = c \cosh(x/c)$ is

A. $1/c$

B. $c/2$

C. c

D. $2c$

Answer: C



[Watch Video Solution](#)

118. If $a > b$, maximum value of $a \sin^2 x + b \cos^2 x$ is

A. a

B. b

C. $a+b$

D. none

Answer: A



[Watch Video Solution](#)

119. The minimum value of $27 \tan^2 \theta + 3 \cot^2 \theta$ is

A. 15

B. 18

C. 24

D. 30

Answer: B



[Watch Video Solution](#)

120. The minimum value of $a^2 \sec^2 \theta + b^2 \cos^2 \theta$ is

A. $a^2 - b^2$

B. $a^2 + b^2$

C. $(a - b)^2$

D. $(a + b)^2$

Answer: D



Watch Video Solution

121. The minimum value of $64 \sec \theta + 27 \cos ec \theta$ where θ lies in $(0, \pi/2)$ is

A. 125

B. 136

C. 142

D. 115

Answer: A



Watch Video Solution

122. The minimum value of $\sqrt{(e^{x^2}) - 1}$ is

A. 0

B. e

C. $1/e$

D. e^{e^2}

Answer: A



Watch Video Solution

123. The minimum value of $px+qy$ when $xy= r^2$ is

A. $2r\sqrt{pq}$

B. $2pq\sqrt{r}$

C. $-2r\sqrt{pq}$

D. none

Answer: A



Watch Video Solution

124. If $f(x) = x - \frac{k}{x}$ has a maximum value at $x=-2$, then $k=$

A. -1

B. -2

C. -3

D. -4

Answer: D



Watch Video Solution

125. If the function $f(x) = x^2 + \alpha/x$ has a local minimum at $x=2$, then the value of α is

- A. 8
- B. 18
- C. 16
- D. none

Answer: C



[Watch Video Solution](#)

126. The constant c of Lagrange's mean value theorem for $f(x) = 2 \sin x + \sin 2x$ in $[0, \pi]$ is

- A. $\frac{-3\sqrt{3}}{2}$

B. 3

C. $\frac{3\sqrt{3}}{2}$

D. 2

Answer: A



Watch Video Solution

127. The minimum value of $(\sin x)^{\sin x}$ is

A. $e^{-1/e}$

B. 1

C. $\pi/2$

D. $1/e$

Answer: A



Watch Video Solution

128. If x is real, then the minimum value of $y = \frac{x^2 - x + 1}{x^2 + x + 1}$ is

A. 1

B. 3

C. $1/3$

D. none

Answer: B



Watch Video Solution

129. if x is real , the maximum value of $\frac{3x^2 + 9x + 17}{3x^2 + 9x + 7}$ is

A. 1

B. $17/7$

C. $1/4$

D. 41

Answer: D



[Watch Video Solution](#)

130. If x, y are strictly positive such that $x+y=1$ then the minimum value of $x \log x + y \log y$ is

A. $\log 2$

B. $-\log 2$

C. $2 \log 2$

D. 0

Answer: B

 [Watch Video Solution](#)

131. If $l^2 + m^2 = 1$ then the max value of $l+m$ is

A. 1

B. $\sqrt{2}$

C. $1/\sqrt{2}$

D. 2

Answer: B

 [Watch Video Solution](#)

132. If x is real and $\frac{a \sin x + b \cos x}{c \sin x + d \cos x}$ has neither maximum nor minimum then

A. $a/d = c/b$

B. $a/b = d/c$

C. $a/c = b/d$

D. $a/c \neq b/d$

Answer: D



[Watch Video Solution](#)

133. The greatest value of the function $f(x) = \sin 2x - x$ on $[-\pi/2, \pi/2]$ is

A. $\frac{\sqrt{3}}{2} - \frac{\pi}{6}$

B. $\frac{\sqrt{3}}{2} + \frac{\pi}{6}$

C. $-\frac{\sqrt{3}}{2} + \frac{\pi}{3}$

D. $\frac{1}{2} - \frac{\pi}{3}$

Answer: A



Watch Video Solution

134. The least value of $f(x) = \sin 2x - x$ on $[-\pi/2, \pi/2]$ is

A. $\frac{\sqrt{3}}{2} - \frac{\pi}{6}$

B. $\frac{\sqrt{3}}{2} + \frac{\pi}{6}$

C. $\frac{\pi}{6} - \frac{\sqrt{3}}{2}$

D. $\frac{1}{2} - \frac{\pi}{6}$

Answer: C



Watch Video Solution

135. The difference between the greatest and least value of the function $f(x) = \sin 2x - x$ on $[-\pi/2, \pi/2]$ is

A. $\frac{\sqrt{3} + \sqrt{2}}{2}$

B. $\frac{\sqrt{3} + \sqrt{2}}{2} + \frac{\pi}{6}$

C. $\sqrt{3} - \frac{\pi}{3}$

D. $\frac{\sqrt{3} + \sqrt{2}}{2} - \frac{\pi}{3}$

Answer: C



Watch Video Solution

136. For $a > 0$, if the function $f(x) = 2x^3 - 9ax^2 + 12a^2x + 1$ attains its maximum value at p and minimum value at q such that $p^2 - q$ then $a =$

A. 3

B. 1

C. 2

D. $1/2$

Answer: C



[Watch Video Solution](#)

137. If m and M respectively denote the minimum and maximum of $f(x) = (x - 1)^2 + 3$ for $x \in [-3, 1]$ then the

ordered pair $(m, M) =$

A. $(-3, 19)$

B. $(3, 19)$

C. $(-19, 3)$

D. $(-19, -3)$

Answer: B



[Watch Video Solution](#)

138. Let $f: \mathbb{R} \rightarrow \mathbb{R}$ be defined, by
$$\begin{cases} k - 2x & \text{if } x \leq -1 \\ 2x + 3 & \text{if } x > -1 \end{cases}$$

If f has a local minimum at $x = -1$, then a possible value of k is

A. 1

B. 0

C. $-\frac{1}{2}$

D. -1

Answer: D



Watch Video Solution

139. p and q are distinct prime numbers and if the equation $x^2 - px + q = 0$ has positive integer as its roots then the roots the roots of the equation are

A. The cubic has minima at $-\frac{\sqrt{p}}{3}$ and maxima at $\sqrt{\frac{p}{3}}$

B. The cubic has minima at both $\frac{\sqrt{p}}{3}$ and $-\sqrt{\frac{p}{3}}$

C. The cubic has maxima at both $\sqrt{\frac{p}{3}}$ and $-\sqrt{\frac{p}{3}}$

D. The cubic has minima at $\sqrt{\frac{p}{3}}$ and maxima at $-\sqrt{\frac{p}{3}}$

Answer: D



Watch Video Solution

140. Given $P(x) = x^4 + ax^3 + bx^2 + cx + d$ such that $x=0$ is the only real root of $P'(x) = 0$. If $P(-1) < P(1)$, then in the interval $[-1,1]$

- A. $P(-1)$ is not minimum but $P(1)$ is the maximum of P
- B. $P(-1)$ is the minimum but $P(1)$ is not the maximum of P
- C. neither $P(-1)$ is the minimum and $P(1)$ is the maximum of P
- D. $P(-1)$ is the minimum and $P(1)$ is the maximum of P

Answer: A



[Watch Video Solution](#)

141. A curve passes through the point $(2,0)$ and the slope of the tangent at any point is $x^2 - 2x$ for all values of x . The point of maximum or minimum of the curve is

A. $(0, 2/3)$

B. $(0, 4/3)$

C. $(0, 1/3)$

D. $(0, 5/3)$

Answer: B

[Watch Video Solution](#)

142. The point on the curve $y = \frac{x}{1+x^2}$ where the tangent to the curve has the greatest slope is

A. (0,0)

B. (1,1)

C. (1,0)

D. (2,3)

Answer: A



[Watch Video Solution](#)

143. The curve $y = ax^2 + bx$ has minimum at (2,-12) on it.

Then (a,b) =

A. (3, - 12)

B. $(-3, 12)$

C. $(-3, -12)$

D. $(3, 12)$

Answer: A



[Watch Video Solution](#)

144. If $f(x) = a/x + bx$ has minimum at $(2,1)$ then $(a,b) =$

A. $(1, 1/4)$

B. $(1/4, 1)$

C. $(1, 4)$

D. $(2, 1)$

Answer: A



Watch Video Solution

145. If $y = \frac{ax + b}{(x - 1)(x - 4)}$ has a maximum value at the point

(2,-1) then

A. $a=10, b=20$

B. $a=1, b=0$

C. $a=5, b=5$

D. none

Answer: B



Watch Video Solution

146. If a quadratic function in x has the value 9 when $x=1$ and has maximum value 10 when $x=2$ then the function is

A. $-x^2 + 4x + 6$

B. $x + 4x - x^2$

C. $8 + 4x - x^2$

D. none

Answer: A



Watch Video Solution

147. A cubic function of x has maximum 10 and minimum $-5/2$, when $x = -3$ and $x = 2$ respectively. Find the function.

$$A. \frac{1}{5}x^3 + \frac{3}{10}x^2 = \frac{18}{5}x + \frac{19}{10}$$

B. $\frac{1}{15}x^3 + \frac{3}{10}x^2 - \frac{18}{5}x + \frac{16}{100}$

C. $ax^2 + bx + c = 0$

D. none

Answer: A



[Watch Video Solution](#)

148. The real number x when added to its inverse gives the minimum value of the sum at $x =$

A. 2

B. 1

C. -1

D. -2

Answer: B



Watch Video Solution

149. If the product of two positive numbers is 400 then the minimum value of their sum is

A. 8

B. 12

C. 32

D. 40

Answer: D



Watch Video Solution

150. The sum of two positive numbers is 12. The numbers so that the sum of the squares is minimum are

A. 6,6

B. 15,38

C. 24,24

D. 38,50

Answer: A



Watch Video Solution

151. The difference of two positive numbers is 10 . If the square of the greater exceeds twice the square of the smaller by maximum value then they are

A. 15,5

B. 20,10

C. 30,20

D. none

Answer: B



Watch Video Solution

152. The sum of three numbers is 30 . The first plus three times the second plus four times the third add up to 80. the numbers so that the product of all three is as large as possible are

A. 12,10,10

B. 10,10,10

C. 12,12,12

D. 10,12,12

Answer: B



Watch Video Solution

153. Divide 64 into two parts such that the sum of the cubes of two parts is minimum. The parts are

A. 32,30

B. 32,32

C. 40,,42

D. 42,42

Answer: B



[Watch Video Solution](#)

154. The maximum value of xy subject to $x+y = 7$ is

A. 12

B. 10

C. $49/4$

D. $55/4$

Answer: C



[Watch Video Solution](#)

155. The sum of two numbers is 20 . If the product of the square of one number and cube of the other is maximum ,

then the numbers are

A. 10,10

B. 11,9

C. 8,12

D. 14,6

Answer: C



[Watch Video Solution](#)

156. The ratio of the two parts of a number 'a' such that the product of the p^{th} power of one and q^{th} power of the other is maximum is

A. $P^2 : q^2$

B. $p:q$

C. $p:p+q$

D. $q:p+q$

Answer: B



Watch Video Solution

157. If x, y, k, m, n are positive and $x + y = k$, then find the maximum value of $x^m y^n$.

A. $\frac{k^{m+n} m^m n^n}{(m+n)^{m+n}}$

B. $(m+n)^n$

C. $k^{m+n} n^m$

D. none

Answer: A



Watch Video Solution

158. The difference of a number and its square is maximum, then the number is

A. $1/2$

B. 2

C. 1

D. 0

Answer: A



Watch Video Solution

159. If α, β are the roots of the quadratic equation $x^2 - (a - 2)x - (a + 1) = 0$, where a is a variable, then the least value of $\alpha^2 + \beta^2$ is

A. 3

B. 5

C. 7

D. none

Answer: B



Watch Video Solution

160. The value of a so that the sum of the squares of roots of the equation $x^2 - (a - 2)x - a + 1 = 0$ assume the least value is

A. 2

B. 0

C. 3

D. 1

Answer: D



Watch Video Solution

161. The function $f(x) = a \sin x + \frac{1}{3} \sin 3x$ has maximum value at $x = \frac{\pi}{3}$. The value of a is

A. 3

B. $\frac{1}{3}$

C. 2

D. $1/2$

Answer: C



[Watch Video Solution](#)

162. The focal distance of the point $(4,2)$ on the parabola $x^2 = 8y$ is

A. $\sqrt{2}$

B. $2\sqrt{2}$

C. $3\sqrt{2}$

D. $4\sqrt{2}$

Answer: B



[Watch Video Solution](#)

163. The equation of the normal to the curve $x^2 = 4y$ at $(2, 1)$ is

- A. (4,4)
- B. (1,2)
- C. (9,6)
- D. (4,5)

Answer: B



[Watch Video Solution](#)

164. The point on the curve $y = x^2$ which is nearest to $(3,0)$ is

A. $(1, -1)$

B. $(-1, 1)$

C. $(-1, -1)$

D. $(1,1)$

Answer: D



Watch Video Solution

165. The point on the curve $x^2 = 2y$ which is closest to the point $(0,5)$ is

A. $(\pm 2\sqrt{2}, 3)$

B. $(\pm 2\sqrt{2}, 4)$

C. $(\pm \sqrt{2}, 3)$

D. $(\pm \sqrt{3}, 4)$

Answer: B

 [Watch Video Solution](#)

166. The minimum distance from the origin to a point on the curve $x^{2/3} + y^{2/3} = a^{2/3}$ ($a > 0$) is

A. a

B. $a/2$

C. $a/\sqrt{8}$

D. $a^{2/3}$

Answer: B

 [Watch Video Solution](#)

167. The shortest distance from $(-6,0)$ to $x^2 - y^2 + 16 = 0$ is

A. $3\sqrt{5}$

B. $\sqrt{34}$

C. 5

D. none

Answer: B



[View Text Solution](#)

168. The points on $y = x^2 + 7x + 2$ which is closest to the line $y=3x-3$ is

A. (-2,-4)

B. (-2,-8)

C. (2,8)

D. (2,4)

Answer: B



Watch Video Solution

169. The point on the curve $y = x^2 + 4x + 3$ which is closest to the line $y=3x+2$ is

A. $\left(\frac{1}{2}, \frac{5}{4}\right)$

B. $\left(-\frac{1}{2}, \frac{5}{4}\right)$

C. $\left(2, -\frac{5}{3}\right)$

D. $\left(2, \frac{5}{3}\right)$

Answer: B

 [Watch Video Solution](#)

170. The shortest distance between the line $y-x = 1$ and the curve $x = y^2$ is

A. $\frac{2\sqrt{3}}{8}$

B. $\frac{3\sqrt{2}}{5}$

C. $\frac{\sqrt{3}}{4}$

D. $\frac{3\sqrt{2}}{8}$

Answer: D

 [Watch Video Solution](#)

171. The longest distance of the point $(a,0)$ from the curve

$$2x^2 + y^2 = 2x \text{ is}$$

A. $1+a$

B. $|1-a|$

C. $\sqrt{1 - 2a + 2a^2}$

D. $\sqrt{1 - 2a + 3a^2}$

Answer: C



Watch Video Solution

172. Point P is $(-2,-3)$ and point Q is $(3,7)$. The point A on the axis for which $PA+AQ$ is least is $(-,0)$. Then A =

A. $(-1, 2, 0)$

B. $(1/2, 0)$

C. $(1, 2)$

D. $(-1, 0)$

Answer: A



[Watch Video Solution](#)

173. A line is drawn through the point $(1, 2)$ to meet the coordinate axes at P and Q such that it forms a triangle OPQ , where O is the origin. If the area of the triangle OPQ is least, then the slope of the line PQ is

A. -2

B. $-1/2$

C. $-1/4$

D. -4

Answer: A



Watch Video Solution

174. If the perimeter of a maximum rectangle is constant , then that rectangle

A. is a square

B. in not a square

C. may or may not be a square

D. none

Answer: A



[Watch Video Solution](#)

175. The maximum area of the rectangle that can be inscribed in a circle of radius r is

- A. is a square
- B. is not a square
- C. may or may not be a square
- D. none

Answer: A



[Watch Video Solution](#)

176. The triangle of maximum area that can be inscribed in a circle is

- A. is a square
- B. is not a square
- C. may or may not be a square
- D. none

Answer: A



[Watch Video Solution](#)

177. The maximum area of the rectangle that can be inscribed in a circle of radius r is

- A. r^2

B. r^3

C. $r^2 / 4$

D. $2r^2$

Answer: D



Watch Video Solution

178. The sides of the greatest rectangle that can be inscribed

in $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$ are

A. $a\sqrt{2}, b\sqrt{2}$

B. \sqrt{a}, \sqrt{b}

C. a, b

D. none

Answer: A



[Watch Video Solution](#)

179. Area of the largest rectangle that can be inscribed in the

ellipse $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$ is

A. πab

B. ab

C. $2 ab$

D. $1/2 ab$

Answer: C



[Watch Video Solution](#)

180. The sides of a rectangle, with maximum perimeter, inscribed in a semicircle of radius R are

A. $\frac{R}{2}, \frac{R}{2}$

B. $\frac{4R}{\sqrt{5}}, \frac{R}{\sqrt{5}}$

C. $\frac{3R}{\sqrt{5}}, \frac{2R}{\sqrt{5}}$

D. none

Answer: B



[View Text Solution](#)

181. A wire of length l is cut into two parts which are bent respectively in the form of a square and a circle. What are the lengths of pieces of wire so that the sum of areas is least ?

A. $\frac{a}{\pi + 4}$

B. $\frac{a^2}{4(\pi + 4)}$

C. $\frac{a^2}{\pi + 4}$

D. $\frac{100}{\pi + 4}$

Answer: B

 [Watch Video Solution](#)

182. A wire of length 20cm is cut into two parts which are bent in the form of a square and a circle, then the least value of the sum of areas so formed is

A. $\frac{400}{\pi + 4}$

B. $\frac{20}{\pi + 4}$

C. $\frac{5}{\pi + 4}$

D. $\frac{100}{\pi + 4}$

Answer: D

 [Watch Video Solution](#)

183. A window is in the shape of a rectangle surmounted by a semi-circle. If the perimeter of the window be 20 feet then find the maximum area.

A. $\frac{k^2}{\pi + 4}$ sq. unit

B. $\frac{k}{\pi + 4}$ sq. unit

C. $\frac{k^2}{2(\pi + 4)}$ sq. unit

D. $\frac{k}{2(\pi + 4)}$ sq. unit

Answer: C



Watch Video Solution

184. A window is in the shape of a rectangle surmounted by a semi-circle. If the perimeter of the window be 20 feet then find the maximum area.

A. $\frac{200}{\pi + 4}$ sq .ft.

B. 200 sq. ft

C. $\frac{\pi}{200}$ sq. ft.

D. none

Answer: A



Watch Video Solution

185. A line segment of length 10 cm is divided into two parts and a rectangle is formed with these as adjacent sides, then the dimensions of the rectangle in order that its area is maximum is

A. 4,6

B. 5,5

C. 2,8

D. none

Answer: B



Watch Video Solution

186. A $(0,a)$, $B(0,b)$ be fixed points , $P(x,0)$ a variable point. The angle $\angle APB$ is maximum if

A. $x^2 = ab$

B. $x=ba$

C. $2x^2 = 2ab$

D. none

Answer: A



[Watch Video Solution](#)

187. A line segment of length 8cm is divided into two parts AP and PB by a point P. If $AP^2 + PB^2$ is minimum then AP=

A. P is the midpoint of AB

B. P is a point of trisection of AB

C. P divides AB in the ratio 1:3

D. none

Answer: A



[Watch Video Solution](#)

188. ABCD is rectangle in which $AB= 9\text{cm}$, $BC =6\text{ cm}$. P is a point in CD such that $PC=x$. If $AP^2 + PB^2$ is minimum then $x=$

A. $2/9\text{ cm}$

B. $9/2\text{ cm}$

C. 9 cm

D. 2 cm

Answer: B



Watch Video Solution

189. The triangle of maximum area that can be inscribed in a circle is

- A. an isosceles
- B. right angled
- C. an equilateral
- D. none

Answer: C



Watch Video Solution

190. The maximum value of the area of the triangle with vertices $(a,0)$, $(a \cos \theta, b \sin \theta)$, $(a \cos \theta, -b \sin \theta)$ is

A. $\frac{3\sqrt{3ab}}{4}$

B. $3\sqrt{ab}$

C. $\frac{\sqrt{3ab}}{4}$

D. $3\sqrt{3ab}$

Answer: A



Watch Video Solution

191. Find the triangle of the greatest area among the right triangles of a given perimeter p .

A. $\frac{p^2}{2}$

B. $\frac{p^2}{2(2 + \sqrt{2})^2}$

C. $\sqrt{2^2}$

D. $2p^2$

Answer: B



Watch Video Solution

192. The maximum area of triangle formed by a tangent line to the curve $x^{2/3} + y^{2/3} = 1$ and the coordinates axes is

A. $1/4$ sq . Units

B. $1/2$ sq. units

C. 1 sq. units

D. none

Answer: A



[View Text Solution](#)

193. The sum of the hypotenuse and a side of a right angled triangle is constant . If the area of the triangle is maximum then the angle between the hypotenuse and the given side is

A. $\pi / 2$

B. $\pi / 3$

C. $\pi / 4$

D. π

Answer: B



[Watch Video Solution](#)

194. Two sides of a triangle is given . If the area of a triangle is maximum, then the angle between the two sides is

A. $\pi / 2$

B. $\pi / 6$

C. π

D. none

Answer: A



Watch Video Solution

195. A triangular park is enclosed on two sides by a fence and on the third side by a straight river bank. The two sides having

fence are of same length x . The maximum area enclosed by the park is

A. $\frac{1}{2}x^2$

B. πx^2

C. $\frac{3}{2}x^2$

D. $\frac{\sqrt{x^3}}{8}$

Answer: A



Watch Video Solution

196. The point P in the first quadrant of the ellipse $x^2/8 + y^2/18 = 1$ so that the area of the triangle formed by the tangent at P and the coordinate axes is least

A. (2,3)

B. $(\sqrt{8}, 0)$

C. $(\sqrt{18}, 0)$

D. none

Answer: A



Watch Video Solution

197. A straight line through the point (3,4) in the first quadrant meets the axes at A and B .

The minimum area of the triangle OAB is

A. 24 sq. unit

B. 42 sq. unit

C. 22 sq .unit

D. 12 sq. unit

Answer: A



Watch Video Solution

198. Through the point $(2,3)$,a straight line is drawn making, positive intercept on the coordinate axes, . The area of the triangle thus formed is least when the ratio of the intercepts on the x and y axes is

A. 1: 2

B. 3: 1

C. 2: 3

D. none

Answer: C



Watch Video Solution

199. The perimeter of a sector is given. The area is maximum when the angle of the sector is

A. $\pi^c / 6$

B. $\pi^c / 4$

C. 4^c

D. 2^c

Answer: D



Watch Video Solution

200. If the perimeter of a sector of a circle is constant then find the angle of the sector, when its area is maximum.

A. $c^2 / 16 \text{sq. cm}$

B. $c^2 / 8 \text{sq. cm}$

C. $c^2 / 4 \text{sq. cm}$

D. none

Answer: A



[Watch Video Solution](#)

201. Twenty meters of wire is available to fence off a flower bed in the form of a sector. If the flower bed has the maximum surface then radius is

A. 10

B. $\frac{5}{2}$

C. 5

D. $15/2$

Answer: C



[View Text Solution](#)

202. A wire of length 20 cm can be bent in the form of a sector then its maximum area is

A. 15 sq. cm

B. 25 sq.cm

C. 5 sq. cm

D. none

Answer: B

 [View Text Solution](#)

203. A box is made from a piece of metal sheet 24 cms square by cutting equal small squares from each corner and tranning up the edges If the volume of the box is maximum then then the dimensions of the box are

A. 2,8,8

B. 2,6,8

C. 4,6,8

D. 2,4,4

Answer: A



[Watch Video Solution](#)

204. An open top box of maximum possible volume from a square piece of tin of side 'a' is to be made by cutting equal squares out of the corners and then folding up the tin to form the sides. The length of a side of square cut out is

A. $a/6$

B. $a/4$

C. $a/3$

D. $a/2$

Answer: A



[View Text Solution](#)

205. From a rectangular sheet of dimensions $30\text{cm} \times 80\text{cm}$, four squares of sides x cm are removed at the corners, and the sides are then turned up so as to form an open rectangular box. What is the value of x , so that the volume of the box is the greatest?

- A. $20/3$
- B. $10/3$
- C. $15/2$
- D. 5

Answer: A



[Watch Video Solution](#)

206. The strength of a beam varies as the product of its breadth b and square of its depth d . A beam cut of a circular log of radius r would be strong when

A. $b = d = \frac{r}{2}$

B. $b^2 = \frac{r}{2}\sqrt{2} = d$

C. $d = \sqrt{2}b = \sqrt{2/3} \cdot 2r$

D. $d = \sqrt{3}b = \sqrt{3/2} \cdot 2r$

Answer: C



[Watch Video Solution](#)

207. The height of the cylinder of maximum volume which can be inscribed in a sphere of radius ' r ' is

A. $\sqrt{3}r$

B. $r / \sqrt{3}$

C. $2r / \sqrt{3}$

D. $r / r\sqrt{3}$

Answer: C



Watch Video Solution

208. The radius of right circular cylinder of maximum volume which can be inscribed in a sphere of radius r is

A. r

B. $r / 2$

C. $\sqrt{2/3}r$

D. $\sqrt{3/2}r$

Answer: C



View Text Solution

209. The maximum volume of the cylinder which can be inscribed in a sphere of radius a

A. $\frac{4\pi a^3}{3\sqrt{3}}$ cubic unit

B. $4\pi a^3$ cubic unit

C. $\frac{4\pi a^3}{\sqrt{3}}$ cubic unit

D. none

Answer: A



View Text Solution

210. The volume of the greatest cylinder which can be inscribed in a cone of height h and semi-vertical angle α is

A. $\frac{4\pi h^3}{27} \tan^2 \alpha$

B. $4\pi h^2 \tan^2 \alpha$

C. $\frac{4\pi h^3}{9} \tan^2 \alpha$

D. none

Answer: A



[Watch Video Solution](#)

211. The height and the radius of the base of a cylinder of maximum volume, given the sum of the height and the

diameter of the base of the cylinder is 3 unit are

A. 2,2

B. 1,1

C. 10,10

D. none

Answer: B



[View Text Solution](#)

212. The dimensions of the greatest cylinder that can be inscribed in a sphere of radius a are

A. $\frac{2a}{\sqrt{3}}, \frac{a\sqrt{2}}{\sqrt{3}}$

B. $\frac{2a}{3}, \frac{a}{3}$

C. $\frac{a}{\sqrt{3}}, \frac{2a}{\sqrt{3}}$

D. none

Answer: A



[View Text Solution](#)

213. Show that when the curved surface of a right circular cylinder inscribed in a sphere of radius R is maximum, then the height of the cylinder is $\sqrt{2R}$.

A. \sqrt{R}

B. $\sqrt{10R}$

C. $\sqrt{2R}$

D. R

Answer: C



Watch Video Solution

214. A cylindrical, gas container is closed at the top and open bottom. If the iron plate of the top is $\frac{5}{4}$ times as thick as the plate forming the cylindrical sides, the ratio of the radius to the height of the cylinder using minimum material for the same capacity is

A. 4

B. 5

C. $\frac{4}{5}$

D. 20

Answer: C



Watch Video Solution

215. The height of the cylinder of maximum volume which can be inscribed in a sphere of radius 'r' is

A. $\frac{R}{3}$

B. $\frac{2R}{3}$

C. $\frac{4R}{3}$

D. $\frac{4R}{\sqrt{3}}$

Answer: C



Watch Video Solution

216. The height of the cone of maximum volume which can be inscribed in a sphere of radius 6 is

A. 8

B. 4

C. 2

D. 24

Answer: A



[View Text Solution](#)

217. The semivertical angle of a cone of maximum volume and of given total surface area is

A. $\sin^{-1} \sqrt{2}$

B. $\sin^{-1} 1/3$

C. $\tan^{-1} \sqrt{2}$

D. $\tan^{-1} 1/3$

Answer: B



[Watch Video Solution](#)

218. A conical tent of given capacity will require the least amount of canvas when the height is times the radius of the tent

A. 1

B. 2

C. $\sqrt{3}$

D. $\sqrt{2}$

Answer: D



View Text Solution

219. The semivertical angle of the cone of maximum volume and of given slant height is

A. $\tan^{-1}(\sqrt{2})$

B. $\cos^{-1}(\sqrt{2})$

C. $\sin^{-1}(\sqrt{2})$

D. none

Answer: A



Watch Video Solution

220. If h is the height of the maximum cone inscribed in a sphere of radius r then $h : r =$

A. 4 : 3

B. 3 : 4

C. 2 : 1

D. 1 : 1

Answer: A



[Watch Video Solution](#)

221. The maximum volume of the right circular cone that can be inscribed in a sphere of radius R

A. $\frac{32}{27}\pi R^3$ cubic units

B. $\frac{32}{81}\pi R^3$ cubic unit

C. $85\pi R^3$ cubic unit

D. none

Answer: B



Watch Video Solution

EXERCISE 2 SET-1 (SPECIAL TYPE QUESTIONS)

1. If $f(x) = x^2 + 3x$, $x = 10$, $\delta x = 0.01$ then

I: $\delta f = 0.2301$

II: $df = 0.23$

III: relative error in x is 1

A. only I, III are true

B. only II, III are true

C. only I, II are true

D. I, II, III are true

Answer: C



[View Text Solution](#)

2. There is an error of 0.02 cm is is made in measuring the radius 10 cm of a circle. Then

I: Approximate error in area is 0.5 sq. cm

II: Approximate percentage error in area is 0.4

A. only I is true

B. only II is true

C. both I and II are true

D. neither I nor II true

Answer: B



[View Text Solution](#)

3. Semivertical angle of a cone is 45° and height is $30 \cdot 05$ cm

I : Error in volume is 45π cubic cm . Approximately

II : Percentage error in volume is $1/2$

A. only I is true

B. only II is true

C. both I and II are true

D. neither I nor II true

Answer: D



[View Text Solution](#)

4. Equation of the tangent to the curve $y = 2x^3 - 6x^2 - 9$ at the point where the curve crosses the y-axis is

- A. only I is true
- B. only II is true
- C. I and II are true
- D. neither I nor II true

Answer: A



[Watch Video Solution](#)

5. Equation of the tangent to the curve $y = 2x^3 - 6x^2 - 9$ at the point where the curve crosses the y-axis is

- A. only I is true
- B. only II is true
- C. I and II are true
- D. neither I nor II true

Answer: B



[Watch Video Solution](#)

6. The angle between the curves $y = x$, $y = 1/x$ at $(1, 1)$ is

- A. only I is true
- B. only II is true

C. I and II are true

D. neither I nor II true

Answer: C



Watch Video Solution

7. Observe the following statements for the curve $y^2 = 4ax$.

I : The length of the subnormal at any point is a constant.

II : the length of the sub- tangent at any point is twice the abscissa of the point of contact

III : Area of triangle formed by tangent normal and x-axis at any point is a constant.

A. only I,II are true

B. only II,III are true

C. only I,III are true

D. I,II,III are true

Answer: A



[View Text Solution](#)

8. Observe the following statements for the curve $x = a (\cos t + \log \tan \frac{1}{2}t)$, $y = a \sin t$

I : Slope of the tangent at any point is $\tan t$

II : Length of the tangent at any point is constant

III : Length of the sub-tangent at any point is $|a \cos t|$

A. only I,II are true

B. only II,III are true

C. only I,III are true

D. I,II,III are true

Answer: D



[View Text Solution](#)

9. If displacement s , time t are related by $s = \sqrt{t}$ then

I : Acceleration is proportional to velocity.

II : Velocity is inversely proportional to displacement.

A. only I is true

B. only II is true

C. both I and II are true

D. neither I nor II true

Answer: B



[View Text Solution](#)

10. I : A particle is projected vertically upward its height h at time t is given by $h = 60t - 16t^2$. The velocity at which it hits the ground is 60 units /sec.

II : A stone is thrown up vertically and the height h reached in time t given by $h = 80t - 16t^2$. The stone reaches the maximum height in $5/2$ secs.

- A. only I is true
- B. only II is true
- C. both I and II are true
- D. neither I nor II true

Answer: C



[View Text Solution](#)

11. A man of height 180 cm walks at a uniform rate of 12 km/hr away from the lamp post of height 450 cm . Then

I : Rate at which the length of shadow increases is 8 km/hr

II : Rate at which the tip of shadow is moving is 20 km /hr

- A. only I is true
- B. only II is true
- C. both I and II are true
- D. neither I nor II true

Answer: C



[View Text Solution](#)

12. I : The function $\log(\log x)$ increases in $(1, \infty)$.

II : The function x^x is decreasing in $(0, 1/e)$.

- A. only I is true
- B. only II is true
- C. both I and II are true
- D. neither I nor II true

Answer: C



[View Text Solution](#)

13. I : The function $f(x) = xe^{-x}$ has maximum at $x=e$.

II : The function $f(x) = \sin x(1 + \cos x)$ has maximum at $x = \pi/3$.

- A. only I is true

B. only II is true

C. both I and II are true

D. neither I nor II true

Answer: B



[View Text Solution](#)

14. The maximum value of $x^4 + 3x^3 - 2x^2 - 9x + 6$ is

A. only I is true

B. only II is true

C. both I and II are true

D. neither I nor II true

Answer: B



Watch Video Solution

EXERCISE 2 SET-2 (SPECIAL TYPE QUESTIONS)

1. If $f(x) = 2x^2 + 3x - 5$, $x = 3$, $\delta x = 0.1$ then $\delta f = A$

(2) If $f(x) = x^2 + 4x$, $x = 2$, $\delta x = 0.1$ then $\delta f = B$

(3) If $f(x) = x^2 + 3x$, $x = 3$, $\delta x = 0.1$ then $\delta f = C$

The ascending order of A, B C is

A. A,B,C

B. B,C,A

C. C,A,B

D. A,C,B

Answer: B

 [Watch Video Solution](#)

2. IF an error of 0.01 cm is made while measuring the radius 2 cm of a circle, then the relative error in the circumference is

A. A,B,C

B. B,C,A

C. C,A,B

D. A,C,B

Answer: C

 [Watch Video Solution](#)

3. Radius of a sphere is 2 cm and error in it is $\frac{1}{10}$ cm then arrange the approximate values of the following in decending order

- A) Error in diameter
- B) Error in Circumference
- C) Error in area
- D) Relative error in radius

A. A,B,C,D

B. D,A,C,B

C. B,D,A,C

D. D,C,B,A

Answer: D



Watch Video Solution

4. If A,B,C,D are the length of tangents to the curves

1) $y = 4x^2$ at $(-1,4)$

2) $y = x^3 + 1$ at $(1,2)$ 3) $y = \frac{x^3}{2-x}$ at $(1,1)$

4) $2x^2 + 3xy - 2y^2 = 8$ at $(2,3)$ then the ascending order of

A,B,C,D is

A. A,B,C,D

B. B,C,D,A

C. A,B,D,C

D. C,B,D,A

Answer: D



[View Text Solution](#)

5. If A,B,C,D are the length of normals to the curves 1) $y = 4x^2$ at (-1,4)

2) $y = x^3 + 1$ at (1,2) 3) $y = \frac{x^3}{2-x}$ at (1,1)

4) $2x^2 + 3xy - 2y^2 = 8$ at (2,3) then the ascending order of A,B,C,D is

A. A,B,C,D

B. B,C,D,A

C. A,B,D,C

D. C,B,D,A

Answer: D



View Text Solution

6. If A,B,C,D are the lengths of subtangents to the curves 1)

$$\sqrt{x} + \sqrt{y} = 3 \text{ at } (4,1)$$

$$2) x^2 y^2 = 1 \text{ at } (-1,1)$$

$$3) y = \frac{x+1}{x} \text{ at } (1,2)$$

4) $x^2 + xy + y^2 = 7$ at $(1,-3)$ then the descending order of

A,B,C,D is

A. A,B,C,D

B. D,C,B,A

C. A,B,D,C

D. C,B,D,A

Answer: B



[View Text Solution](#)

7. The lengths of tangent, subtangent, normal and subnormal for the curve $y = x^2 + x - 1$ at (1, 1) are A, B, C and D respectively, then their increasing order is

A. B,D,A,C

B. B,A,C,D

C. A,B,C,D

D. B,A,D,C

Answer: D



[Watch Video Solution](#)

8. If A,B ,C are the maximum velocities of the particles moving according to the law

$s = 60t - 5t^3$, $s = 6t - \frac{1}{2}t^2$, $s = 10t - 7t^3$ respectively then

the ascending order of A,B,C is

A. A,B,C

B. C,B,A

C. B,A,C

D. B,C,A

Answer: D



[View Text Solution](#)

9. If A,B,C are the maximum heights reached when three stones projected vertically upwards moves according to the law

$s = 128t - 16t^2$, $s = 48t - 16t^2$, $s = 80t - 16t^2$ respectively

then the descending order of A,B,C is

A. A,C,B

B. C,B,A

C. B,A,C

D. B,C,A

Answer: A



[View Text Solution](#)

10. If A,B,C are the minimum values of $2x^3 - 3x^2 - 12x + 5$, $x^3 - 9x^2 + 24x - 12$, $x^3 - 6x^2 + 9x + 1$ then the ascending order of A,B,C is

A. A,B,C

B. B,C,A

C. C,A,B

D. A,C,B

Answer: D



[View Text Solution](#)

11. The functions $y = x^4 - 6x^2 + 8x + 15$ has minimum at $x=A$, $y=x(x-1)(x-2)$ has maximum at $x=B$, $y = 2x^3 - 3x^2 - 12x + 5$ has minimum at $x=C$. The ascending order of A,B,C is

A. A,B,C

B. B,C,A

C. C,A,B

D. A,C,B

Answer: A



View Text Solution

EXERCISE 2 SET-3 (SPECIAL TYPE QUESTIONS)

1. Match the following

- | | |
|---|----------|
| I. The approximate value of $(2 \cdot 001)^4$ | a 0.4983 |
| II. The approximate value of $(1 \cdot 0002)^{3000}$ | b 2.02 |
| III. The approximate value of $\sqrt{4 \cdot 08}$ | c 1.6 |
| IV. The approximate value of $\frac{1}{3\sqrt{8 \cdot 08}}$ | d 16.032 |

A. c,b,d,a

B. d,c,b,a

C. a,c,b,d

D. c,b,d,a

Answer: C



View Text Solution

2. Match the following

- | | | |
|---|---|---------|
| I. The approximate value of $\sin(30^\circ 1')$ | a | 1.00058 |
| II. The approximate value of $\cos(61^\circ)$ | b | 1.0349 |
| III. The approximate value of $\tan(46^\circ)$ | c | 0.4849 |
| IV. The approximate value of $(45^\circ 1')$ | d | 0.50025 |

A. c,b,d,a

B. d,c,b,a

C. c,d,a,b

D. c,b,d,a

Answer: B



View Text Solution

3. If side of a cube is 10 cm and error in it is 0.05 cm then

match the following

- | | | |
|--------------------------------------|---|------|
| I. Error in surface are of cube | a | 15 |
| II. Percentage error in surface area | b | 6 |
| III. Error in volume | c | 1.5 |
| IV. Percentage error in volume | d | 0.05 |
| | e | 1 |

A. b,d,e,a

B. a,c,c,d

C. a,c,b,e

D. b,e,a,c

Answer: D



View Text Solution

4. The slope of the tangent to the curve $y = 6 + x - x^2$ at (2,4) is

A. a,b,c,d

B. b,c,d,a

C. c,d,b,a

D. c,d,a,b

Answer: D



[Watch Video Solution](#)

5. Match the points on the curve $2y^2 = x + 1$ with the slope of normals at those points

I. $(7, 2)$ $a - 4\sqrt{2}$
II. $(0.1/\sqrt{2})$ $b - 8$
III. $(1, -1)$ $c 4$
IV. $(3, \sqrt{2})$ $d 0$
 $e - 2\sqrt{2}$

A. b,d,c,a

B. b,e,c,a

C. b,c,c,a

D. b,e,a,c

Answer: B



[View Text Solution](#)

6. Match the following

I. length of tangent at $(1, 2)$ on the curve $y = x^3 + 1$ is

a) $2\sqrt{5}$

II. length of normal at $(1, 2)$ on the curve $y = x^2 + 1$ is

b) 3

III. length of sub-tangent at any point on the curve $y = be^{x/a}$ is

c) $\frac{2\sqrt{10}}{3}$

IV. length of the normal at the point $(1, 1)$ on the curve $y = x^3$ is

d) $\sqrt{5}$

e) $|a|$

A. a,c,b,d

B. c,b,e,a

C. c,a,d,b

D. c,a,e,b

Answer: D



View Text Solution

7. Match the following

- I. Angle between the curves $y^2 = 4x$, $x^2 = 2y - 3$ at $(1, 2)$ a) 90°
II. Angle between the curves $xy = 4$, $x^2 - y^2 = 15$ at $(-4, -1)$ b) 0°
III. Angle between the curves $y^2 = x$, $x^2 = y$ at $(1, 1)$ c) $\tan^{-1} 3$
IV. Angle between the curves $x^2 = 4y$, $x^2 + y^2 = 5$ at $(-2, 1)$ d) $\tan^{-1} (3/4)$

A. a,b,c,d

B. b,c,d,a

C. c,d,b,a

D. b,a,d,c

Answer: D



View Text Solution

8. The rate of change of radius of a circle is 1 cm / sec. Match the following

- | | |
|--|------------|
| I. The rate of change of area when radius = 5 cm | a) 2π |
| II. The rate of change of area when radius = 3 cm | b) 6π |
| III. The rate of change of area when radius = 4 cm | c) 8π |
| IV. The rate of change of perimeter when radius = 3 cm | d) 10π |

A. d,b,c,a

B. d,c,b,a

C. c,d,b,a

D. a,d,b,c

Answer: A



View Text Solution

9. Match the following

- | | |
|--|--|
| I. Rate of increase in area of equilateral triangle of side 15cm, when each side is increasing at the rate of 0.1 cm/s; is | a) 67.5 c.c/s |
| II. Rate of increase in area of square of side 15 cm and each side is increasing at the rate of 0.1 cm/s is | b) $3\sqrt{3}/2 \text{ cm}^3/\text{s}$ |
| III. Rate of increase in volume of the cube of side 15 cm and each side is increasing at the rate of 0.1 cm/s is | c) $6\text{cm}^2/\text{s}$ |
| | d) $3\text{cm}^2/\text{s}$ |

A. d,a,b

B. c,a,d

C. c,a,b

D. a,b,d

Answer: A



View Text Solution

10. Match the following

I. $f(x) = x^2 - 2x + 5$ is increasing in

II. $f(x) = x^x$ decreases for

III. $f(x) = x + \frac{1}{x}$ is increasing in

IV. $f(x) = 9 - 6x - 2x^2 - x^3$ is decreasing for

a) $x > 1/e$

b) $(-\infty, -1) \cup (1, \infty)$

c) $(1, \infty)$

d) $(-\infty, \infty)$

e) $0 < x < 1/e$

A. b,a,e,c

B. c,e,d,b

C. c,e,d,b

D. d,a,c,b

Answer: C



View Text Solution

11. Match the following

Function

i. $\frac{\log x}{e}$

ii. x^{-1}

iii. $a \sin^2 x + b \cos^2 x$ ($a > b$)

iv. $(\sin x)^{\sin x}$

Maximum value

a) $e^{1/e}$

b) $e^{-1/e}$

c) $1/e$

d) b

e) a

A. a,b,d,e

B. c,d,b,a

C. d,b,e,a

D. c,a,c,b

Answer: D



View Text Solution

12. Match the following

- I. If $l^2 + m^2 = 1$ then maximum value of $l+m$ is a) $\sqrt{2}$
II. Minimum value of $x^3 - 9x^2 + 24x - 12$ is b) 2
III. Least value of $f(x) = \sin 2x - x$ on $\left[-\frac{\pi}{2}, \frac{\pi}{2}\right]$ is c) 4
IV. $f(x) = |x - 2|$ is least when $x =$ d) $\frac{\pi}{6} - \frac{\sqrt{3}}{2}$
e) 0

A. a,c,d,b

B. a,c,d,b

C. a,b,c,d

D. c,a,d,e

Answer: A



View Text Solution

13. Match the following

- | | |
|---|------------------|
| I. The point on the curve $y^2 = 4x$ nearest to $(2, 1)$ is | a) $(1, 1)$ |
| II. The point on the curve $y = x^2$ nearest to $(3, 0)$ is | b) $(1, 2)$ |
| III. The point on $y = x^2 + 7x + 2$ nearest to $y = 3x - 3$ is | c) $(-2, -8)$ |
| IV. The point on $y = x^2 + 4x + 3$ nearest to $y = 3x + 2$ is | d) $(-1/2, 5/4)$ |

A. a,b,c,d

B. b,d,a,c

C. b,a,c,d

D. d,c,b,a

Answer: C



View Text Solution

EXERCISE 2 SET-4 (SPECIAL TYPE QUESTIONS)

1. A : In triangle ABC if a, A and R are fixed then

$$\Delta a \cdot \sec A + \Delta b \cdot \sec B + \Delta c \cdot \sec C = 0$$

R : In any triangle , $A+B+C = \pi$ and so $\Delta A + \Delta B + \Delta C = 0$

- A. A,R are true and R is correct explanation of A
- B. A,R are true of but R is not correct explanation of A
- C. A is true , R is false
- D. A is false, R is true

Answer: A



[View Text Solution](#)

2. A : if semivertical angle of a cone is 45° and height of the cone is 20.025 then approximate value of its volume is 10π cubic units.

R : If semivertical angle of a cone is α and height is h then volume of cone is $\frac{\pi}{3}h^3 \tan^2 \alpha$

- A. A,R are true and R is correct explanation of A
- B. A,R are true of but R is not correct explanation of A
- C. A is true, R is false
- D. A is false, R is true

Answer: D



[View Text Solution](#)

3. Assertion (A) : If semi vertical angle of a cone is 45° and height is 30.05 cm then approximate volume of cone is $9045.08 \pi \text{c.c}$

Reason(R) : When semi vertical angle is 45° approximate error in volume is $\delta v = \pi r^2 \delta h$

- A. A,R are true and R is correct explanation of A
- B. A,R are true of but R is not correct explanation of A
- C. A is true , R is false
- D. A is false, R is true

Answer: D



Watch Video Solution

4. A : The gradient of the curve $y = x^3 - 3x^2 - 2x + 7$ at (1,3) is -5.

R : The gradient of the curve $y = f(x)$ at P is $\left(\frac{dy}{dx}\right)_p$

- A. A and R are true and R is the correct explanation of A
- B. A and R are true and R is not correct explanation of A
- C. A is true but R is false
- D. A is false but R is true

Answer: A



[View Text Solution](#)

5. A : Area of triangle formed by tangent to the curve $xy = c^2$ with coordinates axes is $2c^2$ sq . Units.

R: Area of triangle formed by the line $\frac{x}{a} + \frac{y}{b} = 1$ with coordinate axes is $\frac{1}{2} |ab|$ sq. units

- A. A and R are true and R is the correct explanation of A
- B. A and R are true and R is not correct explanation of A
- C. A is true but R is false
- D. A is false but R is true

Answer: A

 [View Text Solution](#)

6. The curves $y = x^3 - 3x^2 - 8x - 4$, $y = 3x^2 + 7x + 4$ touch at the point $(-1, 0)$. The equation of the common tangent is

- A. A and R are true and R is the correct explanation of A
- B. A and R are true and R is not correct explanation of A
- C. A is true but R is false
- D. A is false but R is true

Answer: A



Watch Video Solution

7. A : The curve $y = x^2$, $6y = 7 - x^3$ cut orthogonally at (1,1) .

R : Two curve cut each other orthogonally at their point of intersection P iff $m_1 m_2 = -1$ where m_1, m_2 are the gradients of the two curves at P .

- A. A and R are true and R is the correct explanation of A
- B. A and R are true and R is not correct explanation of A

C. A is true but R is false

D. A is false but R is true

Answer: A



Watch Video Solution

8. A :A particle moves along a line is given by

$$s = \frac{t^3}{3} - 3t^2 + 8t + 5 \text{ .its direction changes only when } t=2.4$$

R : The direction of a body changes only when sign of velocity changes

A. A,R are true and R is correct explanation of A

B. A,R are true and R is correct explanation of A

C. A is true, R is false

D. A is false ,R is true

Answer: A

 [Watch Video Solution](#)

9. A : The smallest value of $x^2 - 3x + 3$ in $[-3, 3/2]$ is $3/4$

R: The smallest value of $f(x)$ in $[a, b]$ is equal to the local minimum of $f(x)$

A. Both A and R are true and R is correct explanation of A

B. Both A and R are true but R is not correct explanation of

A

C. A is true R is false

D. A is false but R is true

Answer: C



[View Text Solution](#)

10. A : The function $f(x) = 2x^3 - 3x^2 - 12x + 8$ has minimum value

R: For the above function $f'(2) = 0$ and $f''(2) > 0$

A. Both A and R are true and R is correct explanation of A

B. Both A and R are true but R is not correct explanation of

A

C. A is true R is false

D. A is false but R is true

Answer: A



11. A : If $x+y=12$ then the minimum value of $x^2 + y^2$ is 72

R : If $x+y=k$ then the maximum value of xy is k^2

- A. Both A and R are true and R is correct explanation of A
- B. Both A and R are true but R is not correct explanation of A
- C. A is true R is false
- D. A is false but R is true

Answer: A

12. Observe the following statements :

Assertion (A) : $f(x) = 2x^3 - 9x^2 + 12x - 3$ is increasing outside the interval (1,2)

Reason (R) : $f'(x) < 0$ for $x \in (1, 2)$

Then which of the following is true

- A. Both A and R are true, and R is not the correct reason for A
- B. Both A and R are true and R is the correct reason for A
- C. A is true but R is false
- D. A is false but R is true

Answer: A



Watch Video Solution

13. The function $f(x) = xe^{-x}$ ($x \in R$) attains a maximum value at $x = \dots$

- A. Both (A) and (R) are true and (R) is the correct reason for (A).
- B. Both (A) and (R) are true, but (R) is not the correct reason for (A)
- C. (A) is true, (R) is false
- D. (A) is false, (R) is true

Answer: A



[Watch Video Solution](#)

14. Consider the functions, $f(x) = |x - 2| + |x - 5|$, $x \in \mathbb{R}$

Statement-1 : $f'(4) = 0$ Statement-2 : f is continuous in $[2, 5]$,
differentiable in $(2, 5)$ and $F(2) = F(5)$

- A. Statement 1 is true , statement 2 is true, statement 2 is not a correct explanation for statement 1
- B. statement 1 is true, statement 2 is false
- C. statement 1 is false, statement 2 is true
- D. statement 1 is true , statement 2 is true, statement 2 is a correct explanation for statement 1

Answer: A



Watch Video Solution

15. Let $a, b \in R$ be such that the function f given by $f(x) = \ln|x| + bx^2 + ax, x \neq 0$ has extreme values at $x = -1$ and $x = 2$

Statement-I : f has local maximum at $x = -1$ and $x = 2$.

Statement-II: $a = \frac{1}{2}, b = \frac{-1}{4}$

- A. Statement 1 is true , statement 2 is true, statement 2 is not a correct explanation for statement 2
- B. statement 1 is true, statement 2 is false
- C. statement 1 is false, statement 2 is true
- D. statement 1 is true , statement 2 is true, statement 2 is a correct explanation for statement 2

Answer: D



Watch Video Solution

