



## MATHS

### BOOKS - DEEPTI MATHS (TELUGU ENGLISH)

### EAMCET - 2016 TS

#### Questions

1. The length of the segment of the straight line passing through  $(3,3)$  and  $(7,6)$  and off by the coordinate axes is

A.  $4/5$

B.  $5/4$

C.  $7/4$

D.  $4/7$

**Answer: B**



**Watch Video Solution**

2. Find the value of  $k$ , if the straight lines  $y - 3kx + 4 = 0$  and  $(2k - 1)x + (8k - 1)y - 6 = 0$  are perpendicular.

A.  $1/6$

B.  $-1/6$

C. 1

D. 0

**Answer: A**

3. The combined equation of the straight lines of the form  $y = kx + 1$  (where  $k$  is an integer such that the point of intersection of each with the line  $3x + 4y = 9$  has an integer as its  $x$ -coordinate is

A.  $(y+x+1)(y+2x-1) = 0$

B.  $(y+x-1)(y+2x+1) = 0$

C.  $(y+x+1)(y+2x+1) = 0$

D.  $(y+x-1)(y+2x-1) = 0$

**Answer: D**

4. If the axes are rotated anticlockwise through an angle  $90^\circ$  then the equation  $x^2 = 4ay$  is changed to the equation

A.  $y^2 = 4ax$

B.  $x^2 = -4ay$

C.  $y^2 = -4ax$

D.  $x^2 = 4ay$

**Answer: A**



[Watch Video Solution](#)

5. If A(5, -4) and B(7, 6) are points in a plane, then the set of all points P(x, y) in the plane such that AP : PB = 2 : 1 is

A. a circle

B. a hyperbola

C. an ellipse

D. a parabola

**Answer: A**



[View Text Solution](#)

6. If on an average, out of 10 ships, one is drowned, then what is the probability that out of 5 ships atleast 4 reach safely ?

A.  $14(0.9)^5$

B.  $1.4(0.9)^5$

C.  $0.14(0.9)^4$

D.  $1.4(0.9)^4$

**Answer: D**



**Watch Video Solution**

7. In a family with 4 children, the probability that there are at least two girls is

A.  $1/2$

B.  $9/16$

C.  $3/4$

D.  $11/16$

**Answer: D**



**Watch Video Solution**

8. When a pair of six faced fair dice are thrown, the probability that the sum of the numbers on the two dice is greater than 7 , is

- A.  $1/3$
- B.  $5/12$
- C.  $1/2$
- D.  $1/4$

**Answer: B**



[Watch Video Solution](#)

9. A five digit number is formed by the digits 1,2,3,4,5 with no digit being repeated. The probability that the number is divisible by 4, is

A.  $1/5$

B.  $2/5$

C.  $3/5$

D.  $4/5$

**Answer: A**



**Watch Video Solution**

**10.** Two events  $A$  and  $B$  are such that  $P(A) = \frac{1}{2}$ ,  $P(A | B) = \frac{1}{4}$  and  $P(B | A) = \frac{1}{2}$  Consider the

following statements :

(I)  $P(\bar{A} | \bar{B}) = \frac{3}{4}$

(II)  $A$  and  $B$  are mutually exclusive

(III)  $P(A | B) + P(A | \bar{B}) = 1$ . Then



- A. Only (I) is correct
- B. Only (I) and (II) are correct
- C. Only (I) and (III) are correct
- D. Only (II) and (III) are correct

**Answer: A**



**Watch Video Solution**

11. The standard deviation of  $a, a+d, a + 2d, \dots, a + 2nd$  is

A.  $nd$

B.  $n^2d$

C.  $\frac{\sqrt{n(n+1)}}{3}d$

D.  $\frac{\sqrt{n(n+3)}}{3}d$

**Answer: C**



**Watch Video Solution**

12. If the average of the first  $n$  numbers in the sequence 148, 146, 144,....., is 125, then  $n =$

A. 18

B. 24

C. 30

D. 36

**Answer: B**



**Watch Video Solution**

13.  $a$  is perpendicular to both  $b$  and  $c$ . The angle between  $b$  and  $c$  is  $\frac{2\pi}{3}$ . If  $|a| = 2$ ,  $|b| = 3$ ,  $|c| = 4$ , then  $a(b \times c) =$

A.  $18\sqrt{3}$

B.  $12\sqrt{3}$

C.  $8\sqrt{3}$

D.  $6\sqrt{3}$

**Answer: B**



**Watch Video Solution**

14. If  $a, b, c$  are unit vectors satisfying the relation  $a + b + \sqrt{3}c = 0$ , then the angle between  $a$  and  $b$  is

A.  $\pi/6$

B.  $\pi/4$

C.  $\pi/3$

D.  $\pi/2$

**Answer: C**



**Watch Video Solution**

15.  $a, b, c$  are three vectors such that  $|a| = 1, |b| = 2, |c| = 3$  and  $b, c$  are perpendicular. If projection of  $b$  on  $a$  is the same as the projection of  $c$  on  $a$ , then  $|a - b + c| =$

A.  $\sqrt{2}$

B.  $\sqrt{7}$

C.  $\sqrt{14}$

D.  $\sqrt{21}$

**Answer: C**



**Watch Video Solution**

**16.** The vectors  $2i - 3j + k$ ,  $I - 2j + 3k$ ,  $3i + j - 2k$

A. are linearly dependent

B. are linearly independent

C. form sides of a triangle

D. are coplanar

**Answer: B**



**Watch Video Solution**

17. ABCD is a parallelogram and P is the mid point of the side AD.

The line BP meets the diagonal AC in Q. Then the ratio AQ : QC

=

A. 1 : 2

B. 2 : 1

C. 1 : 3

D. 3 : 1

**Answer: A**



[Watch Video Solution](#)

18. If ABCDEF is a regular hexagon with centre O , then P.T

$$\overline{AB} + \overline{AC} + \overline{AD} + \overline{AE} + \overline{AF} = 3\overline{AD} = 6\overline{AO}$$

A.  $2\vec{AO}$

B.  $3\vec{AO}$

C.  $\vec{AO}$

D.  $6\vec{AO}$

**Answer: D**



**Watch Video Solution**

**19.** In  $\triangle ABC$ , if  $2R + r = r_2$  then  $\angle B =$

A.  $\pi/3$

B.  $\pi/4$

C.  $\pi/6$

D.  $\pi/2$

**Answer: D**



**Watch Video Solution**

20. In  $\triangle ABC$ , if  $8R^2 = a^2 + b^2 + c^2$ , then the triangle is a

- A. right angled triangle
- B. equilateral triangle
- C. scalene triangle
- D. obtuse angled triangle

**Answer: A**



**Watch Video Solution**



21. If  $\Delta ABC$  is such that  $\angle A = 90^\circ$ ,  $\angle B \neq \angle C$ , then

$$\frac{b^2 + c^2}{b^2 - c^2} \sin(B - C) =$$

A.  $1/3$

B.  $1/2$

C.  $1$

D.  $3/2$

**Answer: C**



[View Text Solution](#)

22. For  $\theta \in \left(0, \frac{\pi}{2}\right)$ ,  $\sec h^{-1}(\cos \theta) =$

A.  $\log \left| \tan \left( \frac{\pi}{6} + \frac{\theta}{2} \right) \right|$

B.  $\log \left| \tan \left( \frac{\pi}{3} + \frac{\theta}{2} \right) \right|$

C.  $\log \left| \tan \left( \frac{\pi}{4} + \frac{\theta}{2} \right) \right|$

D.  $\log \left| \tan \left( \frac{\pi}{4} - \frac{\theta}{2} \right) \right|$

**Answer: C**



**View Text Solution**

**23.** The value of  $x$  which satisfies

$$\sin(\cot^{-1} x) = \cos(\tan^{-1} (1 + x))$$

A.  $-1/2$

B.  $1/2$

C.  $-1$

D.  $1$

**Answer: A**



**Watch Video Solution**

**24.** If  $f(x) = \cos^2 x + \cos^2 2x + \cos^2 3x$ , then the number of values of  $x \in [0, 2\pi]$  for which  $f(x) = 1$  is

A. 4

B. 6

C. 8

D. 10

**Answer: D**



**View Text Solution**

25. If  $\cos x + \cos y + \cos \alpha = 0$  and  $\sin x + \sin y + \sin \alpha = 0$ ,

then  $\cot\left(\frac{x+y}{2}\right) =$

A.  $\sin \alpha$

B.  $\cos \alpha$

C.  $\tan \alpha$

D.  $\cot \alpha$

**Answer: D**



**Watch Video Solution**

26. 
$$\frac{\cos 13^\circ - \sin 13^\circ}{\cos 13^\circ + \sin 13^\circ} + \frac{1}{\cot 148^\circ} =$$

A. 1

B. -1

C. 0

D.  $1/2$

**Answer: C**



[View Text Solution](#)

27. If  $f(x) = x^2 - 2x + 4$  then the set of values of  $x$  satisfying  $f(x - 1) = f(x + 1)$  is

A.  $\{-1\}$

B.  $\{-1, 1\}$

C.  $\{1\}$

D.  $\{1, 2\}$

**Answer: C**



Watch Video Solution

28. The number of real linear functions  $f(x)$  satisfying

$$f(f(x)) = x + f(x) \text{ is}$$

A. 0

B. 4

C. 5

D. 2

**Answer: D**



Watch Video Solution

29. The remainder when  $7^n - 6n - 50 (n \in \mathbb{N})$  is divided by 36, is

A. 22

B. 23

C. 1

D. 21

**Answer: B**



[View Text Solution](#)

30. Consider the system of equations

$$ax + by + cz = 2$$

$$bx + cy + az = 2$$

$$cx + ay + bz = 2$$

where  $a, b, c$  are real numbers such that  $a + b + c = 0$ . Then the system

- A. has two solutions
- B. is inconsistent
- C. has unique solution
- D. has infinitely many solutions

**Answer: B**



[Watch Video Solution](#)

**31.** Suppose  $A$  and  $B$  are two square matrices of same order. If  $A, B$  are symmetric matrices, then  $AB - BA$  is

- A. a symmetric matrix



B. a skew symmetric matrix

C. a scalar matrix

D. a triangular matrix

**Answer: B**



**Watch Video Solution**

32.  $A(x) = \begin{vmatrix} x + 1 & 2x + 1 & 3x + 1 \\ 2x + 1 & 3x + 1 & x + 1 \\ 3x + 1 & x + 1 & 2x + 1 \end{vmatrix}$  then  $\int_0^1 A(x) dx =$

A.  $-15$

B.  $-15/2$

C.  $-30$

D.  $-5$

**Answer: B**



**Watch Video Solution**

33. If  $z = x + iy$  is complex number such that  $z^{1/3} = a + ib$ , then the value of  $\frac{1}{(a^2 + b^2)} \left( \frac{x}{a} + \frac{y}{b} \right) =$

A.  $-1$

B.  $-2$

C.  $0$

D.  $2$

**Answer: B**



**Watch Video Solution**

34. The locus of  $z$  satisfying  $|z| + |z-1| = 3$  is

- A. a circle
- B. a pair of straight lines
- C. an ellipse
- D. a parabola

**Answer: C**



[View Text Solution](#)

35. If the point  $z = (1 + i)(1 + 2i)(1 + 3i)\dots(1 + 10i)$  lies on a circle with centre at origin and radius  $r$ , then  $r^2 =$

- A.  $10!$
- B.  $2 \times 3 \times 4 \times \dots \times 10$

C.  $2 \times 5 \times 10 \times \dots \times 101$

D.  $11!$

**Answer: C**



**Watch Video Solution**

**36.** The minimum value of  $|z - 1| + |z - 5|$  is

A. 5

B. 4

C. 3

D. 2

**Answer: B**



**Watch Video Solution**

37. The number of real roots of  $|x^2| - 5|x| + 6 = 0$  is

A. 2

B. 3

C. 4

D. 1

**Answer: C**



**Watch Video Solution**

38. If  $\alpha, \beta$  are the roots of  $x^2 - x + 1 = 0$  then the quadratic equation whose roots are  $\alpha^{2015}, \beta^{2015}$  is

A.  $x^2 - x + 1 = 0$

B.  $x^2 + x + 1 = 0$

C.  $x^2 + x - 1 = 0$

D.  $x^2 - x - 1 = 0$

**Answer: A**



**Watch Video Solution**

**39.** If  $\alpha, \beta, \gamma$  are roots of  $x^3 - 5x + 4 = 0$  then

$$(\alpha^3 + \beta^3 + \gamma^3)^2 =$$

A. 12

B. 13

C. 169

D. 144

**Answer: D**



**Watch Video Solution**

**40.** Suppose  $\alpha, \beta, \gamma$  are roots of  $x^3 + x^2 + 2x + 3 = 0$ . If  $f(x) = 0$  is a cubic polynomial equation whose roots are  $\alpha + \beta, \beta + \gamma, \gamma + \alpha$  then  $f(x) =$

A.  $x^3 + 2x^2 - 3x - 1$

B.  $x^3 + 2x^2 - 3x + 1$

C.  $x^3 + 2x^2 + 3x - 1$

D.  $x^3 + 2x^2 + 3x + 1$

**Answer: C**



**View Text Solution**

41. The number of 4 letter words that can be formed with the letters in the word EQUATION with at least one letter repeated is

- A. 2400
- B. 2408
- C. 2416
- D. 2432

**Answer: C**



[Watch Video Solution](#)

42. The number of divisors of  $7!$  is

- A. 24



B. 72

C. 64

D. 60

**Answer: D**



**Watch Video Solution**

**43.** The sum of the series

$$1 + \frac{2}{3} \left( \frac{1}{8} \right) + \frac{2 \times 5 \times 8}{3 \times 6 \times 9} \left( \frac{1}{8} \right)^3 + \dots \text{ is}$$

A.  $\frac{4}{3\sqrt{49}}$

B.  $\frac{3\sqrt{49}}{4}$

C.  $\frac{4}{3\sqrt{81}}$

D.  $\frac{3\sqrt{81}}{4}$

**Answer: A**



**Watch Video Solution**

**44.** If  $C_r$  denotes the binomial coefficient  ${}^n C_r$  then

$$(-1)C_0^2 + 2C_1^2 + 5C_2^2 + \dots + (3n - 1)C_n^2 =$$

A.  $(3n - 2) \cdot {}^{2n} C_n$

B.  $\left(\frac{3n - 2}{2}\right) \cdot {}^{2n} C_n$

C.  $(5 + 3n) \cdot {}^{2n} C_n$

D.  $\left(\frac{3n - 5}{2}\right) \cdot {}^{2n} C_{n+1}$

**Answer: B**



**View Text Solution**

45.

$$\frac{x+1}{x^4(x+2)} = \frac{A}{x} + \frac{B}{x^2} + \frac{C}{x^3} + \frac{D}{x^4} + \frac{E}{x+2} \Rightarrow B + D + E =$$

A. A+C

B. A-C

C. 2A+C

D. 2A+2C

Answer: A



Watch Video Solution

46. If  $\cos^3 \theta + \cos^3 \left( \frac{2\pi}{3} + \theta \right) + \cos^3 \left( \frac{4\pi}{3} + \theta \right) = a \cos 3\theta$ ,

then a =

A. 1/4

B.  $\frac{3}{4}$

C.  $\frac{5}{4}$

D.  $\frac{7}{4}$

**Answer: B**



**Watch Video Solution**

47. The solution of the differential equation  $y' = \frac{1}{e^{-y} - x}$ , is

A.  $x = e^{-y}(y + c)$

B.  $y + e^{-y} = x + c$

C.  $x = e^y(y + c)$

D.  $x + y = e^{-y} + c$

**Answer: A**



Watch Video Solution

48. The solution of the differential equation  $3xy' - 3y + (x^2 - y^2)^{1/2} = 0$ , satisfying the condition  $y(1) = 1$  is

A.  $3 \cos^{-1}\left(\frac{y}{x}\right) = \ln|x|$

B.  $3 \cos\left(\frac{y}{x}\right) = \ln|x|$

C.  $3 \cos^{-1}\left(\frac{y}{x}\right) = 2 \ln|x|$

D.  $3 \sin^{-1}\left(\frac{y}{x}\right) = \ln|x|$

Answer: A



Watch Video Solution

49. Let  $p \in \mathbb{R}$ , then the differential equation of the family of curves  $y = (\alpha + \beta x)e^{px}$ , where  $\alpha, \beta$  are arbitrary constants, is

A.  $y'' = 4py' + p^2y = 0$

B.  $y'' - 2py' + p^2y = 0$

C.  $y'' + 2py' - p^2y = 0$

D.  $y'' + 2py' + p^2y = 0$

**Answer: B**



**Watch Video Solution**

50. If the area bounded by the curves  $y = ax^2$  and  $x = ay^2$  ( $a > 0$ ) is 3 sq. units, then the value of 'a' is

A.  $2/3$

B.  $1/3$

C. 1

D. 4

**Answer: B**



**Watch Video Solution**

51.  $\int_0^{\pi/4} [\sqrt{\tan x} + \sqrt{\cot x}] = dx$

A.  $\frac{\pi}{\sqrt{2}}$

B.  $\frac{\pi}{2}$

C.  $\frac{3\pi}{\sqrt{2}}$

D.  $\pi$

**Answer: A**



Watch Video Solution

$$52. \int_0^{\pi/4} \frac{\sin x + \cos x}{7 + 9 \sin 2x} dx =$$

A.  $\frac{\log 3}{4}$

B.  $\frac{\log 3}{36}$

C.  $\frac{\log 7}{12}$

D.  $\frac{\log 7}{24}$

Answer: D



Watch Video Solution

$$53. \int \frac{2x + 2}{\sqrt{x^2 - 4x - 5}} dx =$$



A.  $\sqrt{x^2 - 4x - 5} + \log|x + \sqrt{x^2 - 4x - 5}| + c$

B.  $\log|\sqrt{x^2 - 4x - 5}| + \sqrt{x^2 - 4x - 5} + c$

C.  $\sqrt{x^2 - 4x - 5} + 6 \log|(x - 2) + \sqrt{x^2 - 4x - 5}| + c$

D.  $2\sqrt{x^2 - 4x - 5} + 6 \log|(x - 2) + \sqrt{x^2 - 4x - 5}| + c$

**Answer: D**



**Watch Video Solution**

54.  $\int \frac{dx}{\cos(x + 4)\cos(x + 2)} =$

A.  $\frac{1}{\sin 2} \log |\cos(x + 4)|^2 + c$

B.  $\frac{1}{2} \log \left| \frac{\sec(x + 2)}{\sec(x + 4)} \right| + c$

C.  $\frac{1}{\sin 2} \log \left| \frac{\sec(x + 4)}{\sec(x + 2)} \right| + c$

D.  $\log \left| \frac{\sec(x + 4)}{\sec(x + 2)} \right| + c$

Answer: C



Watch Video Solution

$$55. \int \frac{x^3}{\sqrt{1+x^2}} dx =$$

A.  $\sqrt{1-x^2} - \frac{x}{3}(1+x^2)^{3/2} + c$

B.  $x\sqrt{1+x^2} + \frac{2}{3}(1+x^2)^{3/2} + c$

C.  $\frac{x^2}{3} - \frac{2}{3}(1+x^2)^{3/2} + c$

D.  $x^2\sqrt{1+x^2} - \frac{1}{3}(1+x^2)^{1/2} + c$

Answer: C



Watch Video Solution

56.  $\int \frac{(x^2 + 1)}{x^4 + 7x^2 + 1} dx =$

A.  $\frac{1}{3} \tan^{-1} \left( \frac{x^2 - 1}{3x} \right) + c$

B.  $\tan^{-1} \left( \frac{x^2 - 1}{x} \right) + c$

C.  $\frac{1}{3} \tan^{-1} \left( \frac{x^2 - 1}{x} \right) + c$

D.  $\frac{1}{\sqrt{3}} \tan^{-1} \left( \frac{x^2 - 1}{\sqrt{3}x} \right) + c$

**Answer: A**



**Watch Video Solution**

57. The smallest value of of the constant  $m > 0$  for which

$f(x) = 9mx - 1 + \frac{1}{x} \geq 0$  for all  $x > 0$ , is

A.  $1/9$

B.  $1/16$

C.  $1/36$

D.  $1/81$

**Answer: C**



**Watch Video Solution**

58. Define  $f(x) = \frac{1}{2}[|\sin x| + \sin x]$ ,  $0 < x \leq 2\pi$ . Then  $f$  is

A. increasing in  $\left(\frac{\pi}{2}, \frac{3\pi}{2}\right)$

B. decreasing in  $\left(0, \frac{\pi}{2}\right)$  and increasing in  $\left(\frac{\pi}{2}, \pi\right)$

C. increasing in  $\left(0, \frac{\pi}{2}\right)$  and decreasing in  $\left(\frac{\pi}{2}, \pi\right)$

D. increasing in  $\left(0, \frac{\pi}{4}\right)$  and decreasing in  $\left(\frac{\pi}{4}, \pi\right)$

**Answer: C**



Watch Video Solution

59. The area of the triangle formed by the positive x-axis, the tangent and normal to the curve  $x^2 + y^2 = 16a^2$  at the point  $(2\sqrt{2}a, 2\sqrt{2}a)$  is

A.  $a^2$

B.  $16a^2$

C.  $4a^2$

D.  $8a^2$

**Answer: D**



Watch Video Solution

60. The length of the segment of the tangent line curve  $x = a \cos^3 t$ ,  $y = a \sin^3 t$ , at any point on the curve cut off by the coordinate axes is

A.  $4a$

B.  $a$

C.  $a^2$

D.  $2a$

**Answer: B**



[View Text Solution](#)

61. If  $y = a \cos(\sin x) + b \sin(\sin x)$  then  $y_2 + (\tan x)y_1 =$

A. 0

B.  $4(\cos^2 2x)y$

C.  $-4(\cos^2 2x)y$

D.  $-(\cos^2 2x)y$

**Answer: C**



**Watch Video Solution**

62.  $\frac{d}{dx} \tan^{-1} \left[ \frac{\sqrt{1 + \sin x} - \sqrt{1 - \sin x}}{\sqrt{1 + \sin x} + \sqrt{1 - \sin x}} \right] =$

A. 1

B.  $-1/2$

C.  $1/2$

D.  $-1$

**Answer: b**



**Watch Video Solution**

63. If  $y = \tan^{-1} \left[ \frac{5 \cos x - 12 \sin x}{12 \cos x + 5 \sin x} \right]$ , then  $\frac{dy}{dx} =$

A. 1

B. -1

C. -2

D. 1/2

**Answer: B**



**Watch Video Solution**



64. continuous at  $x = 0$

$$\text{Let } f(x) = \begin{cases} (1 + |\sin x|)^{\frac{a}{|\sin x|}} & -\frac{\pi}{6} < x < 0 \\ b & x = 0 \\ e^{\tan 2x / \tan 3x} & 0 < x < \frac{\pi}{6} \end{cases}$$

Determine a and b such that  $f(x)$  continuous at  $x = 0$ .

A.  $p = \frac{1}{3}, q = e^{2/3}$

B.  $p = 0, q = e^{2/3}$

C.  $p = \frac{2}{3}, q = e^{-2/3}$

D.  $p = -\frac{2}{3}, q = e^{2/3}$

Answer: D



Watch Video Solution

65.  $\lim_{x \rightarrow \infty} \left[ \frac{x^2 + x + 3}{x^2 - x + 2} \right]^x =$

A.  $\infty$

B.  $e$

C.  $e^4$

D.  $e^2$

**Answer: D**



**Watch Video Solution**

**66.** The image of the point  $(5, 2, 6)$  with respect to the plane  $x + y + z = 9$  is

A.  $(3, -5, 2)$

B.  $\left(\frac{7}{2}, -1, 5\right)$

C.  $\left(\frac{7}{3}, -\frac{2}{3}, \frac{10}{3}\right)$

D.  $\left(\frac{7}{3}, \frac{2}{3}, -\frac{5}{3}\right)$

**Answer: C**



**Watch Video Solution**

**67.** If the angle between the lines whose direction cosines are

$$\left(-\frac{2}{\sqrt{21}}, \frac{C}{\sqrt{21}}, \frac{1}{\sqrt{21}}\right) \text{ and } \left(\frac{3}{\sqrt{54}}, \frac{3}{\sqrt{54}}, -\frac{6}{\sqrt{54}}\right) \text{ is } \frac{\pi}{2},$$

then the value of C is

A. 6

B. 4

C. -4

D. 2

**Answer: B**



Watch Video Solution

68. Points  $A(3, 2, 4)$ ,  $B\left(\frac{33}{5}, \frac{28}{5}, \frac{38}{5}\right)$ , and  $C(9, 8, 10)$  are given.

The ratio in which B divides  $\overline{AC}$  is

A. 5 : 3

B. 2 : 1

C. 1 : 3

D. 3 : 2

**Answer: D**



Watch Video Solution

69.

Let

A

$(2 \sec \theta, 3 \tan \theta)$  and  $B(2 \sec \phi, 3 \tan \phi)$  where  $\theta + \phi = \frac{\pi}{2}$  be

two point on the hyperbola  $\frac{x^2}{4} - \frac{y^2}{9} = 1$ . If  $(\alpha, \beta)$  is the

point of intersection of normals to the hyperbola at A and B

, then  $\beta =$

A.  $\frac{-13}{3}$

B.  $\frac{13}{3}$

C.  $\frac{3}{13}$

D.  $\frac{-3}{13}$

**Answer: A**



**Watch Video Solution**

70. If S and S'' are the foci of the ellipse  $\frac{x^2}{25} + \frac{y^2}{16} = 1$  and if PSP' is a focal chord with SP = 8 then SS'' =

A.  $4 + S'P$

B.  $S'P - 1$

C.  $4 + SP$

D.  $SP - 1$

**Answer: A**



**Watch Video Solution**

71. For the ellipse given by  $\frac{(x - 3)^2}{25} + \frac{(y - 2)^2}{16} = 1$ , match the equations of the lines given in List - I with those in the List -

II

**List - I**

- i) The equation of the major axis
- ii) The equation of a directrix
- iii) The equation of a latus rectum

**List - II**

- a)  $3x = 34$
- b)  $y = 2$
- c)  $x + y = 9$
- d)  $x = 6$
- e)  $x = 3$
- f)  $3y = 34$

The correct matching is

- A. 1-e, ii-a, iii-d,
- B. i-b, ii-f, iii-e
- C. i-b, ii-a, iii-e
- D. i-b,ii-a, iii-d

**Answer: A**



**Watch Video Solution**

72. The points of intersection of the parabolas  $y^2 = 5x$  and  $x^2 = 5y$  lie on the line

A.  $x + y = 10$

B.  $x - 2y = 0$

C.  $x - y = 0$

D.  $2x - y = 0$

**Answer: C**



**Watch Video Solution**

73. From a point  $(C, 0)$  three normals are drawn to the parabola  $y^2 = x$ . Then

A.  $C < \frac{1}{2}$



$$\text{B. } C = \frac{1}{2}$$

$$\text{C. } C > \frac{1}{2}$$

$$\text{D. } \frac{1}{2} > C > \frac{1}{4}$$

**Answer: C**



**Watch Video Solution**

**74.** The radical centre of the circles

$$x^2 + y^2 = 1, x^2 + y^2 - 2x - 3 = 0 \quad \text{and}$$

$$x^2 + y^2 - 2y - 3 = 0 \text{ is}$$

A. (1,1)

B. (1,-1)

C. (-1,1)

D. (-1, -1)

**Answer: D**



[Watch Video Solution](#)

75. If the circles  $x^2 + y^2 - 2\lambda x - 2y - 7 = 0$  and  $3(x^2 + y^2) - 8x + 29y = 0$  are orthogonal then  $\lambda =$

A. 4

B. 3

C. 2

D. 1

**Answer: D**



[Watch Video Solution](#)

76. For all real values of  $k$ , the polar of the point  $(2k, k-4)$  with respect to  $x^2 + y^2 - 4x - 6y + 1 = 0$  passes through the point

- A. (1, 1)
- B. (1, -1)
- C. (-3, 1)
- D. (3, 1)

**Answer: D**



[Watch Video Solution](#)

77. The normal to the circle given by  $x^2 + y^2 - 6x + 8y - 144 = 0$  at  $(8, 8)$  meets the circle again

at the point

A. (2, -16)

B. (2, 16)

C. (-2, 16)

D. (-2, -16)

**Answer: D**



**Watch Video Solution**

**78.** The point where the line  $4x - 3y + 7 = 0$  touches the circle

$$x^2 + y^2 - 6x + 4y - 12 = 0 \text{ is}$$

A. (1, 1)

B. (1, -1)

C. (-1, 1)

D. (-1, -1)

**Answer: C**



**Watch Video Solution**

**79.** The combined equation of three sides of a triangle is  $(x^2 - y^2)(2x + 3y - 6) = 0$ . If  $(-2, \alpha)$  is an interior point of the triangle, then

A.  $-2 < \alpha < 0$

B.  $-2 < \alpha < 2$

C.  $0 < \alpha < 2$

D.  $\alpha \geq 2$

**Answer: C**



**Watch Video Solution**

**80.** The equation on the pair of straight lines through the point (1, 1) and perpendicular to the pair of straight lines  $3x^2 - 8xy + 5y^2 = 0$  is

A.  $5x^2 + 8xy + 3y^2 - 14x - 18y + 16 = 0$

B.  $5x^2 + 8xy + 3y^2 - 18x - 14y + 16 = 0$

C.  $5x^2 - 8xy + 3y^2 - 18x - 14y + 32 = 0$

D.  $5x^2 - 8xy + 3y^2 - 14x - 18y + 32 = 0$

**Answer: B**



**Watch Video Solution**

