



India's Number 1 Education App

MATHS

BOOKS - DEEPTI MATHS (TELUGU ENGLISH)

LIMITS AND CONTINUITY

Exercise 1A

$$1. \lim_{x \rightarrow -2} \frac{x^4 + 2x^3 + 3x^2 + 5x - 2}{x^5 + 3x^4 + 2x^3 + 3x^2 + 7x + 2} =$$

A. 5

B. -5

C. 8

D. 16

Answer: B



doubtnut | doubt, not out



Watch Video Solution

2. $\lim_{x \rightarrow -2} \frac{(1+x)^2(1-x)^2}{(1+x)^3 - (1-x)^3} =$

A. 1

B. $1/2$

C. $2/3$

D. $3/2$

Answer: C



View Text Solution

3. $\lim_{x \rightarrow 2} \frac{x^5 - 32}{x^3 - 8} =$

A. $20/3$

B. $3/2$

C. $\sqrt{3}/5$

D. $8/9$

Answer: A



Watch Video Solution

4. $\lim_{x \rightarrow 1} \frac{x^{-2/3} - 1}{x^{-3/4} - 1} =$

A. $20/3$

B. $3/2$

C. $\sqrt{3}/5$

D. $8/9$

Answer: D



Watch Video Solution

5. If $f(x) = \frac{4 - 7x}{3x + 4}$ and $\lim_{x \rightarrow 2} f(x) = m$ then the equation whose roots are $1, 1/m$ is

A. $(x - 1)^2 = 0$

B. $(x + 1)^2 = 0$

C. $x^2 - 1 = 0$

D. none

Answer: C



Watch Video Solution

6. $\lim_{x \rightarrow 1} \frac{(2x - 3)(\sqrt{x} - 1)}{2x^2 + x - 3} =$

A. 0

B. 1

C. 2

D. $-\frac{1}{10}$

Answer: D



Watch Video Solution

7. Evaluate $\lim_{x \rightarrow 0} \frac{\sqrt{1+x} - 1}{x}$

A. 0

B. 1

C. 2

D. 1/2

Answer: D



Watch Video Solution

8. $\lim_{x \rightarrow 0} \frac{\sqrt[n]{1+x} - 1}{x} =$

A. 0

B. n

C. $1/n$

D. $1/\sqrt{n}$

Answer: C



Watch Video Solution

$$9. \lim_{x \rightarrow 0} \frac{\sqrt[3]{x+8}-2}{x} =$$

A. 0

B. 1

C. $1/12$

D. $-1/12$

Answer: C



Watch Video Solution

$$10. \lim_{x \rightarrow 0} \frac{\sqrt{1+x+x^2}}{x} =$$

A. 0

B. 1

C. 2

D. $1/2$

Answer: D



Watch Video Solution

$$11. \lim_{x \rightarrow a} \frac{\sqrt{(x+a)} - \sqrt{2a}}{x-a} =$$

A. $2a$

B. $2\sqrt{2a}$

C. $1/2\sqrt{2a}$

D. $1/\sqrt{2}a$

Answer: C



Watch Video Solution

12. $\lim_{x \rightarrow 0} \frac{\sqrt{1+x} - \sqrt{1-x}}{x} =$

A. 0

B. 1

C. $\sqrt{2}$

D. $1/\sqrt{2}$

Answer: B



Watch Video Solution

13. $\lim_{x \rightarrow 0} \frac{\sqrt{a+x^2} - \sqrt{a-x^2}}{x^2} =$

A. 0

B. 1

C. \sqrt{a}

D. $1 / \sqrt{a}$

Answer: D



Watch Video Solution

$$14. \lim_{x \rightarrow 0} \frac{\sqrt{1+x^3} - \sqrt{1-x^3}}{x^3} =$$

A. 0

B. 1

C. 2

D. $1/2$

Answer: B



Watch Video Solution

$$15. \lim_{x \rightarrow 0} \frac{\sqrt[3]{1+x} - \sqrt[3]{1-x}}{x} =$$

A. 1

B. $1/2$

C. $2/3$

D. $3/2$

Answer: C



Watch Video Solution

$$16. \lim_{x \rightarrow 0} \frac{(1+x)^{1/4} - (1-x)^{1/4}}{x} =$$

A. $1/2$

B. 0

C. -1

D. $-1/2$

Answer: A



Watch Video Solution

17. $\lim_{x \rightarrow 0} \frac{x}{\sqrt{x+4} - 2} =$

A. 4

B. $\sqrt{2}$

C. $2\sqrt{2}$

D. $\frac{1}{\sqrt{2}}$

Answer: A



Watch Video Solution

18. $\lim_{x \rightarrow 1} \frac{x-1}{\sqrt{x^2+3} - 2} =$

A. 2

B. 3

C. - 2

D. - 3

Answer: A



Watch Video Solution

19. $\lim_{x \rightarrow 0} \frac{x}{1 - \sqrt{1-x}} =$

A. 2

B. 3

C. - 2

D. - 3

Answer: A



Watch Video Solution

$$20. \lim_{x \rightarrow 0} \frac{x^2}{1 - \sqrt[3]{1 - x^2}} =$$

A. 2

B. 3

C. -2

D. -3

Answer: B



Watch Video Solution

$$21. \lim_{x \rightarrow -1} \frac{x + 1}{2 - \sqrt{4 + x + x^2}} =$$

A. 1/4

B. 4

C. 1/2

D. 2

Answer: B



Watch Video Solution

$$22. \lim_{x \rightarrow 0} \frac{x}{\sqrt{4-x} - \sqrt{4+x}} =$$

A. 2

B. 3

C. - 2

D. - 3

Answer: C



Watch Video Solution

$$23. \lim_{x \rightarrow -8} \frac{\sqrt{1+\sqrt{1+x}} - 2}{x - 8} =$$

A. $\frac{3}{2}$

B. $\frac{1}{4}$

C. $\frac{1}{24}$

D. ∞

Answer: C



Watch Video Solution

24. $\lim_{x \rightarrow a} \frac{\sqrt{a+2x} - \sqrt{3x}}{\sqrt{1+x^2} - \sqrt{1+x}} =$

A. $\frac{2}{\sqrt{3}}$

B. $\frac{1}{\sqrt{3}}$

C. $\frac{-2}{3\sqrt{3}}$

D. $\frac{2}{3\sqrt{3}}$

Answer: D



View Text Solution

$$25. \lim_{x \rightarrow 0} \frac{\sqrt{1-x} - \sqrt{1-x^2}}{\sqrt{1+x^2} - \sqrt{1+x}} =$$

A. 1

B. -1

C. 0

D. 2

Answer: A



Watch Video Solution

$$26. \lim_{x \rightarrow 0} \frac{\sqrt{4+x} - \sqrt{4+3x}}{x} =$$

A. -1/2

B. 1/2

C. -3

D. 0

Answer: A



Watch Video Solution

$$27. \text{ Lt}_{x \rightarrow 1} \frac{\sqrt[3]{x^2} - 2\sqrt[3]{x} + 1}{(x - 1)^2} =$$

A. $1/6$

B. $1/9$

C. $-1/6$

D. $-1/9$

Answer: B



Watch Video Solution

$$28. \text{ Lt}_{x \rightarrow 2} \left[\left(\frac{x^3 - 4x}{x^3 - 8} \right)^{-1} - \left(\frac{x + \sqrt{2}x}{x - 2} - \frac{\sqrt{2}}{\sqrt{x} - \sqrt{2}} \right)^{-1} \right]$$

A. $1/2$

B. 2

C. 1

D. none

Answer: A



Watch Video Solution

$$29. \lim_{x \rightarrow 0} \frac{5x - \sin 5x}{x} =$$

A. 0

B. 1

C. 2

D. $1/2$

Answer: A



Watch Video Solution

30. $\lim_{x \rightarrow 0} \frac{5x - \sin 5x}{x} =$

A. ab/c

B. $a^2 b^3 / c^4$

C. c/ab

D. $c^4 / a^2 b^3$

Answer: B



Watch Video Solution

31. $\lim_{x \rightarrow 0} \frac{x \tan 7x}{\sin 3x \sin 4x} =$

A. 0

B. 1

C. 2/9

Answer: D**Watch Video Solution**

$$32. \lim_{x \rightarrow 0} \frac{\tan 6x \cdot \tan 5x}{\sin 3x \sin 4x} =$$

A. 5

B. $5/2$ C. $3/7$ D. $5/3$ **Answer: B****Watch Video Solution**

$$33. \lim_{x \rightarrow 0} \frac{3 \sin x - \sin 3x}{x^3} =$$

A. 4

B. - 4

C. 8

D. - 8

Answer: A



Watch Video Solution

$$34. \lim_{\theta \rightarrow 0} \frac{3 \tan \theta - \tan 3\theta}{2\theta^3}$$

A. $1/4$

B. $3/4$

C. 4

D. - 4

Answer: D



Watch Video Solution

$$35. \lim_{\theta \rightarrow 0} \frac{a \sin ax - b \sin bx}{\tan ax - \tan bx} =$$

A. $a^2 - b^2$

B. 0

C. a+b

D. a-b

Answer: C



Watch Video Solution

$$36. \lim_{x \rightarrow 0} \frac{\sin 5x - \sin 3x}{\sin 4x - \sin 2x} =$$

A. 0

B. 1

C. 2/9

Answer: B**Watch Video Solution**

$$37. \lim_{x \rightarrow 0} \frac{\sin 7x - \sin 5x + \sin 3x - \sin x}{\sin 6x - \sin 4x + \sin 2x} =$$

A. -1

B. 0

C. 1

D. 2

Answer: C**Watch Video Solution**

$$38. \lim_{x \rightarrow 0} \frac{1 - \cos x}{x} =$$

A. 0

B. 1

C. 2

D. $1/2$

Answer: A



Watch Video Solution

$$39. \text{ Lt}_{x \rightarrow 0} \frac{1 - \cos x}{x^2} =$$

A. 0

B. 1

C. 2

D. $1/2$

Answer: D



Watch Video Solution

$$40. \lim_{x \rightarrow 0} \frac{1 - \cos x}{x \sin 2x} =$$

A. $1/2$

B. $3/2$

C. $3/4$

D. $1/4$

Answer: D



Watch Video Solution

$$41. \lim_{x \rightarrow 0} \frac{1 - \cos mx}{1 - \cos nx} =$$

A. m/n

B. n/m

C. m^2 / n^2

D. n^2 / m^2

Answer: C



Watch Video Solution

42. $\lim_{x \rightarrow 0} \frac{\operatorname{cosec} x - \cot x}{x} =$

A. 0

B. 1

C. 2

D. $1/2$

Answer: D



Watch Video Solution

43. $\lim_{x \rightarrow 0} \frac{\tan x - \sin x}{x^3} =$

A. 0

B. 1

C. 2

D. $1/2$

Answer: D



Watch Video Solution

$$44. \lim_{x \rightarrow 0} \frac{\tan x - \sin x}{x^2} =$$

A. 0

B. 1

C. $\frac{1}{2}$

D. $-\frac{1}{2}$

Answer: A



Watch Video Solution

$$45. \lim_{x \rightarrow 0} \frac{\cos ax - \cos bx}{x^2} =$$

A. 0

B. 1

C. $\frac{a^2 - b^2}{2}$

D. $\frac{b^2 - a^2}{2}$

Answer: D



Watch Video Solution

$$46. \lim_{x \rightarrow 0} \frac{\cos 5x - \cos 3x}{x^2} =$$

A. 4

B. -4

C. 8

Answer: D**Watch Video Solution**

$$47. \lim_{x \rightarrow 0} \frac{\cos 7x - \cos 9x}{\cos 3x - \cos 7x} =$$

A. $5/2$ B. $2/5$ C. $7/5$ D. $4/5$ **Answer: D****Watch Video Solution**

$$48. \lim_{x \rightarrow 0} \frac{1 - 2\cos x + \cos 2x}{x^2} =$$

A. 0

B. 1

C. -1

D. none

Answer: C



Watch Video Solution

49. $\lim_{x \rightarrow 0} \frac{(1 - \cos 2x)(3 + \cos x)}{x \tan 4x}$ is equal to

A. 1

B. 2

C. -1/4

D. 1/2

Answer: B



Watch Video Solution

50. $\lim_{x \rightarrow 0} \frac{1 - \cos^3 x}{x \sin x \cos x} =$

A. $2/5$

B. $3/5$

C. $3/2$

D. $3/4$

Answer: C



[Watch Video Solution](#)

51. $\lim_{x \rightarrow 0} \frac{\sin(\pi \cos^2 x)}{x^2}$ is equal to

A. $-\pi$

B. π

C. $\frac{\pi}{2}$

D. 1

Answer: B



Watch Video Solution

52. $\lim_{x \rightarrow 0} \frac{\tan^3 x - \sin^3 x}{x^5} =$

A. $\frac{5}{2}$

B. $\frac{3}{2}$

C. $\frac{3}{5}$

D. $\frac{2}{5}$

Answer: B



View Text Solution

53. The quadratic equation whose roots are l and m where

$$l = \lim_{\theta \rightarrow 0} \left(\frac{3 \sin \theta - 4 \sin^2 \theta}{\theta} \right) \text{ and } m = \lim_{\theta \rightarrow 0} \frac{2 \tan \theta}{\theta(1 - \tan^2 \theta)} \text{ is}$$

A. $x^2 + 5x + 6 = 0$

B. $x^2 - 5x + 6 = 0$

C. $x^2 - 5x - 6 = 0$

D. $x^2 + 5x - 6 = 0$

Answer: B



Watch Video Solution

54. $\lim_{x \rightarrow 0} \frac{3 \sin x^\circ - \sin 3x^\circ}{x^3} =$

A. 0

B. 1

C. $\left(\frac{\pi}{180}\right)^3$

$$\text{D. } 4. \left(\frac{\pi}{180} \right)^3$$

Answer: D



Watch Video Solution

$$55. \lim_{x \rightarrow 0} \frac{1 - \cos 2x^\circ}{x^2} =$$

A. 0

B. 1

$$\text{C. } \left(\frac{\pi}{180} \right)^2$$

$$\text{D. } 2. \left(\frac{\pi}{180} \right)^2$$

Answer: D



Watch Video Solution

$$56. \lim_{x \rightarrow \pi/2} \frac{\cos x}{\pi/2 - x} =$$

A. 0

B. -1

C. 1

D. $1/2$

Answer: C



Watch Video Solution

$$57. \lim_{x \rightarrow \pi/2} \frac{\cot x}{x - \pi/2} =$$

A. 0

B. -1

C. 1

D. $1/2$

Answer: B



Watch Video Solution

$$58. \lim_{x \rightarrow \pi/2} \frac{(\pi/2 - x) \sec x}{\operatorname{cosec} x} =$$

A. 0

B. -1

C. 1

D. 1/2

Answer: C



[Watch Video Solution](#)

$$59. \lim_{x \rightarrow \pi/2} \frac{\sec x - \tan x}{\pi/2 - x} =$$

A. 0

B. 1

C. 2

D. $1/2$

Answer: B



[View Text Solution](#)

60. $\lim_{x \rightarrow \pi/2} \sec 5x \cos 7x =$

A. $5/7$

B. $-5/7$

C. $7/5$

D. $-7/5$

Answer: D



[Watch Video Solution](#)

61. $\lim_{x \rightarrow \pi/2} \frac{\cot x - \cos x}{(\pi/2 - x)^3} =$

A. 0

B. 1

C. 2

D. $1/2$

Answer: D



Watch Video Solution

$$62. \lim_{\theta \rightarrow \pi/2} \frac{1 - \sin^2 \theta}{\cos \theta (\pi/2 - \theta)} =$$

A. 1

B. -1

C. $-1/2$

D. $1/2$

Answer: A



Watch Video Solution

$$63. \lim_{\theta \rightarrow \pi/2} \frac{[1 - \tan(x/2)][1 - \sin x]}{[1 + \tan(x/2)][\pi - 2x]^3} =$$

A. $1/8$

B. 0

C. $1/32$

D. ∞

Answer: C



[View Text Solution](#)

$$64. \lim_{x \rightarrow \pi/2} \frac{1 - \sin^3 x}{\cos^2 x} =$$

A. $3/2$

B. $2/3$

C. $2/5$

D. $-3/2$

Answer: A



Watch Video Solution

$$65. \lim_{x \rightarrow 3\pi/2} \frac{\cos^2 x}{1 + \sin^3 x} =$$

A. $3/2$

B. $2/3$

C. $2/5$

D. $-3/2$

Answer: B



Watch Video Solution

$$66. \lim_{x \rightarrow 1} \frac{\tan^2 \pi x}{1 + \sec \pi x} =$$

A. 0

B. 1

C. 2

D. -2

Answer: D



Watch Video Solution

$$67. \lim_{x \rightarrow 0} \frac{\sec x - 1}{x^2(\sec x + 1)^2} =$$

A. 1/8

B. 11/4

C. 2

D. 0

Answer: A



Watch Video Solution

68. $\lim_{x \rightarrow 1} (1 - x) \tan(\pi x / 2) =$

A. $2/\pi$

B. $3/\pi$

C. $4/\pi$

D. none

Answer: A



[Watch Video Solution](#)

69. $\lim_{x \rightarrow a} \frac{\operatorname{cosec}(x - a)}{x - a} =$

A. 1

B. $-\infty$

C. -1

D. does not exist

Answer: B



Watch Video Solution

$$70. \lim_{x \rightarrow a} \frac{\tan x - \tan a}{\sin a - \sin x} =$$

A. $\sec a$

B. $-\sec^2 a$

C. $\sec^3 a$

D. $-\sec^3 a$

Answer: D



Watch Video Solution

$$71. \lim_{y \rightarrow x} \frac{\sin^2 y - \sin^2 x}{y - x} =$$

A. $\sin 2y$

B. $\sin 2x$

C. $\cos^2 y$

D. $\cos^2 x$

Answer: B



Watch Video Solution

$$72. \lim_{x \rightarrow 1} \frac{3 \sin \pi x - \sin 3\pi x}{(x - 1)^3} =$$

A. $-4\pi^3$

B. 0

C. 1

D. x^3

Answer: A



Watch Video Solution

$$73. \lim_{x \rightarrow \pi/6} \left(\frac{3 \sin x - \sqrt{3} \cos x}{6x - \pi} \right) =$$

A. $\sqrt{3}$

B. $1/\sqrt{3}$

C. $-\sqrt{3}$

D. $-1/\sqrt{3}$

Answer: B



Watch Video Solution

$$74. \lim_{x \rightarrow \pi/4} \frac{\sec x \tan(4x - \pi)}{\sin(\pi - 4x)} =$$

A. 2

B. $\sqrt{2}$

C. $-\sqrt{2}$

D. $2\sqrt{2}$

Answer: C



Watch Video Solution

$$75. \lim_{x \rightarrow \pi/4} \frac{1 - \cot^3 x}{2 - \cot x - \cot^3 x} =$$

A. $11/4$

B. $3/4$

C. $1/2$

D. none

Answer: B



Watch Video Solution

$$76. \lim_{x \rightarrow -2} \frac{\tan \pi x}{x + 2} + \lim_{x \rightarrow \infty} (1 + 1/x^2) =$$

A. 3

B. π

C. $\pi + 1$

D. $\pi + 3$

Answer: C



Watch Video Solution

$$77. \lim_{x \rightarrow 0} \frac{1 - \cos(1 - \cos x)}{x^4} =$$

A. $1/2$

B. $1/4$

C. $1/8$

D. $1/12$

Answer: C



Watch Video Solution

78. Let α and β be the distinct roots of $ax^2 + bx + c = 0$, then $\lim_{x \rightarrow a} \frac{1 - \cos(ax^2 + bx + c)}{(x - \alpha)^2}$

A. $\frac{a^2}{2}(\alpha - \beta)^2$

B. 0

C. $-\frac{a^2}{2}(\alpha - \beta)^2$

D. $\frac{1}{2}(\alpha - \beta)^2$

Answer: A



Watch Video Solution

79. $\lim_{x \rightarrow 0} \frac{\cos^2 x - \sin^2 x - 1}{\sqrt{x^2 + 1} - 1} =$

A. 0

B. -4

C. 1

D. 4

Answer: B



Watch Video Solution

$$80. \lim_{x \rightarrow 0} \frac{\sqrt{1 + \sin x} - \sqrt{1 - \sin x}}{x} =$$

A. 0

B. -1

C. 2

D. 1

Answer: D



Watch Video Solution

$$81. \lim_{x \rightarrow 0} \frac{\sqrt{1 + \tan x} - \sqrt{1 - \tan x}}{\sin x} =$$

A. 0

B. 1

C. 2

D. $1/2$

Answer: B



Watch Video Solution

$$82. \lim_{x \rightarrow 0} \frac{\sqrt{1 + \sin x + \sin^2 x} - 1}{x} =$$

A. 0

B. 1

C. 2

D. $1/2$

Answer: D



Watch Video Solution

$$83. \lim_{x \rightarrow \pi/2} \frac{\sqrt{1 + \cos^3 x} - \sqrt{1 - \cos^3 x}}{(\pi/2 - x)^3} =$$

A. 0

B. 1

C. $\pi/2$

D. $\pi/4$

Answer: B



View Text Solution

$$84. \lim_{x \rightarrow \pi/2} \frac{\sqrt{1 - \sin x}}{(\pi/2 - x)\sqrt{1 + \sin x}} =$$

A. $1/2$

B. $\pi / 2$

C. 0

D. $-1 / 2$

Answer: A



[View Text Solution](#)

$$85. \lim_{x \rightarrow 0} \frac{\sin^{-1} x}{x} =$$

A. 0

B. 1

C. -1

D. $1 / 2$

Answer: B



[Watch Video Solution](#)

$$86. \lim_{x \rightarrow 0} \frac{\tan^{-1} x}{x} =$$

A. 0

B. 1

C. -1

D. 1/2

Answer: B



[Watch Video Solution](#)

$$87. \lim_{x \rightarrow 0} \frac{\sin^{-1} x - \tan^{-1} x}{x} =$$

A. 2

B. 1

C. -1

D. 1/2

Answer: D



View Text Solution

$$88. \lim_{x \rightarrow -1} \frac{\sqrt{x} - \sqrt{\cos^{-1} x}}{\sqrt{x} + 1} =$$

A. $1/\sqrt{2}$

B. $1/\sqrt{2x}$

C. $1/\sqrt{x}$

D. none

Answer: B



View Text Solution

$$89. \lim_{x \rightarrow 0} \frac{\sqrt[3]{1 + \tan^{-1} 3x} - \sqrt[3]{1 - \sin^{-1} 3x}}{\sqrt{1 - \sin^{-2} 2x} - \sqrt{1 + \tan^{-1} 2x}}$$

A. -1

B. 0

C. 1

D. 2

Answer: A



[View Text Solution](#)

$$90. \lim_{x \rightarrow 0} \frac{\sin^{-1} ax - \sin^{-1} bx}{x} =$$

A. 0

B. 1

C. a-b

D. none

Answer: C



[Watch Video Solution](#)

$$91. \lim_{x \rightarrow 0} \frac{x(1 - \sqrt{1 - x^2})}{\sqrt{1 - x^2} \cdot (\sin^{-1} x)^3} =$$

- A. 1
- B. -1
- C. 1/2
- D. 2

Answer: C



Watch Video Solution

$$92. \lim_{x \rightarrow 0} \frac{\sinh x}{x} =$$

- A. 0
- B. ∞
- C. $-\infty$
- D. 1

Answer: D



Watch Video Solution

$$93. \lim_{x \rightarrow 0} \frac{\tan hx}{x} =$$

A. 0

B. ∞

C. $-\infty$

D. 1

Answer: D



Watch Video Solution

$$94. \lim_{x \rightarrow 0} \frac{\sqrt[5]{2+x} - \sqrt[5]{2-x}}{\sin hx} =$$

A. 1

B. $\sqrt[5]{2}$

C. $\sqrt{5}/2$

D. $\sqrt[5]{2}/5$

Answer: D



Watch Video Solution

95. $\lim_{x \rightarrow 1} \frac{1 + \log x - x}{1 - 2x + x^2} =$

A. 1

B. 0

C. -1

D. -1/2

Answer: D



Watch Video Solution

$$96. \lim_{x \rightarrow 1} \left\{ \frac{1}{x-1} - \frac{2}{x^2-1} \right\} =$$

A. $1/2$

B. $1/8$

C. $7/2$

D. $1/32$

Answer: A



Watch Video Solution

$$97. \lim_{x \rightarrow 1} \frac{1}{x-1} \left\{ \frac{1}{x+3} - \frac{2}{3x+5} \right\} =$$

A. $1/2$

B. $1/8$

C. $7/2$

D. $1/32$

Answer: D



Watch Video Solution

$$98. \lim_{x \rightarrow 2} \left\{ \frac{1}{x^2 + x - 6} - \frac{1}{2x^2 - 3x - 2} \right\} =$$

A. $1/15$

B. $1/25$

C. $1/30$

D. $1/40$

Answer: B



Watch Video Solution

$$99. \lim_{x \rightarrow 1/2} \left\{ \frac{8x - 3}{2x - 1} - \frac{4x^2 + 1}{4x^2 - 1} \right\} =$$

A. $1/2$

B. $1/8$

C. $7/2$

D. $1/32$

Answer: C



Watch Video Solution

$$100. \lim_{x \rightarrow 0} \frac{1}{\sin^2 x} - \frac{1}{\sinh^2 x} =$$

A. $2/3$

B. 0

C. $1/3$

D. $-2/3$

Answer: A



View Text Solution

$$101. \lim_{x \rightarrow \infty} \frac{(x+1)(2x+3)}{(x+2)(3x+4)} =$$

A. $7/2$

B. $-5/4$

C. $2/3$

D. $2/7$

Answer: C



Watch Video Solution

$$102. \lim_{x \rightarrow \infty} \frac{(x+1)(2x+3)}{(x+2)(3x+4)} =$$

A. 0

B. ∞

C. 1

D. a

Answer: C



Watch Video Solution

103. $\lim_{x \rightarrow \infty} \frac{2x + 7}{x^2 + 5x + 4} =$

A. 0

B. 1

C. 2

D. $1/2$

Answer: A



Watch Video Solution

104. $\lim_{x \rightarrow \infty} \frac{3x^2 + 5x + 2}{2x^2 - 3x + 1} =$

A. $2/3$

B. $3/2$

C. 3

D. $1/2$

Answer: B



Watch Video Solution

$$105. \lim_{x \rightarrow \infty} \frac{3x^3 + x^2 - 1}{x^2 - x + 7} =$$

A. 0

B. 1

C. ∞

D. $-\infty$

Answer: C



Watch Video Solution

$$106. \lim_{x \rightarrow \infty} \frac{-x^3 + 8}{2x^2 + 5x + 7} =$$

A. 0

B. 1

C. ∞

D. $-\infty$

Answer: D



Watch Video Solution

$$107. \lim_{x \rightarrow \infty} \frac{|3x^2 + 1|}{2x^2 + 1} =$$

A. $3/2$

B. $2/3$

C. $-3/2$

D. $-2/3$

Answer: A



Watch Video Solution

$$108. \lim_{x \rightarrow \infty} \frac{\sqrt{3x^2 + 5}}{5x + 3} =$$

A. $20/3$

B. $3/2$

C. $\sqrt{3}/5$

D. $8/9$

Answer: C



Watch Video Solution

$$109. \lim_{x \rightarrow 0} 2(\sqrt{x}) =$$

A. 0

B. ∞

C. $1/2$

D. $1/\sqrt{2}$

Answer: A



Watch Video Solution

$$110. \lim_{x \rightarrow \infty} \frac{\sqrt{1 + 25x^2} + \sqrt{9x^2 - 1}}{\sqrt{1 + 25x^2} - \sqrt{9x^2 - 1}}$$

A. 1

B. 2

C. 3

D. 4

Answer: D



Watch Video Solution

$$111. \lim_{x \rightarrow \infty} \frac{\sqrt{1+9x^2} + \sqrt{x^2-1}}{\sqrt{1+9x^2} - \sqrt{x^2-1}} =$$

A. 1

B. 2

C. 3

D. 4

Answer: B



[Watch Video Solution](#)

$$112. \lim_{x \rightarrow \infty} \left(\frac{\sqrt{x^2+1} - \sqrt[3]{x^2+1}}{\sqrt[4]{x^4+1} - \sqrt[5]{x^4+1}} \right) =$$

A. 0

B. 1

C. $1/2$

D. $1/4$

Answer: B



Watch Video Solution

$$113. \lim_{x \rightarrow \infty} \frac{(x+1)^{10} + (x+2)^{10} + \dots + (x+10)^{10}}{x^{10} + 10^{10}} = \dots$$

A. 0

B. 1

C. -1

D. 10

Answer: D



Watch Video Solution

$$114. \lim_{x \rightarrow \infty} \left(\frac{x^3}{3x^2 - 1} - \frac{x^2}{3x + 1} \right) =$$

A. 0

B. $1/9$

C. ∞

D. none

Answer: B



Watch Video Solution

$$115. \lim_{x \rightarrow \infty} \left\{ \sqrt{x^2 + ax + b} - x \right\} =$$

A. 0

B. a

C. $a/2$

D. $-a/2$

Answer: B



Watch Video Solution

$$116. \lim_{x \rightarrow \infty} \left\{ x - \sqrt{x^2 + x} \right\} =$$

A. 0

B. 1

C. $1/2$

D. $-1/2$

Answer: D



[Watch Video Solution](#)

$$117. \lim_{x \rightarrow \infty} \sqrt{x}(\sqrt{x+c} - \sqrt{x}) =$$

A. 1

B. ∞

C. c

D. $c/2$

Answer: D



Watch Video Solution

$$118. \lim_{x \rightarrow \infty} x^{3/2} \left(\sqrt{x^3 + 1} - \sqrt{x^3 - 1} \right) =$$

- A. 1
- B. -1
- C. 0
- D. none

Answer: A



Watch Video Solution

$$119. \lim_{x \rightarrow \infty} \left[\sqrt{x^2 + 2x - 1} - x \right] =$$

- A. ∞

B. $1/2$

C. 4

D. 1

Answer: D



Watch Video Solution

120. If $\lim_{x \rightarrow \infty} \left(\sqrt{x^2 - ax} - x \right) = \frac{1}{2}$ then a=

A. 2

B. 0

C. -1

D. $1/2$

Answer: C



Watch Video Solution

121. The values of constants a so that $\lim_{x \rightarrow \infty} \left(\frac{x^2 + 1}{x + 1} - ax - b \right) = 0$

are

A. 2

B. -2

C. 0

D. 1

Answer: D



Watch Video Solution

122. $\lim_{x \rightarrow \infty} \left\{ \sin \sqrt{x+1} - \sin \sqrt{x} \right\} =$

A. 0

B. 1

C. 1/2

D. -1/2

Answer: C



Watch Video Solution

$$123. \lim_{x \rightarrow \infty} \left\{ \sqrt{x^2 + 1} - \sqrt{x^2 - 1} \right\} =$$

A. 0

B. 1

C. $1/2$

D. $-1/2$

Answer: A



Watch Video Solution

$$124. \lim_{n \rightarrow \infty} \left(\sqrt{n^4 + 1} - \sqrt{n^4 - 1} \right) =$$

A. 2

A. 0

C. $1/2$

D. none

Answer: B



Watch Video Solution

$$125. \lim_{x \rightarrow \infty} \left[\sqrt{x + \sqrt{x + \sqrt{x}}} - \sqrt{x} \right] =$$

A. 0

B. $1/2$

C. $\log 2$

D. e_4

Answer: B



Watch Video Solution

$$126. \lim_{n \rightarrow \infty} \frac{1 + 2 + 3 + \dots + n}{n^2} =$$

A. 1

B. $1/2$

C. $1/3$

D. $1/4$

Answer: B



Watch Video Solution

$$127. \lim_{n \rightarrow \infty} \frac{1^2 + 2^2 + 3^2 + \dots + n^2}{n^3} =$$

A. 1

B. $1/2$

C. $1/3$

D. $1/4$

Answer: C



Watch Video Solution

$$128. \lim_{n \rightarrow \infty} \frac{1^3 + 2^3 + 3^3 + \dots + n^3}{n^4} =$$

A. 1

B. $1/2$

C. $1/3$

D. $1/4$

Answer: D



Watch Video Solution

$$129. \lim_{n \rightarrow \infty} \frac{1 + 3 + 5 + \dots + (2n - 1)}{2 + 4 + 6 + \dots + 2n}$$

A. 1

B. $1/2$

C. $1/3$

D. $1/4$

Answer: A



Watch Video Solution

130. $\lim_{n \rightarrow \infty} \frac{1}{n^3} \sum_{k=1}^n [k^2 x] =$

A. x

B. $x/2$

C. $x/3$

D. $x/4$

Answer: C



Watch Video Solution

$$131. \lim_{n \rightarrow \infty} \frac{n(1^3 + 2^3 + \dots + n^3)^2}{(1^2 + 2^2 + \dots + n^2)^3} =$$

A. $27/16$

B. $16/27$

C. $9/4$

D. $4/9$

Answer: A



[Watch Video Solution](#)

$$132. \lim_{n \rightarrow \infty} \frac{(1 + 2 + \dots + n \text{ terms})(1^2 + 2^2 + \dots + n \text{ terms})}{n(1^3 + 2^3 + \dots + n \text{ terms})} =$$

A. $3/2$

B. $2/3$

C. 1

D. 0

Answer: B



Watch Video Solution

$$133. \lim_{n \rightarrow \infty} \frac{1 + 3 + 6 + \dots + n(n+1)/2}{n^3} =$$

A. $1/2$

B. $1/3$

C. $1/6$

D. $1/8$

Answer: C



Watch Video Solution

$$134. \lim_{n \rightarrow \infty} \frac{1}{n^4} [1^2 + (1^2 + 2^2) + \dots + (1^2 + 2^2 + \dots + n^2)] =$$

A. $1/6$

B. $1/16$

C. $1/12$

D. 0

Answer: C



Watch Video Solution

$$135. \lim_{n \rightarrow \infty} \left\{ \frac{1}{2} + \frac{1}{2^2} + \frac{1}{2^3} + \dots + \frac{1}{2^n} \right\} =$$

A. 1

B. $1/2$

C. $1/3$

D. 3

Answer: A



Watch Video Solution

136. If $0 < r < 1$ then $\lim_{n \rightarrow \infty} (a + ar + \dots + ar^{n-1}) =$

A. 0

B. $\frac{1}{1-r}$

C. $\frac{a}{1-r}$

D. none

Answer: C



Watch Video Solution

137. $\lim_{n \rightarrow \infty} \frac{1 + 3 + 3^2 + \dots + 3^n}{1 + 2 + 2^2 + \dots + 2^n} =$

A. 0

B. $3/2$

C. ∞

D. none

Answer: C



Watch Video Solution

$$138. \lim_{n \rightarrow \infty} \frac{1 + a + a^2 + \dots + a^{n-1}}{1 + b + b^2 + \dots + b^{n-1}} = 0 \text{ if}$$

A. $a=b$

B. $a < b, b > 1$

C. $a > b, a > 1$

D. $1 < a < b$

Answer: B



Watch Video Solution

$$139. \lim_{n \rightarrow \infty} \left\{ \frac{1}{1.2} + \frac{1}{2.3} + \frac{1}{3.4} + \dots + \frac{1}{n(n+1)} \right\} =$$

A. 0

B. 1

C. -1

D. $1/2$

Answer: B



Watch Video Solution

$$140. \lim_{n \rightarrow \infty} \left\{ \frac{1}{1.3} + \frac{1}{3.5} + \frac{1}{5.7} + \dots + \frac{1}{(2n-1)(2n+1)} \right\} =$$

A. 1

B. $1/2$

C. $1/13$

D. $1/6$

Answer: B



Watch Video Solution

$$141. \lim_{n \rightarrow \infty} \left\{ \frac{1}{1.4} + \frac{1}{4.7} + \frac{1}{7.10} + \dots + \frac{1}{(3n-2)(3n+1)} \right\} =$$

A. 1

B. $1/2$

C. $1/3$

D. $1/6$

Answer: C



Watch Video Solution

$$142. \lim_{n \rightarrow \infty} \frac{1.1! + 2.2! + 3.3! + \dots + n.n!}{(n+1)!} =$$

A. 0

B. ∞

C. 1

D. none

Answer: C



Watch Video Solution

$$143. \lim_{n \rightarrow \infty} \frac{2^n - n}{2^n} =$$

A. 0

B. 1

C. $-\infty$

D. ∞

Answer: B



Watch Video Solution

$$144. \lim_{n \rightarrow \infty} \frac{2^n - 1}{3^n + 1} =$$

A. 0

B. 1

C. $-\infty$

D. ∞

Answer: A



Watch Video Solution

$$145. \lim_{n \rightarrow \infty} \frac{5^n + 1}{3^n - 1} =$$

A. 0

B. 1

C. $-\infty$

D. ∞

Answer: D



Watch Video Solution

$$146. \lim_{n \rightarrow \infty} \frac{3^{x+1} + 4}{3^{x+2} + 4} =$$

- A. 3
- B. 1/3
- C. 1
- D. ∞

Answer: B



Watch Video Solution

$$147. \lim_{n \rightarrow \infty} \frac{2^{1/n} - 1}{2^{1/n} + 1} =$$

- A. 6/5
- B. 2/7
- C. -2/7
- D. -6/7

Answer: D



Watch Video Solution

148. $\lim_{n \rightarrow \infty} \frac{2^{1/n} - 1}{2^{1/n} + 1} =$

A. 0

B. 1

C. -1

D. 1/2

Answer: A



Watch Video Solution

149. If $a > 1$, then $\lim_{x \rightarrow \infty} \frac{a^x - a^{-x}}{a^x + a^{-x}} =$

A. e

B. a

C. 1

D. 0

Answer: C



Watch Video Solution

150. If $a > 1$, then $\frac{a^x - x^{-x}}{a^x + a^{-x}} =$

A. -1

B. 0

C. 1

D. ∞

Answer: C



Watch Video Solution

$$151. \lim_{n \rightarrow \infty} \frac{{}^nP_n}{{}^{n+1}P_{n+1} - {}^nP_n} =$$

A. 1

B. 0

C. -1

D. none

Answer: B



Watch Video Solution

$$152. \text{ If } |x| < 1 \text{ then } \lim_{n \rightarrow \infty} (1+x)(1+x^2)(1+x^4) \dots (1+x^{2n}) =$$

A. $1/(x-1)$

B. $1/(1-x)$

C. 0

D. $1-x$

Answer: B



Watch Video Solution

$$153. \text{ Lt}_{x \rightarrow \infty} \frac{3\sqrt{x} + \log x - \sin^2 x}{5\sqrt{x} + 6\cos^2 x - 100\log x} =$$

A. $1/3$

B. $3/5$

C. $-5/3$

D. $-3/5$

Answer: B



Watch Video Solution

$$154. \text{ Lt}_{x \rightarrow 1} \frac{\log(1-x)}{\cot \pi x} =$$

A. -1

A. 0

C. 1

D. 2

Answer: B



Watch Video Solution

$$155. \lim_{x \rightarrow \infty} 5^x \sin\left(\frac{a}{5^x}\right) =$$

A. 0

B. 5

C. $\log 5$

D. a

Answer: D



Watch Video Solution

$$156. \text{ Lt}_{x \rightarrow 0} \frac{(2+x)\sin(2+x) - 2\sin 2}{x} =$$

- A. $\sin 2$
- B. $\cos 2$
- C. $2 \sin 2 + \cos 2$
- D. $2 \cos 2 + \sin 2$

Answer: D



Watch Video Solution

$$157. \text{ Lt}_{x \rightarrow 0} \frac{\sin(x+a) + \sin(a-x) - 2\sin a}{x \sin x} =$$

- A. $\sin a$
- B. $-\sin a$
- C. $\cos a$
- D. $-\cos a$

Answer: B



Watch Video Solution

$$158. \text{ Lt}_{x \rightarrow 0} \frac{\sin(x + a) + \sin(a - x) - 2 \sin a}{x \sin x} =$$

A. $a(a \cos a + 2 \sin a)$

B. $a(a \cos a - 2 \sin a)$

C. $(a \cos a + 2 \sin a)$

D. $(a \cos a - 2 \sin a)$

Answer: A



Watch Video Solution

$$159. \text{ If } \text{Lt}_{x \rightarrow 0} \frac{x(1 + a \cos x) - b \sin x}{x^3} = 1 \text{ then } (a, b) =$$

A. $(0, 3/2)$

B. (1,2/3)

C. (5/2,-2/3)

D. (-5/2,-3/2)

Answer: D



Watch Video Solution

160. If $\lim_{x \rightarrow 0} \left(\frac{\cos 4x + a \cos 2x + b}{x^4} \right)$ is finite, then the value of a, b are respectively.

A. 5,-4

B. - 5, - 4

C. - 4, 3

D. 4,5

Answer: C



Watch Video Solution

161. $\lim_{x \rightarrow 1} \frac{x + x^2 + \dots + x^n - n}{x - 1} =$

A. $\frac{n(n + 1)}{2}$

B. $\frac{n(n + 1)(2n + 1)}{6}$

C. $\frac{n^2(n + 1)^2}{4}$

D. none

Answer: A



[Watch Video Solution](#)

162. $\lim_{x \rightarrow 0} \frac{2}{x} \log(1 + x) =$

A. 1

B. 2

C. 3

D. 4

Answer: B



Watch Video Solution

163. $\lim_{x \rightarrow 0} \frac{\log_e(1 + x)}{3^x - 1} =$

A. $\log_e 3$

B. 0

C. 1

D. $\log_3 e$

Answer: D



Watch Video Solution

164. $\lim_{x \rightarrow 0} \left\{ \frac{1}{x} - \frac{\log(1 + x)}{x^2} \right\} =$

A. 0

B. 1

C. $1/2$

D. $-1/2$

Answer: C



Watch Video Solution

$$165. \lim_{x \rightarrow 0} \frac{x \cos - \log(1 + x)}{x^2} =$$

A. $1/2$

B. 0

C. 1

D. none

Answer: A



Watch Video Solution

$$166. \lim_{x \rightarrow 0} \left(\frac{1}{\log x} - \frac{x}{x-1} \right) =$$

A. 2

B. -2

C. 4

D. -1/2

Answer: D



Watch Video Solution

$$167. \lim_{x \rightarrow 0} \left(\frac{1}{x} - \frac{1}{e^x - 1} \right) =$$

A. 1/2

B. 2

C. 1/3

D. 3

Answer: A



Watch Video Solution

$$168. \lim_{x \rightarrow \infty} x[\log(x+1) - \log x] =$$

A. 0

B. -1

C. 1

D. e

Answer: C



Watch Video Solution

$$169. \lim_{x \rightarrow 0} \log \left| \frac{\log(1+x)}{x} \right| =$$

A. 0

B. 1

C. e

D. $1/e$

Answer: A



Watch Video Solution

170. If $\lim_{x \rightarrow 0} \frac{\log(3+x) - \log(3-x)}{x} = k$ then $k =$

A. 0

B. $-1/3$

C. $2/3$

D. $-2/3$

Answer: C



Watch Video Solution

$$171. \lim_{x \rightarrow 0} \frac{\sin x - x}{x^3} =$$

A. $1/6$

B. $-1/6$

C. $1/5$

D. $-1/5$

Answer: B



Watch Video Solution

$$172. \lim_{x \rightarrow 0} \left\{ \operatorname{cosec} x - \frac{1}{x} \right\}$$

A. 0

B. 1

C. -1

D. 2

Answer: A



Watch Video Solution

173. $\lim_{x \rightarrow 0} \frac{\operatorname{cosec} x - \frac{1}{x} - \frac{x}{6}}{x^3} =$

A. $3/360$

B. $-2/360$

C. $7/360$

D. $11/360$

Answer: C



View Text Solution

174. $\lim_{x \rightarrow 0} \frac{\operatorname{cosec} x - \frac{1}{x} - \frac{x}{6}}{x^3} =$

A. $1/120$

B. $1/110$

C. $1/100$

D. $1/90$

Answer: A



[View Text Solution](#)

175. Lt $\frac{y^y - x^x}{y - x} =$

A. xy^{x-1}

B. x^x

C. $x^x(1 + \log x)$

D. none

Answer: C



[Watch Video Solution](#)

176. If $a > 0$ and $\lim_{x \rightarrow a} \frac{a^x - x^a}{x^x - a^a} = -1$ then $a =$

A. 0

B. 1

C. 3

D. 2e

Answer: B



Watch Video Solution

177. $\lim_{x \rightarrow -\infty} e^x =$

A. 0

B. 1

C. $-\infty$

D. ∞

Answer: A



Watch Video Solution

178. $\lim_{x \rightarrow 0} \frac{e^{mx} - 1}{x} =$

A. e

B. m

C. e^m

D. $m \log a$

Answer: B



Watch Video Solution

179. $\lim_{y \rightarrow 0} \frac{a^y - 1}{b^y - 1} =$

A. $\frac{\log a}{\log b}$

B. $\log \frac{a}{b}$

C. $\log(a - b)$

D. $\log a^b$

Answer: A



Watch Video Solution

180. $\lim_{x \rightarrow 0} \frac{a^x - b^x}{x} =$

A. 0

B. 1

C. $\log\left(\frac{a}{b}\right)$

D. $\frac{\log a}{\log b}$

Answer: C



Watch Video Solution

$$181. \lim_{x \rightarrow 0} \frac{a^{\tan x} - 1}{x} =$$

A. 1

B. a

C. $2 \log a$

D. $\log a$

Answer: D



Watch Video Solution

$$182. \lim_{x \rightarrow 0} \frac{e^{\sin x} - 1}{x} =$$

A. 1

B. e

C. e^{-1}

D. 0

Answer: A



Watch Video Solution

$$183. \lim_{x \rightarrow 0} \frac{e^{\tan x} - 1}{\sin x} =$$

A. 1

B. e

C. e^{-1}

D. 0

Answer: A



Watch Video Solution

$$184. \lim_{x \rightarrow 0} \frac{(x 2^{(x)-x})/(1-\cos x)}{=}$$

A. 0

B. 1

C. $\log 2$

D. $\log 4$

Answer: D



Watch Video Solution

$$185. \lim_{x \rightarrow 0} \frac{x \cdot 10^x - x}{1 - \cos x} =$$

A. $\log 10$

B. $2 \log 10$

C. $3 \log 10$

D. $4 \log 10$

Answer: B



Watch Video Solution

$$186. \text{ Lt}_{x \rightarrow 0} \frac{2^x - 1}{(\sqrt{1+x} - 1)} =$$

A. 0

B. 1

C. $\log 2$

D. $\log 4$

Answer: D



Watch Video Solution

$$187. \text{ Lt}_{x \rightarrow 0} \frac{a^x - 1}{\sqrt{a+x} - \sqrt{a}} =$$

A. $2\sqrt{a} \log a$

B. $\sqrt{a} \log a$

C. $\log a$

D. none

Answer: A



Watch Video Solution

$$188. \lim_{x \rightarrow 0} \frac{\sqrt{1+x^2} - \sqrt{1-x+x^2}}{3^x - 1} =$$

A. $\log 9$

B. $\frac{1}{\log 9}$

C. $\log 3$

D. $\frac{1}{\log 3}$

Answer: B



Watch Video Solution

$$189. \lim_{x \rightarrow 0} \frac{5^x + 3^x - 2^x - 1}{x} =$$

A. 5

B. 1

C. $\log 20$

D. $\log 15/2$

Answer: D



Watch Video Solution

$$190. \lim_{x \rightarrow 0} \frac{(ab)^x - a^x - b^x + 1}{x^2} =$$

A. $\log a \log b$

B. $\log(ab)$

C. $\log a/b$

D. $\log b/a$

Answer: A



Watch Video Solution

$$191. \text{ Lt}_{x \rightarrow 0} \frac{10^x - 2^x - 5^x + 1}{x^2} =$$

- A. $\log 5$
- B. $\log 2$
- C. $\log 5 \cdot \log 2$
- D. $\log 5 / \log 2$

Answer: C



Watch Video Solution

$$192. \text{ Lt}_{x \rightarrow 0} \left(\frac{a^x + b^x}{2} \right)^{1/x} =$$

- A. ab
- B. $a+b$
- C. \sqrt{ab}

D. none

Answer: C



Watch Video Solution

193. $\lim_{x \rightarrow 0} \left(\frac{16^x + 9^x}{2} \right)^{1/x} =$

A. 25/2

B. 12

C. 1

D. 0

Answer: B



Watch Video Solution

194. $\lim_{x \rightarrow \infty} \left[\frac{a^{1/x} + b^{1/x} + c^{1/x}}{3} \right]^x$, where a,b,c are real and nonzero=

A. 0

B. $(abc)^{1/3}$

C. $(abc)^{-1/3}$

D. $abc/3$

Answer: B



Watch Video Solution

195. The value of $\lim_{x \rightarrow 0} \left(\frac{a_1^{1/x} + a_2^{1/x} + \dots + a_n^{1/x}}{n} \right)^{nx}$ is

A. $a_1 + a_2 + \dots + a_n$

B. $e^{a_1 + a_2 + \dots + a_n}$

C. $\frac{a_1 + a_2 + \dots + a_n}{n}$

D. $a_1 a_2 \dots a_n$

Answer: D



www.smartickmethod.com



Watch Video Solution

$$196. \text{ Lt}_{x \rightarrow 0} \left(\frac{1^x + 2^x + 3^x + \dots + n^x}{n} \right)^{1/x}$$

A. $(n!)^n$

B. $(n!)^{1/n}$

C. $n!$

D. $\ln(n!)$

Answer: B



Watch Video Solution

$$197. \text{ Lt}_{x \rightarrow 0} \frac{e^{x^2} - \cos x}{x^2} =$$

A. 0

B. 1

C. 1/2

D. 3/2

Answer: D



Watch Video Solution

198. $\lim_{x \rightarrow 0} \frac{(1 - e^x)\sin x}{x^2 + x^3} =$

A. -1

B. 0

C. 1

D. 2

Answer: A



Watch Video Solution

199. $\lim_{x \rightarrow 0} \frac{e^{\tan x} - e^x}{\tan x - x} =$

A. 1

B. e

C. e^{-1}

D. 0

Answer: A



Watch Video Solution

$$200. \lim_{x \rightarrow 0} \frac{e^x - e^{\sin x}}{2(x - \sin x)} =$$

A. $-1/2$

B. $1/2$

C. 1

D. $3/2$

Answer: B



Watch Video Solution

$$201. \text{ Lt}_{x \rightarrow 0} \frac{x^a - a^a}{a^x - a^n} =$$

A. 0

B. a^a

C. $\log_e a$

D. $\log_a e$

Answer: D



[Watch Video Solution](#)

$$202. \text{ Lt}_{x \rightarrow a} \frac{e^a - e^x}{e^{x+a}(x-a)} =$$

A. $-1/e^a$

B. e^{-a}

C. a^a

D. none

Answer: A



Watch Video Solution

$$203. \text{ Lt}_{x \rightarrow 1} \frac{\sin(e^{x-1} - 1)}{\log x} =$$

A. 1

B. 0

C. e

D. none

Answer: A



Watch Video Solution

$$204. \text{ Lt}_{x \rightarrow 0} (1 + 2x)^{1/x} =$$

A. e

B. e^2

C. e^3

D. $-e$

Answer: B



Watch Video Solution

205. $\lim_{x \rightarrow 0} (1 + x)^{2/x} =$

A. e

B. e^2

C. e^3

D. $-e$

Answer: B



Watch Video Solution

206. $\lim_{x \rightarrow 0} (1 + \sin x)^{\cot x} =$

A. e

B. e^2

C. e^3

D. e^4

Answer: D



Watch Video Solution

207. $\lim_{x \rightarrow 0} (1 + \tan x)^{\cot x} =$

A. -1

B. 0

C. 1

D. e

Answer: D



Watch Video Solution

208. $\lim_{x \rightarrow 0} \left(1 + \frac{3}{x}\right)^x =$

A. e

B. e^2

C. e^3

D. $-e$

Answer: C



Watch Video Solution

209. The value of $\lim_{x \rightarrow 0} \frac{(1+x)^{1/x} - e}{x}$ is

A. 1

B. $e/2$

C. $-e/2$

D. $2/e$

Answer: C



Watch Video Solution

$$210. \lim_{x \rightarrow \infty} \left(\frac{x+1}{x+2} \right)^{x+3} =$$

A. 0

B. 1

C. e

D. $1/e$

Answer: D



Watch Video Solution

$$211. \lim_{x \rightarrow \infty} \left(\frac{x+5}{x+2} \right)^{x+3} =$$

A. e

B. e^2

C. e^3

D. e^5

Answer: C



Watch Video Solution

$$212. \lim_{x \rightarrow \infty} \left(\frac{x+6}{x+1} \right)^{x+4} =$$

A. e^4

B. e^6

C. e^5

D. e

Answer: C



Watch Video Solution

$$213. \lim_{x \rightarrow 0} \left(\frac{x+a}{x+b} \right)^{x+b} =$$

A. 1

B. e^{b-a}

C. e^{a-b}

D. e^b

Answer: C



Watch Video Solution

$$214. \lim_{x \rightarrow \infty} \left(\frac{x^2 + 5x + 3}{x^2 + x + 2} \right)^x =$$

A. e^4

B. $-e$

C. e

D. e^2

Answer: A



Watch Video Solution

$$215. \lim_{x \rightarrow \infty} \left(\frac{x}{1+x} \right)^x =$$

A. 0

B. 1

C. e

D. $1/e$

Answer: D



Watch Video Solution

216. If a, b, c, d are positive real numbers then $\lim_{n \rightarrow \infty} \left(1 + \frac{1}{a + bn}\right)^{c + dn} =$

A. $e^{d/b}$

B. $e^{c/a}$

C. $e^{\frac{c+d}{a+b}}$

D. e

Answer: A



Watch Video Solution

217. If $\lim_{x \rightarrow 0} \left(1 + \frac{a}{x} + \frac{b}{x^2}\right)^{2x} = e^2$, then the value of a and b are

A. $a \in R, b \in R$

B. $a = 1$ and $b = 2$

C. $a \in R, b = 2$

D. $a = 1, b \in R$

Answer: D



Watch Video Solution

218. $\lim_{x \rightarrow 0} (1 + \sin x)^{\cot x} =$

A. e

B. e^2

C. e^3

D. e^4

Answer: A



Watch Video Solution

219. $\lim_{x \rightarrow 0} (\cos x + a \sin bx)^{1/x} =$

A. e^a

B. e^b

C. $e^{a/b}$

D. e^{ab}

Answer: D



Watch Video Solution

220. $\lim_{x \rightarrow \infty} \left(\frac{x^2 + 2x - 1}{2x^2 - 3x - 2} \right)^{\frac{2x+1}{2x-1}} =$

A. $-1/2$

B. $-1/4$

C. $e^{1/4}$

D. $e^{-1/4}$

Answer: A



Watch Video Solution

221. $\lim_{x \rightarrow 0} \frac{e^{3x} - 1}{\log(1 + 5x)}$

A. 1

B. 3/5

C. 1

D. none

Answer: B

[Watch Video Solution](#)

222. $\lim_{x \rightarrow 1} (\log_2 2x)^{\log_x 5} =$

A. 5/2

B. $e^{\log_2 5}$

C. $e^{\log_e 2}$

D. $\log 5 / \log 2$

Answer: B



Watch Video Solution

223. $\lim_{x \rightarrow 0} x^x =$

A. 0

B. 1

C. e

D. $1/e$

Answer: B



Watch Video Solution

224. $\lim_{x \rightarrow 0} (\sin x)^{\tan x} =$

A. -1

B. 0

C. 1

D. ∞

Answer: C



Watch Video Solution

$$225. \lim_{x \rightarrow 0} (x + \sin x)^{\tan x} =$$

A. -1

B. 0

C. 1

D. 2

Answer: C



Watch Video Solution

226. $\lim_{x \rightarrow 0} \{(1 - x^2) \frac{1}{\log(1 - x)}\} =$

A. e

B. e^2

C. e^3

D. e^4

Answer: A



[View Text Solution](#)

227. $\lim_{x \rightarrow 1} (1 - x)^{\tan \pi x} =$

A. -1

B. 0

C. 1

D. 2

Answer: C



Watch Video Solution

228. If $f(x) = \left(\frac{x}{2+x}\right)^{2x}$, then

A. $\lim_{x \rightarrow \infty} f(x) = e^{-6}$

B. $\lim_{x \rightarrow \infty} f(x) = 2$

C. $\lim_{x \rightarrow \infty} f(x) = e^{-4}$

D. $\lim_{x \rightarrow \infty} f(x) = 1/9$

Answer: C



Watch Video Solution

229. $\lim_{x \rightarrow 0} \left(\frac{\tan x}{x} \right)^{1/x^2} =$

A. e^2

B. $e^{1/2}$

C. $e^{-1/2}$

D. $e^{1/3}$

Answer: D



Watch Video Solution

230. $\lim_{x \rightarrow 0} \left(\frac{\sin x}{x} \right)^{\frac{\sin x}{x - \sin x}} =$

A. 0

B. 1

C. e

D. 1/e

Answer: D



Watch Video Solution

$$231. \lim_{x \rightarrow 0} (\sin x + \cos x)^{1/x} =$$

A. e^{-1}

B. e

C. e^2

D. e^{-2}

Answer: B**Watch Video Solution**

$$232. \lim_{x \rightarrow \frac{\pi}{2}} \left(\frac{1 + \cos x}{1 - \cos x} \right)^{1/\cos x} =$$

A. e

B. e^2

C. e^3

D. e^4

Answer: B



Watch Video Solution

$$233. \lim_{x \rightarrow 0} \left(\frac{1 + \tan x}{1 + \sin x} \right)^{1/x} =$$

A. e^2

B. e^{-2}

C. e^6

D. 1

Answer: D



Watch Video Solution

$$234. \lim_{x \rightarrow 0} \left(\frac{1 + \tan x}{1 + \sin x} \right)^{\operatorname{cosec} x} =$$

A. 1

B. e

C. e^{-1}

D. none

Answer: A



Watch Video Solution

Exercise 1B(One Sided Limits)

1. $\lim_{x \rightarrow 0} \sqrt{x} =$

A. 0

B. $\sqrt{2}$

C. $\sqrt{3}$

D. does not exist

Answer: A



Watch Video Solution

2. $\lim_{x \rightarrow 1} \sqrt{1-x} =$

A. 0

B. $\sqrt{2}$

C. $\sqrt{3}$

D. does not exist

Answer: A



Watch Video Solution

3. $\lim_{x \rightarrow 2^-} \sqrt{2-x} =$

A. 0

B. $\sqrt{2}$

C. $\sqrt{3}$

D. does not exist

Answer: A



Watch Video Solution

$$4. \lim_{x \rightarrow 3^+} \sqrt{x - 3} =$$

A. 0

B. $\sqrt{2}$

C. $\sqrt{3}$

D. does not exist

Answer: A



Watch Video Solution

$$5. \lim_{x \rightarrow 2} \sqrt{4 - x^2} =$$

A. 0

B. $\sqrt{2}$

C. $\sqrt{3}$

D. does not exist

Answer: A



[Watch Video Solution](#)

$$6. \lim_{x \rightarrow 0} |x| =$$

A. 0

B. 1

C. -1

D. does not exist

Answer: A



Watch Video Solution

$$7. \lim_{x \rightarrow 1} \frac{|x|}{x} =$$

A. 0

B. 1

C. -1

D. does not exist

Answer: B



Watch Video Solution

$$8. \lim_{x \rightarrow -2} \frac{|x|}{x} =$$

A. 0

B. 1

C. -1

D. does not exist

Answer: C



Watch Video Solution

$$9. \lim_{x \rightarrow 3^-} \frac{|x - 3|}{x - 3} =$$

A. 0

B. 1

C. -1

D. does not exist

Answer: C



Watch Video Solution

$$10. \lim_{x \rightarrow 0^+} \left\{ \frac{|x|}{x} + x^2 + 3 \right\} =$$

A. 2

B. 3

C. 4

D. 6

Answer: C



Watch Video Solution

$$11. \lim_{x \rightarrow 0} \frac{3|x| + x}{7|x| - 5x} =$$

A. $3/2$

B. $3/10$

C. $1/8$

D. does not exist

Answer: D



Watch Video Solution

$$12. \lim_{x \rightarrow \infty} \frac{3|x| + x}{7|x| - 5x} =$$

A. 0

B. 1

C. 2

D. does not exist

Answer: C



Watch Video Solution

$$13. \lim_{x \rightarrow -\infty} \frac{2|x| + x}{5|x| - 3x} =$$

A. $3/2$

B. $3/10$

C. $1/8$

D. does not exist

Answer: C



Watch Video Solution

$$14. \lim_{x \rightarrow 0^-} \frac{5|x| + 2x}{7|x| - 3x} =$$

A. $3/2$

B. $3/10$

C. $1/8$

D. does not exist

Answer: B



Watch Video Solution

$$15. \lim_{x \rightarrow 1} \frac{x^2 - 1}{|x - 1|} =$$

- A. 2
- B. -2
- C. ∞
- D. does not exist

Answer: D



Watch Video Solution

$$16. \lim_{x \rightarrow 0} \frac{\sin x}{|x|} =$$

- A. 1
- B. -1
- C. 0
- D. does not exist

Answer: D



Watch Video Solution

$$17. \lim_{x \rightarrow 0} \frac{|\sin x|}{x} =$$

A. 1

B. -1

C. 0

D. does not exist

Answer: D



Watch Video Solution

$$18. \lim_{x \rightarrow 0} \frac{\sin|x|}{|x|} =$$

A. 1

B. -1

C. 0

D. does not exist

Answer: A



Watch Video Solution

$$19. \lim_{x \rightarrow 0} \frac{\sin x^2}{|x|} =$$

A. 0

B. 1

C. -1

D. does not exist

Answer: A



Watch Video Solution

$$20. \text{ Lt}_{x \rightarrow 2} \left(\frac{\sqrt{1 - \cos\{2(x - 2)\}}}{x - 2} \right)$$

A. equals $-\sqrt{2}$

B. equals $1/\sqrt{2}$

C. does not exist

D. equals $\sqrt{2}$

Answer: C



Watch Video Solution

$$21. \text{ Lt}_{x \rightarrow 0} \sin\left(\frac{1}{x}\right) =$$

A. 0

B. 1

C. -1

D. does not exist

Answer: D



Watch Video Solution

22. $\lim_{x \rightarrow 0} x \sin\left(\frac{1}{x}\right) =$

A. 0

B. 1

C. does not exist

D. cannot be determine

Answer: A



Watch Video Solution

23. $\lim_{x \rightarrow 0} x \sin\left(\frac{1}{x}\right) =$

A. 0

B. 1

C. -1

D. does not exist

Answer: B



Watch Video Solution

$$24. \lim_{x \rightarrow 0} x \sin\left[\frac{1}{x^2}\right] =$$

A. 0

B. 1

C. -1

D. does not exist

Answer: A



Watch Video Solution

$$25. \lim_{x \rightarrow 0} x^2 \sin(\pi/x) =$$

A. 1

B. 0

C. does not exist

D. ∞

Answer: B



Watch Video Solution

$$26. \lim_{x \rightarrow 0} \cos\left(\frac{1}{x}\right) =$$

A. 0

B. 1

C. -1

D. does not exist

Answer: D



Watch Video Solution

27. $\lim_{x \rightarrow 0} x \cos\left(\frac{1}{x}\right) =$

A. 0

B. 1

C. -1

D. does not exist

Answer: A



Watch Video Solution

28. $\lim_{x \rightarrow \pi/2} \tan x$

A. 0

B. ∞

C. $-\infty$

D. does not exist

Answer: C



Watch Video Solution

$$29. \lim_{x \rightarrow 1^-} \left(\frac{|x|}{x} \right)$$

A. 0

B. 1

C. -1

D. does not exist

Answer: C



Watch Video Solution

30. $\lim_{x \rightarrow 3} |x|$

- A. 3
- B. ∞
- C. $-\infty$
- D. does not exist

Answer: D



Watch Video Solution

31. $\lim_{x \rightarrow \infty} x \tan \frac{1}{x} =$

- A. 0
- B. 1
- C. -1
- D. does not exist

Answer: B



Watch Video Solution

32. $\lim_{x \rightarrow 0} \cot x =$

A. 0

B. ∞

C. $-\infty$

D. does not exist

Answer: B



Watch Video Solution

33. $\lim_{x \rightarrow 0} \cot x =$

A. 0

B. ∞

C. $-\infty$

D. does not exist

Answer: C



Watch Video Solution

$$34. \lim_{x \rightarrow \infty} \sqrt{\frac{x - \sin x}{x + \cos^2(x)}} =$$

A. 0

B. 1

C. -1

D. ∞

Answer: B



Watch Video Solution

$$35. \lim_{n \rightarrow \infty} \frac{\sin n\theta}{\sqrt{n}} =$$

- A. 0
- B. 1
- C. -1
- D. does not exist

Answer: A



Watch Video Solution

$$36. \lim_{x \rightarrow \pi/2} \left(e^{-\tan \theta} \right) \sec^2 \theta =$$

- A. -1
- B. 0
- C. 1
- D. does not exist

Answer: D



Watch Video Solution

$$37. \lim_{x \rightarrow -\infty} \frac{x^4 \cdot \sin(1/x) + x^2}{1 + |x|^3} =$$

A. 1

B. -1

C. 2

D. -3

Answer: B



Watch Video Solution

$$38. \lim_{x \rightarrow 0} \frac{1}{x} =$$

A. 0

B. ∞

C. $-\infty$

D. does not exist

Answer: D



Watch Video Solution

$$39. \lim_{x \rightarrow \infty} \frac{1}{x} =$$

A. 0

B. ∞

C. $-\infty$

D. does not exist

Answer: A



Watch Video Solution

$$40. \lim_{x \rightarrow 0} \frac{1}{x^2} =$$

- A. 0
- B. ∞
- C. $-\infty$
- D. does not exist

Answer: B



[Watch Video Solution](#)

$$41. \lim_{x \rightarrow \infty} \frac{1}{x^2} =$$

- A. 0
- B. ∞
- C. $-\infty$
- D. does not exist

Answer: A



Watch Video Solution

$$42. \lim_{x \rightarrow \infty} \frac{1}{x^2} =$$

A. 0

B. ∞

C. $-\infty$

D. does not exist

Answer: A



Watch Video Solution

$$43. \lim_{x \rightarrow 0} e^{1/x} =$$

A. 0

B. 1

C. -1

D. does not exist

Answer: D



Watch Video Solution

$$44. \lim_{x \rightarrow \infty} \frac{e^{1/x} - e^{-1/x}}{e^{1/x} + e^{-1/x}} =$$

A. -1

B. 1

C. 0

D. does not exist

Answer: C



Watch Video Solution

$$45. \lim_{x \rightarrow 0} \frac{2^{1/x} - 1}{2^{1/x} + 1} =$$

A. -1

B. 1

C. 0

D. ∞

Answer: B



Watch Video Solution

$$46. \lim_{x \rightarrow 3} [x] =$$

A. -1

B. 3

C. 8

D. does not exist

Answer: D



Watch Video Solution

47. $\lim_{x \rightarrow 4^+} \{[x] + x\} =$

A. -1

B. 3

C. 8

D. does not exist

Answer: C



Watch Video Solution

48. $\lim_{x \rightarrow 2^-} \{[x] + x\} =$

A. -1

B. 3

C. 8

D. does not exist

Answer: A



Watch Video Solution

49. $\lim_{x \rightarrow 5^+} \{x - [x]\} =$

A. 0

B. $1/2$

C. $-1/2$

D. does not exist

Answer: A



View Text Solution

$$50. \text{ Lt}_{x \rightarrow 0^+} [x] \sin \frac{1}{x} =$$

A. 0

B. 1

C. -1

D. does not exist

Answer: A



[View Text Solution](#)

$$51. \text{ Lt}_{x \rightarrow \pi/2} [\sin x] =$$

A. 1

B. 0

C. -1

D. none

Answer: B



View Text Solution

52. If $a > 0$ then $\underset{x \rightarrow \infty}{\text{Lt}} ([ax+b])/(x) =$

A. 0

B. 1

C. a

D. b

Answer: C



View Text Solution

53. $\underset{x \rightarrow \infty}{\text{Lt}} \frac{[x] + [2x] + [3x] + \dots + [nx]}{n^2} =$

A. $x/2$

B. $x/3$

C. x

D. 0

Answer: A



[View Text Solution](#)

$$54. \lim_{x \rightarrow 0} \frac{[1^2x] + [2^2x] + [3^2x] + \dots + [n^2x]}{n^3} =$$

A. $x/2$

B. $x/3$

C. $x/6$

D. 0

Answer: B



[View Text Solution](#)

55. $\lim_{x \rightarrow 1} \{1 - x + [x - 1] + [1 - x]\}$ is

- A. 0
- B. 1
- C. -1
- D. none

Answer: C



[View Text Solution](#)

56. $\lim_{x \rightarrow 0} \{1 - x + [x - 1] + [1 - x]\}$ is

- A. 0
- B. 1
- C. 2
- D. 3

Answer: D



View Text Solution

57.

If

$f: R \rightarrow R$ is defined by $f(x) = [x - 3] + [x - 4]$ for $x \in R$ then $\lim_{x \rightarrow 3^-} f(x)$

A. -2

B. -1

C. 0

D. 1

Answer: C



View Text Solution

58.

If

$f(x) = \frac{\sin[x]}{[x]}$ at $[x] \neq 0$ and $f(x) = 0$ at $x = 0$ then $\lim_{x \rightarrow 0} f(x)$

A. 0

B. 1

C. -1

D. does not exist

Answer: D



View Text Solution

59. If $f(x) = \begin{cases} \frac{\sin(1 + [x])}{x} & \text{for } [x] \neq 0 \\ 0 & \text{for } [x]=0 \end{cases}$ where $[x]$ denotes the greatest integer not exceeding x then $\lim_{x \rightarrow 0^-} f(x) =$

A. -1

B. 0

C. 1

D. 2

Answer: B

 View Text Solution

60. Let $f: R \rightarrow R$ be a positive increasing function with $\lim_{x \rightarrow \infty} \frac{f(3x)}{f(x)} = 1$

A. 1

B. $2/3$

C. $3/2$

D. 3

Answer: A

 View Text Solution

61. If $f(x) = \sqrt{9 - x^2}$ then $\lim_{x \rightarrow 2} \frac{f(x) - f(2)}{x - 2}$ is

A. $3/\sqrt{5}$

B. $-2/\sqrt{5}$

C. 0

D. none

Answer: B



Watch Video Solution

62. If $f(x) = -\sqrt{25 - x^2}$, then $\lim_{x \rightarrow 1} \frac{f(x) - f(1)}{x - 1} =$

A. $1/24$

B. $1/5$

C. $-\sqrt{24}$

D. $1/\sqrt{24}$

Answer: D



Watch Video Solution

63. If $f(x) = x \tan^{-1} x$ then $\lim_{x \rightarrow 1} \left(\frac{f(x) - f(1)}{x - 1} \right) =$

A. $\frac{\pi}{4}$

B. $\frac{\pi + 1}{4}$

C. $\frac{\pi + 2}{4}$

D. $\frac{\pi + 3}{4}$

Answer: C



Watch Video Solution

64.

If

$f: \mathbb{R} \rightarrow \mathbb{R}$ is defined by $f(x) \{ : ((x-2)/(x^2-3x+2)), \text{ if } x \in \mathbb{R}-\{1,2\}), ($

A. 0

B. -1

C. 1

D. $-1/2$

Answer: B



View Text Solution

65. If $f(x) = \begin{cases} x-5, & \text{for } x \leq 1 \\ 4x^2 - 8, & \text{for } 1 < x \leq 2 \\ 3x+4, & \text{for } x > 2 \end{cases}$ then $f(2+) - f(2-) =$

A. 0

B. 2

C. 3

D. 4

Answer: C



View Text Solution

66. If $f(x) = xf$ or $x < 0 = 0f$ or $x = 0=x^2$ for "x gt 0" then

"underset(x to 0)"Lt" f(x)=`

A. -1

B. 0

C. 1

D. 2

Answer: B



[View Text Solution](#)

67. If $f(x) = \begin{cases} 2x + b & (x < a) \\ x + d & (x \geq a) \end{cases}$ is such that $\lim_{x \rightarrow a^-} f(x) = l$ then $l =$

A. $2d-b$

B. $2b-d$

C. $2d+b$

D. b-2d

Answer: A



Watch Video Solution

68.

$$f(x) = \begin{cases} \sin x & x \neq n\pi \quad n \in \mathbb{Z} \\ 2 & \text{otherwise} \end{cases} \quad \text{and} \quad g(x) = \begin{cases} x^2 + 1 & x \neq 0, 2 \\ 4 & x = 0 \\ 5 & x = 2 \end{cases}$$

If

A. 2

B. 4

C. 5

D. 1

Answer: D



Watch Video Solution

69. If $f(x) = \begin{cases} 1 & (x < 0) \\ 2x + 1 & (x \leq 1) \\ 3x & (x > 1) \end{cases}$ then $\lim_{x \rightarrow 1} f(x) =$

A. 2

B. 0

C. 3

D. -1

Answer: C



Watch Video Solution

70. $f(x) = \begin{cases} x + 2 & (1 < x \leq 2) \\ x^2 & (2 < x < 5) \end{cases}$ then $\lim_{x \rightarrow 2} f(x) =$

A. 2

B. 4

C. 3

D. -1

Answer: B



Watch Video Solution

71. If $f(x) = \begin{cases} x/2 & (x < 2) \\ x^2/3 & (x \geq 2) \end{cases}$ then $\lim_{x \rightarrow 2^+} f(x) =$

A. 1

B. $4/3$

C. 3

D. -1

Answer: B



View Text Solution

Exercise 1C(Continuity)

1.

The

function

$f(x) = \tan x$ for $x \neq \pi/2$, $f(\pi/2) = 0$ at $x = \pi/2$ is

- A. continuous
- B. discontinuous
- C. not determined
- D. none

Answer: B



[View Text Solution](#)

2. The function $f(x) = \frac{\sin^2 x}{x}$ for $x \neq 0$, $f(0) = 1$ at $x = 0$ is

- A. continuous
- B. discontinuous
- C. not determined

D. none

Answer: B



View Text Solution

3. The function $f(x) = \frac{\tan x}{x}$ for $x \neq 0$, $f(0) = 1$ at $x = 0$ is

A. continuous

B. discontinuous

C. not determined

D. none

Answer: C



View Text Solution

4. If $f(x) = \frac{|x|}{x}$ for $x \neq 0$, $f(0) = 0$, then $f(x)$ at $x=0$ is

A. continuous

B. discontinuous

C. not determined

D. none

Answer: B



[View Text Solution](#)

5. The function $f(x) = \sqrt{x}$ at $x=0$ is

A. continuous

B. discontinuous

C. not determined

D. none

Answer: A



[View Text Solution](#)

6. The function $f(x) = \sqrt{x}$ at $x=0$ is

- A. continuous
- B. discontinuous
- C. not determined
- D. none

Answer: A



[View Text Solution](#)

7. The function $f(x) = x + [x]$ at $x = 1/2$ is

- A. continuous
- B. discontinuous
- C. not determined

D. none

Answer: A



Watch Video Solution

8. The function $f(x) = x + [x]$ at $x = 1/2$ is

A. continuous

B. discontinuous

C. not determined

D. none

Answer: A



Watch Video Solution

9. The function $f(x) = x - [x]$ when $x=2$ is

A. continuous

B. discontinuous

C. not determined

D. none

Answer: B



Watch Video Solution

10. The function $f(x) = (1 + x)^{5/x}$, $f(0) = e^5$ at $x = 0$ is

A. continuous

B. discontinuous

C. not determined

D. none

Answer: A



Watch Video Solution

11.

The

function

$$f(x) = \frac{x \tan 2x}{\sin 3x \sin 5x} \text{ for } x \neq 0, f(0) = 2/17. \text{ At } x=0 \text{ is}$$

- A. continuous
- B. discontinuous
- C. not determined
- D. none

Answer: B



Watch Video Solution

12.

The

function

$$f(x) = \frac{\cos 3x - \cos 4x}{x \sin 2x} \text{ for } x \neq 0, f(0) = 7/4 \text{ at } x=0 \text{ is}$$

- A. continuous
- B. discontinuous

C. not determined

D. none

Answer: A



[View Text Solution](#)

13. The function $f(x) = \frac{1 - \cos x}{x^2}$ for $x \neq 0$, $f(0) = 1$ at $x=0$ is

A. continuous

B. discontinuous

C. not determined

D. none

Answer: A



[Watch Video Solution](#)

14. The function $f(x) = \frac{3 \sin x - \sin 3x}{x^3}$ for $x \neq 0$, $f(0) = 1$ at $x=0$ is

- A. continuous
- B. discontinuous
- C. not determined
- D. none

Answer: B



Watch Video Solution

15. The function $f(x) = \frac{1 - \cos ax}{1 - \cos bx}$ for $x \neq 0$, $f(0) = \frac{a}{b}$ at $x=0$ is

- A. continuous
- B. discontinuous
- C. not determined
- D. none

Answer: B



Watch Video Solution

16.

The

function

$$f(x) = \left(\sqrt{1+x^2} - \frac{\sqrt{1-x^2}}{x^2} \text{ for } x \neq 0, f(0) = 1 \text{ at } x = 0 \right)$$

- A. continuous
- B. discontinuous
- C. not determined
- D. none

Answer: B



Watch Video Solution

17.

The

function

$$f(x) = \frac{\sqrt{x^2 + 3} - 2}{x - 1} \text{ for } x \neq 1, f(1) = 1/2 \text{ at } x=1 \text{ is}$$

A. continuous

B. discontinuous

C. not determined

D. none

Answer: A



Watch Video Solution

18. The function $f(x) = \frac{7|x| + 5x}{7|x| - 5x}$ for $x \neq 0$, $f(0) = 6$ at $x = 0$ is

A. continuous

B. discontinuous

C. not determined

D. none

Answer: B



Watch Video Solution

19. If $f(x) = \frac{1}{3^{1/x} + 1}$, then at $x=0$, $f(x)$ is

- A. continuous
- B. discontinuous
- C. not determined
- D. none

Answer: B



Watch Video Solution

20. The function

$$f(x) = \left[\frac{x-1}{1+e^{\frac{1}{x-1}}} \right] \quad \text{for } x \neq 1, f(1) = 0 \text{ at } x=1 \text{ is}$$

- A. continuous
- B. discontinuous

C. not determined

D. none

Answer: A



Watch Video Solution

21. The function $f(x) = \begin{cases} \frac{e^{1/x^2}}{e^{1/x^2} - 1} & \text{for } x \neq 0, \\ f(0) = 1 & \text{at } x = 0 \end{cases}$ is

A. continuous

B. discontinuous

C. not determined

D. none

Answer: A



Watch Video Solution

22. $f(x) = (x - [x])\sin\left(\frac{1}{X}\right)$, at $x = 0$, f is

- A. continuous
- B. discontinuous
- C. not determined
- D. none

Answer: B



Watch Video Solution

23. If $f(x) = x \sin \frac{1}{x}$ for $x \neq 0$, $f(0) = 0$ then at $x = 0$, f(x) is

- A. continuous
- B. discontinuous
- C. not determined
- D. none

Answer: A



Watch Video Solution

24. If $f(x) = x \cos \frac{1}{x^2}$ for $x \neq 0$, $f(0) = 1$ then at $x=0$, $f(x)$ is

- A. continuous
- B. discontinuous
- C. not determined
- D. none

Answer: B



Watch Video Solution

25. If $f(x) = \frac{x}{1 + e^{1/x}}$ for $x \neq 0$, $f(0) = 0$ then at $x=0$, $f(x)$ is

- A. continuous

B. discontinuous

C. not determined

D. none

Answer: A



Watch Video Solution

26.

The

function

$$f(x) = \frac{xe^{1/x}}{1 + e^{1/x}} + \sin \frac{1}{x} \quad \text{for } x \neq 0, f(0) = 0 \text{ at } x = 0 \text{ is}$$

A. continuous

B. discontinuous

C. not determined

D. none

Answer: B



Watch Video Solution

27. If $f(x) = \frac{e^{1/x} - 1}{e^{1/x} + 1}$ for $x \neq 0$, $f(0) = 0$ then at $x=0$, $f(x)$ is

- A. continuous
- B. discontinuous
- C. not determined
- D. none

Answer: B



Watch Video Solution

28. If $f(x) = \frac{\sin^2 ax}{x^2}$ for $x \neq 0$, $f(0) = 1$ is continuous at $x=0$ then $a =$

- A. 0
- B. 1
- C. ± 1

D. 2

Answer: C



Watch Video Solution

29. if $f(x) = \frac{1 - \cos ax}{x \sin x}$ for $x \neq 0$, $f(0) = 1/2$ is continuous at $x=0$

then a=

A. 0

B. 1

C. ± 1

D. 2

Answer: C



Watch Video Solution

30. A function f is defined by

$$f(x) = \begin{cases} \frac{\sin(a+1)x + \sin x}{x} & \text{if } x < 0 \\ c & \text{if } x = 0 \\ \frac{(x+bx^2)^{\frac{1}{2}} - x^{\frac{1}{2}}}{bx^{\frac{3}{2}}} & \text{if } x > 0 \end{cases}$$

If $f(x)$ is continuous at $x = 0$, find the values of a, b and c .

A. 1

B. 2

C. $3/2$

D. $-3/2$

Answer: D



[Watch Video Solution](#)

31. $f(x) = |x|$ is

A. everywhere continuous

B. continuous only at 0

C. discontinuous at $x=0$

D. nowhere continuous

Answer: A



Watch Video Solution

32. If $f(x) = x^2$, for x rational

$= -x^2$, for x irrational, then

A. f is continuous at $x=0$ and $x=1/2$

B. f is discontinuous at $x=0$ and $x=1/2$

C. f is continuous at $x=0$ and discontinuous at $x=1/2$

D. f is discontinuous at $x=0$ and continuous at $x=1/2$

Answer: C



View Text Solution

33. If $f(x) = \frac{\tan^{-1}(x+2)}{|x+2|}$, $x \neq -2$ and $f(-2) = 2$ then

- A. f is continuous at -2
- B. f is not continuous at -2
- C. $\lim_{x \rightarrow -2} f(x) = 1$
- D. none

Answer: B



Watch Video Solution

34. $f(x)=1$ for x is rational $f(x)=-1$ for x irrational, f is continuous on

- A. R
- B. \emptyset
- C. $(-1, 1)$
- D. none

Answer: B



Watch Video Solution

35. If $a < b$ then $f(x) = \sqrt{((x-a)/(b-x))}$ is continuous on

A. (a,b)

B. $[a,b]$

C. $[a,b)$

D. $(a,b]$

Answer: C



Watch Video Solution

36. The function $f(x) = \begin{cases} 3x - 1 & \text{if } x < 1 \\ x^2 & \text{if } x > 2 \end{cases}$ is continuous

A. $(-\infty, 1)$

B. $(2, \infty)$

C. $(-\infty, 1) \cup (2, \infty)$

D. $(1, 2)$

Answer: C



Watch Video Solution

37. If $f: \mathbb{R} \rightarrow \mathbb{R}$ is defined by

$$f(x) = \begin{cases} \frac{x+2}{x^2+3x+2} & \text{if } x \in \mathbb{R} - \{-1, -2\} \\ -1 & \text{if } x = -2 \\ 1 & \text{if } x = -1 \end{cases} \quad \text{then is a continuous}$$

on the set

A. \mathbb{R}

B. $\mathbb{R} - \{-2\}$

C. $\mathbb{R} - \{-1\}$

D. $\mathbb{R} - \{-1, -2\}$

Answer: C



Watch Video Solution

38. Let ' f ' be a non-zero real valued continuous function satisfying $f(x + y) = f(x), f(y) \cdot f(y)$ for all $x, y \in R$ if $f(2) = 9$, then $f(6) =$

A. 3^2

B. 3^6

C. 3^4

D. 3^3

Answer: B



Watch Video Solution

39. If $f(x+y)=f(x)f(y)$ and
 $f(x) = 1 + g(x), h(x)$ where $\lim_{x \rightarrow 0} g(x) = \lim_{x \rightarrow 0} h(x)$ exists, then $f(x)$ is co

A. ϕ

B. R

C. $R - \{0\}$

D. $R - \{1\}$

Answer: B



[View Text Solution](#)

40. Let $f(x)=f(x)$ $f(y)$ for all x and y . If $f(x)$ is continuous at $x=1$. then $f(x)$ is continuous

A. ϕ

B. R

C. $R - \{0\}$

D. $R - \{1\}$

Answer: B



View Text Solution

41. A discontinuous function $y=f(x)$ satisfying

$x^2 + y^2 = 4$ is given by $f(x) =$

A. $f(x) = \sqrt{4 - x^2}$

B. $f(x) = \sqrt{4 + x^2}$

C. $f(x) = \sqrt{x^2 - 4}$

D. none

Answer: A



Watch Video Solution

42. The discontinuous points of $f(x) = \frac{1}{\log|x|}$ are

A. $0, \pm 2$

B. $1, \pm 2$

C. 0, ± 1

D. 0, ± 3

Answer: C



Watch Video Solution

43. If $f : \mathbb{R} \rightarrow \mathbb{R}$ is defined by $f(x) = x - [x]$, where $[x]$ is the greatest integer not exceeding x , then the set of discontinuous of f is

A. the empty set

B. \mathbb{R}

C. \mathbb{Z}

D. \mathbb{N}

Answer: C



Watch Video Solution

44. If $f: R \rightarrow R$ is a continuous defined by $f(x) = [x]\cos\left(\frac{2x - 1}{2}\right)\pi$,

where $[x]$ denotes the greatest integer function, then f is

- A. discontinuous only at non-zero integral values of x
- B. continuous only at y
- C. continuous only at x
- D. discontinuous only

Answer: C



[View Text Solution](#)

45. $f(x) = \sqrt{x - [x]}$ is discontinuous at

- A. square of integers
- B. every integer
- C. every non integer
- D. none

Answer: B



Watch Video Solution

46. The value of $f(0)$ so that $f(x) = (1 + x)^{5/x}$ is continuous at $x=0$ is

A. e^2

B. e^3

C. $e^{3/2}$

D. e^5

Answer: D



Watch Video Solution

47. The value of $f(0)$ so that $f(x) = \frac{\sin x}{x}$ is continuous at $x=0$ is

A. 1

B. 2

C. 3

D. $3/2$

Answer: A



Watch Video Solution

48. The value of $f(0)$ so that $f(x) = \frac{3x + 4 \tan x}{x}$ is continuous at $x=3$ is

A. $3/2$

B. 2

C. 3

D. 7

Answer: D



Watch Video Solution

49. The value of $f(0)$ so that $f(x) = \frac{3x + 4 \tan x}{x}$ is continuous at $x=0$ is

A. 5

B. 6

C. 7

D. none

Answer: C



[Watch Video Solution](#)

50. The value of $f(0)$ so that $f(x) = \frac{\sqrt{1+x} - 1}{x}$ is continuous at $x=0$ is

A. 1

B. 2

C. $1/2$

D. 3

Answer: C



Watch Video Solution

51. If $f(x) = x^{\frac{1}{x-1}}$ for $x \neq 1$ and if f is continuous at $x=1$ then $f(1)=$

A. 0

B. 1

C. e

D. $1/e$

Answer: C



Watch Video Solution

52. The value of $f(0)$ so that the function

$$f(x) = \frac{(27 - 2x)^{1/3} - 3}{9 - 3(243 + 5x)^{1/5}}$$
 is continuous at $x=0$ is

A. 2 / 3

B. 6

C. 2

D. none

Answer: C



Watch Video Solution

53. In order that the function $f(x) = (x + 1)^{\cot x}$ is continuous at $x=0$, $f(0)$ must be

A. 0

B. e

C. $1/e$

D. none

Answer: B



Watch Video Solution

54. The value of $f(0)$ so that the function $f(x) = \frac{2x - \sin^{-1} x}{2x + \tan^{-1} x}$ is continuous at $x=0$, is

A. 2

B. $1/3$

C. $2/3$

D. $-1/3$

Answer: B



Watch Video Solution

55. The value of $f(0)$ so that the function $f(x) = \frac{\sqrt{1+x} - \sqrt[3]{1+x}}{x}$ becomes continuous at $x=0$ is

A. $1/6$

B. $1/4$

C. 2

D. $1/3$

Answer: A



Watch Video Solution

56. The value of $f(0)$ so that $f(x) = \frac{\cos^2 x - \sin^2 x - 1}{\sqrt{x^2 + 1} - 1}$ is continuous at $x=0$ is

A. 2

B. 4

C. -2

D. -4

Answer: D



Watch Video Solution

57. Extend the definition of the following by continuity

$$f(x) = \frac{1 - \cos 7(x - \pi)}{5(x - \pi)^2} \text{ at the point } x = \pi$$

A. 0

B. 1

C. π

D. $\pi/2$

Answer: A



Watch Video Solution

58.

Let

$$f(x) = \frac{1 - \tan x}{4x - \pi}, x \neq \frac{\pi}{4}, x \in \left[0, \frac{\pi}{2}\right]. \quad \text{If } f(x) \text{ is continuous in } \left[0, \frac{\pi}{4}\right],$$

A. 1

B. -1

C. $-1/2$

D. $1/2$

Answer: C



Watch Video Solution

$$59. f(x) = \begin{cases} \frac{1 - \sqrt{2} \sin x}{\pi - 4x} & \text{if } x \neq \frac{\pi}{4} \\ k & \text{if } x = \frac{\pi}{4} \end{cases}$$

continuous at $x=(\pi)/(4)$, then $K=$

A. 4

B. 2

C. 1

D. $1/4$

Answer: D



Watch Video Solution

60. If $f: R \rightarrow R$ is defined by $f(x) = \begin{cases} \frac{1+3x^2-\cos 2x}{x^2} & \text{for } x \neq 0 \\ k & \text{for } x=0 \end{cases}$

A. 1

B. 5

C. 6

D. 0

Answer: B



Watch Video Solution

61. If $f: R \rightarrow R$ is defined by $f(x) = \begin{cases} \frac{\cos 3x - \cos x}{x^2} & \text{for } x \neq 0 \\ \lambda & \text{for } x = 0 \end{cases}$ and if
f is continuous at $x=0$, then $\lambda =$

A. -2

B. -4

C. -6

D. - 8

Answer: B



Watch Video Solution

62. If $f: R \rightarrow R$ is defined by $f(x) = \begin{cases} \frac{2 \sin x - \sin 2x}{2x \cos x} & \text{if } x \neq 0 \\ \alpha & \text{for } x = 0 \end{cases}$ then the value of 'alpha' so that f is continuous at 0 is

A. 2

B. 1

C. - 1

D. 0

Answer: D



Watch Video Solution

63. The value of $f(0)$ so that the function $f(x) = \frac{1 - \cos(1 - \cos x)}{x^4}$ is continuous every-where is

A. $1/8$

B. $1/2$

C. $1/4$

D. none

Answer: A



Watch Video Solution

64. Find the value of a so that the function $f(x) = \frac{(a + x)^2 \sin(a + x) - a^2 \sin a}{x}$ is continuous at $x = 0$.

A. $a^2 \cos a + a \sin a$

B. $a^2 \cos a + 2a \sin a$

C. $2a^2 + \cos a + a \sin a$

D. none

Answer: B



Watch Video Solution

65. The value of $f(2)$ so that $f(x) = \frac{2^{x+2} - 16}{4^x - 16}$ is continuous at $x=2$ is

A. 2

B. $1/2$

C. -2

D. $-1/2$

Answer: B



Watch Video Solution

66. If the function $f(x) = \frac{\log(1 + ax) - \log(1 - bx)}{x}$ is continuous at $x=0$, then $f(0)=$

- A. $a-b$
- B. $a+b$
- C. $\log a + \log b$
- D. $\log a - \log b$

Answer: B



Watch Video Solution

67. The value of $f(0)$ so that $f(x) = \frac{\log(1 + x/a) - \log(1 - x/b)}{x}$ is continuous at $x=0$ is

- A. $\frac{a+b}{ab}$
- B. $(a - b)$
- C. $\frac{ab}{a+b}$

D. $\frac{ab}{a - b}$

Answer: A



Watch Video Solution

68. Let $f(x) = \left(\log(1 + x + x^2) + \frac{\log(1 - x + x^2)}{\sec x - \cos x} \right)$, $x \neq 0$. Then the

value of $f(0)$ so that f is continuous at $x=0$ is

A. 1

B. 0

C. 2

D. none

Answer: A



View Text Solution

69. The value of $f(0)$ so that $f(x) = \frac{(4^x - 1)^3}{\sin(x/4)\log(1 + x^2/3)}$ is continuous everywhere is

A. $3(\log 4)^3$

B. $4(\log 4)^3$

C. $12(\log 4)^3$

D. $15(\log 4)^3$

Answer: C



Watch Video Solution

70. If $f(x) = \frac{(a^x - 1)^3}{\sin(x \log a) \log(1 + x^2 \log a^2)}$ is continuous at $x=0$, then

$f(0) =$

A. $\log a$

B. $2 \log a$

C. $\log a^{-1}$

D. $\log \sqrt{a}$

Answer: D



View Text Solution

71. If $f(x) = \begin{cases} \frac{72^x - 9^x - 8^x + 1}{\sqrt{2} - \sqrt{1 + \cos x}} & x \neq 0 \\ k \log 2 \log 3 & x = 0 \end{cases}$ is a continuous function then k is equal

to

A. $\sqrt{2}$

B. 24

C. $18\sqrt{3}$

D. $24\sqrt{2}$

Answer: D



View Text Solution

72. If $f: R \rightarrow R$ is continuous such that

$$f(x + y) = f(x) + f(y) \quad x \in R, y \in R \text{ and}$$

$$f(1) = 2 \text{ then } f(100) =$$

A. 100

B. 50

C. 200

D. 0

Answer: C



Watch Video Solution

73. Let f be a non zero continuous function satisfying $f(x+y)=f(x)f(y)$ for all

$x, y \in R$. If $f(2)=9$ then $f(3)$ is

A. 1

B. 27

C. 200

D. 0

Answer: B



Watch Video Solution

74. Given that the function f defined by $f(x) = \begin{cases} 2x - 1 & \text{if } x > 2 \\ k & \text{if } x=2 \\ x^2 - 1 & \text{if } x < 2 \end{cases}$ is continuous. Then the value of k is

A. 2

B. 3

C. 4

D. -3

Answer: B



Watch Video Solution

75. $f(x) = \frac{x^3 + x^2 - 16x + 20}{(x - 2)^2}$ if $x \neq 2$, $= k$ if $x = 2$, $f(x)$ is continuous at $x = 2$ then

A. 5

B. 7

C. 11

D. 15

Answer: B



[Watch Video Solution](#)

76. If $f(x) = \frac{x^2 - 10x + 25}{x^2 - 7x + 10}$ for $x \neq 5$ and f is continuous at $x=5$ then

$f(5) =$

A. 0

B. 5

C. 10

D. 25

Answer: A



Watch Video Solution

77. Let $f(x) = \frac{x + x^2 + \dots + x^n - n}{x - 1}$, $x \neq 1$, the value of $f(1)$

A. n

B. $\frac{n+1}{2}$

C. $\frac{n(n+1)}{2}$

D. $\frac{n(n-1)}{2}$

Answer: C



Watch Video Solution

78. If the function $f(x) = \frac{\sin 3x}{x}$ for $x \neq 0$ and $f(0) = \frac{k}{2}$ is

continuous at $x=0$ then $k=$

A. 3

B. 6

C. 9

D. 12

Answer: B



Watch Video Solution

79. If $f(x) = \begin{cases} \frac{\sqrt{1+kx} - \sqrt{1-kx}}{x} & \text{for } -1 \leq x < 0 \\ 2x^2 + 3x - 2 & \text{for } 0 \leq x \leq 1 \end{cases}$ is continuous at $x = 0$

then find k .

A. -4

B. -3

C. - 2

D. - 1

Answer: C



Watch Video Solution

80. If : [- 2, 2] → R is defined by

$$f(x) = \begin{cases} \frac{\sqrt{1+cx} - \sqrt{1-cx}}{x} & \text{for } -2 \leq x < 0 \\ \frac{x+3}{x+1} & \text{for } 0 \leq x \leq 2 \end{cases}$$

continuous on [-2, 2] then c is equal to

A. 3

B. $\frac{3}{2}$

C. $\frac{3}{\sqrt{2}}$

D. $\frac{2}{\sqrt{3}}$

Answer: B



Watch Video Solution

81. Let $f(x) = -2 \sin x$, if $x > -\frac{\pi}{2}$
 $f(x) = a \sin x + b$, if $-\frac{\pi}{2} \leq x \leq \frac{\pi}{2}$,
if $x < -\frac{\pi}{2}$, $f(x) = a \sin x + b$, if $\left(-\frac{\pi}{2}, \frac{\pi}{2} \right)$.
The value of d and b so that $f(x)$ is continuous everywhere are

A. $a=0, b=1$

B. $a=-1, b=1$

C. $a=-1, b=0$

D. $a=-1, b=1$

Answer: D



[View Text Solution](#)

82. If $[x]$ denotes the greatest integer not exceeding x and if the function f defined by $f(x) = \begin{cases} \frac{a+2 \cos x}{x^2} & x < 0 \\ b \tan \frac{\pi}{x+4} & x \geq 0 \end{cases}$ is continuous at $x = 0$, then the ordered pair (a,b) is equal to

A. (-2,1)

B. (-2,-1)

C. (- 1, $\sqrt{3}$)

D. (- 2, - $\sqrt{3}$)

Answer: B



Watch Video Solution

83. The values of p and q for which the function

$$f(x) = \begin{cases} \frac{\sin(p+1)x + \sin x}{x} & x < 0 \\ \frac{q^x + x^2\sqrt{-\sqrt{x}}}{x^{3/2}} & x > 0 \end{cases}$$

is continuous for all x in R, are

A. $p = \frac{3}{2}, q = \frac{1}{2}$

B. $p = \frac{1}{2}, q = \frac{3}{2}$

C. $p = \frac{1}{2}, q = -\frac{3}{2}$

D. $p = \frac{5}{2}, q = \frac{1}{2}$

Answer: A



[View Text Solution](#)

84. Let $f: R \rightarrow R$ is defined by $f(x) = \{(\alpha + (\sin [x])/(x), \text{if } x > 0), (2, \text{if } x = 0), (\beta + [(\sin x - x)/(x^3)]), \text{if } x < 0\}$
where $[y]$ denotes the integral part of y . If f is continuous at $x = 0$, then
 $\beta - \alpha =$

A. -1

B. 1

C. 0

D. 2

Answer: B



[View Text Solution](#)

85. If $R \rightarrow R$ b e defined $f(x) = \begin{cases} a^2 \cos^2 x + b^2 \sin^2 x & \text{if } x \leq 0 \\ e^{ax+b} & \text{if } x > 0 \end{cases}$ is a continuous function show that

A. $b=2 \log |a|$

B. $2b=\log |a|$

C. $b=\log |2a|$

D. $b^2 = \log|a|$

Answer: A



Watch Video Solution

86. $f: R - \{0\} \rightarrow R$ given by $f(x) = \frac{1}{x} - \frac{2}{e^{2x} - 1}$

can be made continuous at $x = 0$ by defining $f(0)$ as

A. 2

B. -1

C. 0

D. 1

Answer: D



Watch Video Solution

Exercise 2(Special Type Questions)

I. $\lim_{x \rightarrow 0} \frac{(1+x)^2 - (1-x)^2}{(1+x)^3 - (1-x)^3} - 2$

II. If $f(x) = \frac{4-7x}{3x+4}$ and $\lim_{x \rightarrow 2} f(x) = l$, $\lim_{x \rightarrow 0} f(x) = m$ then the equation whose roots are $1/l, 1/m$ is $x^2 - 1 = 0$

A. only I is true

B. only II is true

C. both I and II are true

D. neither I nor II are true

Answer: C

2. $\lim_{x \rightarrow 0} x \sin\left(\frac{1}{x}\right) =$

- A. only I is true
- B. only II is true
- C. both I and II are true
- D. neither I nor II are true

Answer: B



Watch Video Solution

3. I. If $a > 0$ then $\lim_{x \rightarrow \infty} \frac{[ax + b]}{x} = a$

II. $\lim_{x \rightarrow \pi/2} [\sin x] =$

- A. only I is true
- B. only II is true

C. both I and II are true

D. neither I nor II are true

Answer: C



[View Text Solution](#)

4. I. $\lim_{x \rightarrow 0} \sqrt{s}$ does not exist

II. $\lim_{x \rightarrow 1} \frac{|x|}{x}$ does not exist.

A. only I is true

B. only II is true

C. both I and II are true

D. neither I nor II are true

Answer: D



[View Text Solution](#)

$$5. \text{ I. } \lim_{x \rightarrow 0} \frac{\log_e(1+x)}{3^x - 1} = \log_3 e$$

$$\text{II. } \lim_{x \rightarrow 1} (\log_5 5x)^{\log_x 5} = e$$

- A. only I is true
- B. only II is true
- C. both I and II are true
- D. neither I nor II are true

Answer: C



[View Text Solution](#)

$$6. \lim_{n \rightarrow \infty} \frac{1.1! + 2.2! + 3.3! + \dots + n.n!}{(n+1)!} =$$

- A. only I is true
- B. only II is true
- C. both I and II are true
- D. neither I nor II are true

Answer: C



Watch Video Solution

7. I. The function $f(x) = \sqrt{x}$ is continuous at $x=0$

II. The function $f(x) = |x|$ is continuous at $x=0$.

A. only I is true

B. only II is true

C. both I and II are true

D. neither I nor II are true

Answer: C



View Text Solution

8. The function $f(x) = \frac{1 - \cos ax}{1 - \cos bx}$ for $x \neq 0$, $f(0) = \frac{a}{b}$ at $x=0$ is

- A. only I is true
- B. only II is true
- C. both I and II are true
- D. neither I nor II are true

Answer: B



Watch Video Solution

9. I. $\lim_{x \rightarrow \infty} \frac{\sin x}{x} = 1$

II. Every identity function is continuous on R.

- A. only I is true
- B. only II is true
- C. both I and II are true
- D. neither I nor II are true

Answer: B



Watch Video Solution

10. I. $\lim_{\theta \rightarrow 0} \frac{\sin(\theta^2)}{\theta} = \frac{\pi}{200}$

II. $f(x) = x^2 \sin(1/x)$ ($x \neq 0$) is continuous at $x=0$

A. only I is true

B. only II is true

C. both I and II are true

D. neither I nor II are true

Answer: C



View Text Solution

SET-3 Match The Following

1. Match the following

I. $f(x) = \frac{7|x| + 5x}{7|x| - 5x}$ for $x \neq 0, f(0) = 6$ at $x = 0$ is

a) continuous

II. $f(x) = \frac{\cos 3x - \cos 4x}{x \sin 2x}$ for $x \neq 0, f(0) = \frac{7}{4}$ at $x = 0$ is

b) discontinuous

III. If $f(x) = \frac{\sin^2 ax}{x^2}$ for $x = 0, f(0) = 4$ is continuous at

c) ± 1

$x = 0$ then $a =$

IV. If $f(x) = \frac{1 - \cos ax}{x \sin x}$ for $x = 0, f(0) = 1/2$ is continuous at

d) ± 2

$x = 0$ then $a =$

A. a,b,c,d

B. c,b,d,a

C. b,a,d,c

D. a,b,d,c

Answer: B



View Text Solution

2.

Match

the

following

- I. If $f(x) = \frac{\log(1+ax) - \log(1-bx)}{x}$ is continuous at $x=0$ then $f'(0) =$ a) 1
- II. If $f(x) = \frac{\log(1+x/a) - \log(1-x/b)}{x}$ is continuous at $x=0$ then $f'(0) =$ b) $a+b$
- III. If $f(x) = \frac{\log(1+x+x^2) + \log(1-x+x^2)}{\sec x - \cos x}$ is continuous at $x=0$ then $f'(0) =$ c) $\frac{a+b}{ab}$
- d) 0

A. a,c,b

B. b,c,a

C. d,c,b

D. c,d,a

Answer: A



[View Text Solution](#)

Set -4

1. Assertion (A) : $Lt_{x \rightarrow 0} \frac{|x|}{x} = 1$

Reason (R) : Limit of a function doesn't exist if left and right limits exists and are not equal the correct answer is

- A. Both A and R are true and R is the correct explanation of A
- B. Both A and R are true and R is not the correct explanation of A
- C. A is true but R is false
- D. A is false but R is true

Answer: D



Watch Video Solution

2. A: $Lt_{x \rightarrow 0} \left(\frac{\sin(x^2)}{x} \right) = \frac{\pi}{200}$

R: $Lt_{x \rightarrow 0} \frac{\sin(ax)}{x} = a$ if x is measured in radains

- A. Both A and R are true and R is the correct explanation of A
- B. Both A and R are true and R is not the correct explanation of A

C. A is true but R is false

D. A is false but R is true

Answer: A



[View Text Solution](#)

3. A: $\lim_{x \rightarrow \pi/2} \left(\frac{\cot x}{\pi/2 - x} \right) = 1$

R: If $\lim_{x \rightarrow a} f(x)$ exists, then $\lim_{x \rightarrow a} f(x) = \lim_{x \rightarrow 0} f(a + x) = \lim_{x \rightarrow a} f(a - x)$

A. Both A and R are true and R is the correct explanation of A

B. Both A and R are true and R is not the correct explanation of A

C. A is true but R is false

D. A is false but R is true

Answer: A



[View Text Solution](#)

4. A: $\lim_{x \rightarrow \infty} x \sin \frac{1}{x} = 0$

R: If $\lim_{x \rightarrow a} f(x) = 0$ and $g(x)$ is bounded on a deleted neighbourhood of a

then $\lim_{x \rightarrow a} f(x)g(x) = 0$

A. Both A and R are true and R is the correct explanation of A

B. Both A and R are true and R is not the correct explanation of A

C. A is true but R is false

D. A is false but R is true

Answer: A



[View Text Solution](#)

5. Show that the function $f(x) = \frac{x}{1 + e^{1/x}}, x \neq 0, f(0) = 0$ is continuous at $x = 0$

A. Both A and R are true and R is the correct explanation of A

B. Both A and R are true and R is not the correct explanation of A

C. A is true but R is false

D. A is false but R is true

Answer: A



Watch Video Solution

6. A: $f(x)=x-[x]$ is discontinuous at $x=2$

R: $\lim_{x \rightarrow a} f(x) = f(a)$

A. Both A and R are true and R is the correct explanation of A

B. Both A and R are true and R is not the correct explanation of A

C. A is true but R is false

D. A is false but R is true

Answer: B



View Text Solution

7. A: $f(x) = \frac{1}{1 + e^{1/x}}$ ($x \neq 0$) and $f(0) = 0$ is right continuous at $x=0$

R: $\lim_{x \rightarrow 0} \frac{1}{1 + e^{1/x}} = 0$

- A. Both A and R are true and R is the correct explanation of A
- B. Both A and R are true and R is not the correct explanation of A
- C. A is true but R is false
- D. A is false but R is true

Answer: A



[View Text Solution](#)

8. Let $f: R \rightarrow R$ be a continuous function defined by $f(x) = \frac{1}{e^x + 2e^{-x}}$

Statement -1: $f(x) = \frac{1}{3}$, for some $c \in R$.

Statement -2, $0 < f(x) \leq \frac{1}{2\sqrt{2}} \forall x \in R$.

- A. Statement-1 is true, Statement-2 is true, Statement -2 is a correct explanation for statement-1

B. Statement-1 is true, Statement-2 is true, Statement -2 is not a

correct explanation for statement-1

C. Statement-1 is true, Statement-2 is false

D. Statement-1 is false, Statement- is true

Answer: A



[Watch Video Solution](#)