



MATHS

BOOKS - DEEPTI MATHS (TELUGU ENGLISH)

PARABOLA

Examples

1. The equation to the parabola having focus (1,2) qbe and directrix $x + 2y + 5 = 0$ is

A. $4x^2 - 4xy + y^2 - 20x - 40y = 0$

B. $9x^2 - 24xy + 16y^2 - 76x + 18y + 91 = 0$

C. $x^2 + 6xy + 9y^2 - 28x - 16y + 46 = 0$

$$D. x^2 + 4xy + 4y^2 - 50x + 30y + 40 = 0$$

Answer: A



Watch Video Solution

2. The length of the latus rectum of the parabola

$$y^2 + 8x - 4y - 4 = 0.$$

A. 2

B. 1

C. 8

D. 3

Answer: C





Watch Video Solution

3. The equation of the directrix of the parabola $x^2 - 4x + 16y + 52 = 0$ is

A. $x-2=0$

B. $y+3=0$

C. $y-1=0$

D. $x+2=0$

Answer: C



Watch Video Solution

4. The point of contact of $x-y+2=0$ to the parabola $y^2 = 8x$ is

A. (2,4)

B. (-2,4)

C. (2,-4)

D. (-2,-4)

Answer: A



Watch Video Solution

5. The equation of the common tangent to $x^2 + y^2 = 18$ and $y^2 = 24x$ is

A. $y = \pm (x + 3)$

B. $y = \pm (x + 6)$

C. $y = \pm (x + 9)$

D. $y = \pm (x + 2)$

Answer: B



Watch Video Solution

6. An equilateral is inscribed in the parabola $y^2 = 8x$ with one of its vertices is the vertex of the parabola. Then the length of the side of that triangle is

A. $2\sqrt{3}$

B. $4\sqrt{3}$

C. $8\sqrt{3}$

D. $16\sqrt{3}$

Answer: D



Watch Video Solution

7. If $(\frac{1}{2}, 2)$ is one extremity of a focalchord of the parabola $y^2 = 8x$. Find the co-ordinates of the other extremity.

A. (8,8)

B. (-8,8)

C. (8,-8)

D. (-8,-8)

Answer: C



Watch Video Solution

8. The line $y = 2x - 12$ is a normal to the parabola $y^2 = 4x$ at the point P whose coordinates are

A. (4,-4)

B. (-2,-2)

C. (3,1)

D. (0,-4)

Answer: A



Watch Video Solution

9. The length of the chord of the parabola $y^2 = x$ are ends of the chord Mid point (2,1) is

A. 3

B. $\sqrt{14}$

C. $\sqrt{6}$

D. $2\sqrt{5}$

Answer: D

 [Watch Video Solution](#)

10. The diameter of the parabola $y^2 = 6x$ corresponding to the system of parallel chords $3x-y+c=0$ is

A. $y-1=0$

B. $y-2=0$

C. $y+1=0$

D. $y+2=0$

Answer: A



Watch Video Solution

11. If the tangent at $P(at^2, 2at)$ to the parabola $y^2 = ax$ intersects X-axis at A and the normal at P meets it at B then area of triangle PAB is

A. $4a^2|t|\sqrt{1+r^2}$

B. $2a^2|t|(1+r^2)$

C. $4a^2|t|(1 + t^2)$

D. $\frac{2a^2(1 + t^2)}{|t|}$

Answer: B



View Text Solution

Exercise 1 A

1. The parabola with directrix $x+2y-1=0$ and focus $(1,0)$ is

A. $4x^2 - 4xy + y^2 - 8x + 4y + 4 = 0$

B. $4x^2 + 4xy + y^2 - 8x + 4y + 4 = 0$

C. $4x^2 + 4xy + y^2 + 8x - 4y + 4 = 0$

D. $4x^2 - 4xy + y^2 - 8x - 4y + 4 = 0$

Answer: A



Watch Video Solution

2. The equation of the parabola with focus (1,-1) and directrix $x+y+3=0$ is

A. $x^2 + y^2 - 10x - 2y - 2xy - 5 = 0$

B. $x^2 + y^2 + 10x - 2y - 2xy - 5 = 0$

C. $x^2 + y^2 + 10x + 2y - 2xy - 5 = 0$

D. $x^2 + y^2 + 10x + 2y + 2xy - 5 = 0$

Answer: A



Watch Video Solution

3. The equation to the parabola having focus $(-1, -1)$ and directrix $2x - 3y + 6 = 0$ is

A. $x^2 - 2xy + y^2 - 18x - 10y - 45 = 0$

B. $9x^2 + 12xy + 4y^2 + 2x + 62y - 10 = 0$

C. $x^2 + 6xy + 9y^2 + 28x - 16y + 46 = 0$

D. $x^2 + 4xy + 4y^2 - 50x + 30y + 40 = 0$

Answer: B

 [Watch Video Solution](#)

4. The equation of the parabola whose vertex is origin axis along x-axis and which passes through the point $(-2,4)$ is

A. $y^2 = -8x$

B. $x^2 = 4y$

C. $y^2 = 8x$

D. $y^2 = -12x$

Answer: A



Watch Video Solution

5. The equation of the parabola whose vertex is origin axis is along y-axis and which passes through the point (2,1) is

A. $y^2 = -8x$

B. $x^2 = 4y$

C. $y^2 = 8x$

$$D. y^2 = -12x$$

Answer: B



Watch Video Solution

6. The equation of the parabola whose vertex is (3,-2) axis is parallel to x-axis and latus rectum 4 is

A. $(y + 2)^2 = \pm 4(x - 3)$

B. $(x + 2)^2 = \pm 6(y - 1)$

C. $(y - 1)^2 = 16(x + 2)$

D. $(y - 2)^2 = -8(x - 5)$

Answer: A



[Watch Video Solution](#)

7. The equation of the parabola having focus $(2, 1)$, and vertex $(-2, 1)$ is

A. $(y + 2)^2 = \pm 4(x - 3)$

B. $(x + 2)^2 = \pm 6(y - 1)$

C. $(y - 1)^2 = 16(x + 2)$

D. $(y - 2)^2 = -8(x - 5)$

Answer: C

[Watch Video Solution](#)

8. The equation of the parabola whose vertex is at $(0,0)$ and focus is the point of intersection of $x+y=2, 2x-y=4$ is

A. $y^2 = 2x$

B. $y^2 = 4x$

C. $y^2 = 8x$

D. $x^2 = 8y$

Answer: C

 [Watch Video Solution](#)

9. The equation of parabola whose vertex and focus are on x-axis at distances a and a' respectively from the origin is

A. $y^2 = 4(a' + a)(x - a)$

B. $y^2 = 4(a' - a)(x + a)$

C. $y^2 = 4(a' + a)(x + a)$

D. $y^2 = 4(a' - a)(x - a)$

Answer: A



Watch Video Solution

10. Find the equation of the parabola whose axis is parallel to X-axis and which passes through these points.

$(-2,1), (1,2),$ and $(-1,3)$

A. $5y^2 + 2x - 21y + 20 = 0$

B. $15y^2 + 12x - 11y + 10 = 0$

$$C. 18y^2 - 12x + 21y + 56 = 0$$

$$D. 25y^2 - 2x - 65y + 120 = 0$$

Answer: A



Watch Video Solution

11. Find the equation of the parabola whose axis is parallel to Y-axis and which passes through the points (4,5),(-2,11) and (-4,21).

$$A. x^2 - 4x - 2y + 10 = 0$$

$$B. x^2 - 2x - y + 5 = 0$$

$$C. x^3 - 4x - 2y + 10 = 0$$

$$D. y^2 - 2x - 3y + 4 = 0$$

Answer: A



Watch Video Solution

12. The equation of parabola whose latus rectum is the line segment joining the points $(-3,1)$, $(1,1)$ is

A. $(x + 1)^2 = 4y$

B. $(x - 1)^2 = 4y$

C. $(x + 1)^2 = 2y$

D. $(x - 1)^2 = 2y$

Answer: A



Watch Video Solution

13. The equation of the parabola with latusrectum joining the points (6,7) and (6,-1) is

A. $(y - 3)^3 = 8(x - 4)$

B. $(y + 3)^2 = 8(x + 4)$

C. $(y - 3)^2 = -8(x - 8)$

D. $y^2 = 4ax$

Answer: B

 [Watch Video Solution](#)

14. The vertex of a parabola is the point (a,b) and latusrectum is of length 1. If the axis of the parabola is along the positive direction of y-axis then its equation is

$$\text{A. } (x + a)^2 = \frac{1}{2}(2y - 2b)$$

$$\text{B. } (x - a)^2 = \frac{l}{2}(2y - 2b)$$

$$\text{C. } (x + a)^2 = \frac{l}{4}(2y - 2b)$$

$$\text{D. } (x - a)^2 = \frac{l}{2}(2y - 2b)$$

Answer: A



Watch Video Solution

15. The equation of the parabola having the vertex $(-1,-2)$ and whose axis is vertical and which passes through $(3,6)$ is

$$\text{A. } x^2 + 2x - 2y - 3 = 0$$

$$\text{B. } x^2 + 4x + 8y - 13 = 0$$

$$\text{C. } y^2 + 4y - 16x - 12 = 0$$

D. none

Answer: A



Watch Video Solution

16. The focus of a parabola is $(2,3)$ and the foot of the perpendicular from the focus to the directrix is $(4,5)$. The equation to the parabola is .

A. $(x - 2)^2 + (y - 3)^2 = (1/2)(x - y + 9)^2$

B. $(x - 2)^2 + (y - 3)^2 = (1/2)(x + y + 9)^2$

C. $(x - 2)^2 + (y - 3)^2 = (1/2)(x + y - 9)^2$

D. none

Answer: D



Watch Video Solution

17. The vertex of the parabola $y^2 + 4x - 2y + 3 = 0$ is

A. $(-3, 1)$

B. $(-3, 1)$

C. $(-3/2, 3)$

D. $(-1/2, 1)$

Answer: D



Watch Video Solution

18. The vertex of the parabola $x^2 + 8x + 12y + 4 = 0$ is

A. (-4,1)

B. (4,-1)

C. (-4,-1)

D. (4,1)

Answer: A

 [Watch Video Solution](#)

19. The vertex of the parabola $x^2 + 12x - 9y = 0$ is

A. (6,-4)

B. (-6,4)

C. (6,4)

D. (-6,-4)

Answer: D



Watch Video Solution

20. A parabola has the origin as its focus and the line $x=2$ as the directrix . Then the vertex of the parabola is at

A. (1,0)

B. (0, 1)

C. (2,0)

D. (0,2)

Answer: A

 [Watch Video Solution](#)

21. If the focus is $(1,-1)$ and the directrix is the line $x+2y-9=0$,
the vertex of the parabola is

- A. $(1,2)$
- B. $(2,1)$
- C. $(1,-2)$
- D. $(2,-1)$

Answer: B

 [Watch Video Solution](#)

22. The locus of the vertices of the family of parabolas

$$y = \frac{a^3 x^2}{3} + \frac{a^2 x}{2} - 2a \text{ is}$$

A. $xy = \frac{35}{16}$

B. $xy = \frac{64}{105}$

C. $xy = \frac{105}{64}$

D. $xy = \frac{3}{4}$

Answer: C



Watch Video Solution

23. The focus of the parabola $y^2 - x - 2y + 2 = 0$ is

A. $(1/4, 0)$

B. (1,2)

C. (3/4,0)

D. (5/4,1)

Answer: D



Watch Video Solution

24. The focus of the parabola $y^2 - 4y - 8x - 4 = 0$ is

A. (1,1)

B. (1,2)

C. (2,0)

D. (2,2)

Answer: B



Watch Video Solution

25. The focus of the parabola $x^2 - 2x - 8y - 23 = 0$ is

- A. (1,1)
- B. (1,-1)
- C. (-1,1)
- D. (-1,-1)

Answer: B



Watch Video Solution

26. The distance between the vertex and the focus of the parabola $x^2 - 2x + 3y - 2 = 0$ is

A. $\frac{4}{5}$

B. $\frac{3}{4}$

C. $\frac{1}{2}$

D. $\frac{5}{6}$

Answer: B



Watch Video Solution

27. The two ends of latusrectum of a parabola are the points $(3,6)$ and $(-5,6)$. The focus is

A. (1,6)

B. (-1,6)

C. (1,-6)

D. (-1,-6)

Answer: B



Watch Video Solution

28. If $(2,0)$ is the vertex and y -axis is the directrix of a parabola then its focus is

A. $(-4, 0)$

B. $(4,0)$

C. $(-2,0)$

D. (2,0)

Answer: B



Watch Video Solution

29. The length of the latus rectum of the parabola

$$y^2 + 8x - 2y + 17 = 0 \text{ is}$$

A. 2

B. 4

C. 8

D. 16

Answer: C





Watch Video Solution

30. The length of the latus rectum of the parabola

$$4y^2 + 12x - 20y + 67 = 0 \text{ is}$$

A. 2

B. 1

C. 8

D. 3

Answer: D



Watch Video Solution

31. The length of the latus rectum of the parabola

$$x^2 - 4x + 8y + 28 = 0 \text{ is}$$

A. 16

B. 4

C. 2

D. 8

Answer: D



Watch Video Solution

32. The length of the latus rectum of the parabola

$$3x^2 - 9x + 5y - 2 = 0 \text{ is}$$

A. 5

B. 4

C. 16

D. $\frac{5}{3}$

Answer: D



Watch Video Solution

33. The point (3,4) is the focus and $2x-3y+5=0$ is the directrix of a parabola . Its latusrectum is

A. $\frac{2}{\sqrt{13}}$

B. $\frac{4}{\sqrt{13}}$

C. $\frac{1}{\sqrt{13}}$

D. $\frac{3}{\sqrt{13}}$

Answer: A



Watch Video Solution

34. The length of the latusrectum of the parabola whose

focus is $\left(\frac{\mu^2}{2g} \sin 2\alpha, -\frac{\mu^2}{2g} \cos 2\alpha \right)$

A. $\frac{\mu^2}{g} \cos^2 \alpha$

B. $\frac{\mu^2}{g} \cos 2\alpha$

C. $\frac{2\mu^2}{g} \cos 2\alpha$

D. $\frac{2\mu^2}{g} \cos^2 \alpha$

Answer: D

 [Watch Video Solution](#)

35. Latus rectum of the parabola whose axis is parallel to the y-axis and which passes through the points (0,4), (1,9) and (-2,6) is equal to

A. 2

B. 1

C. $1/2$

D. $1/4$

Answer: C

 [View Text Solution](#)

36. The point (3,4) is the focus and $2x-3y+5=0$ is the directrix of a parabola . Its latusrectum is

A. $2/\sqrt{13}$

B. $4/\sqrt{13}$

C. $1/\sqrt{13}$

D. none

Answer: A



Watch Video Solution

37. If the parabola $y^2 = 4ax$ passes through (-3,2) then the length of its latusrectum is

A. $2/3$

B. $1/3$

C. $4/3$

D. 4

Answer: C



Watch Video Solution

38. If the straight line $y=mx+c$ is parallel to the axis of the parabola $y^2 = lx$ and intersects the parabola at $\left(\frac{c^2}{8}, c\right)$ then the length of the latus rectum is

A. 2

B. 3

C. 4

D. 8

Answer: D



Watch Video Solution

39. The ends of the latus rectum of the parabola

$$(x - 2)^2 = -6(y + 1) \text{ are}$$

A. (2,7),(3,-7)

B. (0,5),(0,-5)

C. (0,7),(0,-5)

D. (5,-5/2),(-1,-5/2)

Answer: D





Watch Video Solution

40. The equation of the latus rectum of the parabola

$$(y - 2)^2 = -4(x + 2) \text{ is}$$

A. $y=4$

B. $x=4$

C. $x+3=0$

D. $x+y=0$

Answer: C



Watch Video Solution

41. The equation of the directrix to the parabola

$$y^2 - 2x - 6y - 5 = 0 \text{ is}$$

A. $2x+15=0$

B. $x+5=0$

C. $2x+3=0$

D. $x+2=0$

Answer: A

 [Watch Video Solution](#)

42. The equation of the directrix of the parabola

$$y^2 + 4y + 4x + 2 = 0 \text{ is}$$

A. $x = -1$

B. $x=1$

C. $x=-3/2$

D. $x=3/2$

Answer: D



Watch Video Solution

43. The parabola $(y + 1)^2 = a(x - 2)$ passes through point (1,-2). The equation of its directrix is

A. $4x+1=0$

B. $4x-1=0$

C. $4x+9=0$

D. $4x-9=0$

Answer: D



Watch Video Solution

44. The equation of the directrix of the parabola

$$y^2 - 4ay - 2ax = 0 \text{ is}$$

A. $2y-13=0$

B. $2x+5a=0$

C. $2x+25=0$

D. $2x-13=0$

Answer: B





Watch Video Solution

45. The equation of the latus rectum of the parabola

$$x^2 - 12x - 8y + 52 = 0 \text{ is}$$

A. $x=4$

B. $y=4$

C. $x=6$

D. $y=2$

Answer: B



Watch Video Solution

46. The equation of the directrix of the parabola

$$3x^2 - 9x + 5y - 2 = 0 \text{ is}$$

A. $4y+9=0$

B. $y+4=0$

C. $y-3=0$

D. $6y-13=0$

Answer: D



Watch Video Solution

47. The equation of the directrix of the parabola whose vertex $(3,2)$ and focus $(2,-1)$ is

A. $x+3y-19=0$

B. $y-2y-9=0$

C. $2x+6y-24=0$

D. $x-3y-19=0$

Answer: A



Watch Video Solution

48. The equation of the axis of the parabola

$(y + 3)^2 = 4(x - 2)$ is

A. $x-5=0$

B. $y+3=0$

C. $2x-1=0$

D. $y-1=0$

Answer: B



Watch Video Solution

49. Axis of the parabola $x^2 - 3y - 6x + 6 = 0$ is

A. $x=-3$

B. $y=-1$

C. $x=3$

D. $y=1$

Answer: C



Watch Video Solution

50. The equation of the axis of the parabola

$$3x^2 - 9x + 5y - 2 = 0 \text{ is}$$

A. $x-2=0$

B. $x-1=0$

C. $x-3=0$

D. $2x-3=0$

Answer: D



Watch Video Solution

51. The parabola $y=px^2 + px + q$ is symmetrical about the line

A. $x=q$

B. $x=p$

C. $2x=1$

D. $2x+1=0$

Answer: D



Watch Video Solution

52. The focal distance of the point $(9, 6)$ on the parabola

$y^2 = 4x$ is

A. 4

B. 8

C. 10

D. 16

Answer: C



Watch Video Solution

53. The focal distance of the point (4,2) on the parabola

$$x^2 = 8y \text{ is}$$

A. 10

B. 4

C. 15

D. 12

Answer: B





Watch Video Solution

54. the eccentricity of the parabola $y^2 - 2x - 6y + 5 = 0$ is

A. 0

B. 1

C. $1/2$

D. 2

Answer: B



Watch Video Solution

55. If the vertex of the parabola $y = x^2 - 8x + c$ lies on x-axis then the value of c is

A. -16

B. -4

C. 4

D. 16

Answer: D

 [Watch Video Solution](#)

56. If the ordinate of a point on the parabola $y^2=4x$ is twice the latus rectum then the point is

A. (16,8)

B. (16,-8)

C. (- 16, 8)

D. (- 16, - 8)

Answer: A



Watch Video Solution

57. In the parabola $y^2 - 2y + 8x - 23 = 0$ the length of double ordinate at a distance of 3 from its vertex is

A. $4\sqrt{6}$

B. $2\sqrt{6}$

C. $\sqrt{6}$

D. none

Answer: A



Watch Video Solution

58. For a parabola the distance between the focus and the directrix is equal to

A. a

B. $4a$

C. $2a$

D. none

Answer: C





Watch Video Solution

59. If $(9,12)$ is one end of a double ordinate of the parabola $y^2 = 16x$ then its equation is

A. $x+9=0$

B. $y+9=0$

C. $y-9=0$

D. $x-9=0$

Answer: D



Watch Video Solution

60. The focal distance of a point on the parabola $y^2 = 8x$ whose focal distance is 10. Its coordinates are

A. $(2, \pm 2)$

B. $(3, \pm 3)$

C. $(5, \pm 5)$

D. $(8, \pm 8)$

Answer: D

 [Watch Video Solution](#)

61. The coordinates of the parabola $y^2 = 2x$ whose focal distance is $5/2$ are

A. $(2, \pm 2)$

B. $(3, \pm 3)$

C. $(5, \pm 5)$

D. $(8, \pm 8)$

Answer: A



Watch Video Solution

62. The point on the parabola $y^2 = 36x$ whose ordinate is three times its abscissa is

A. $(4,12)$

B. $(-4, 12)$

C. $(4, -12)$

D. $(-4, -12)$

Answer: A



Watch Video Solution

63. An equilateral is inscribed in the parabola $y^2 = 8x$ with one of its vertices is the vertex of the parabola. Then the length of the side of that triangle is

A. $24\sqrt{3}$

B. $16\sqrt{3}$

C. $8\sqrt{3}$

D. $4\sqrt{3}$

Answer: B



Watch Video Solution

64. If Q is the foot of the perpendicular from a point p on the parabola $y^2 = 8(x - 3)$ to its directrix. S is an equilateral triangle then find the length of side of the triangle.

A. $(4\sqrt{3}, 8)$

B. $(8, 4\sqrt{3})$

C. $(9, 4\sqrt{3})$

D. $(4\sqrt{3}, 9)$

Answer: C

 [Watch Video Solution](#)

65. L and L are the ends of the latus rectum of the parabola $x^2 = 6y$. The equation of OL and OL where O is the origin is

A. $x^2 + 4y^2 = 0$

B. $x^2 - 4y^2 = 0$

C. $x^2 + 2y^2 = 0$

D. $x^2 - 2y^2 = 0$

Answer: B

 [Watch Video Solution](#)

66. The coordinates of an endpoint of the latusrectum of the parabola $y^2 = 4ax$ at its vertex is

A. (0,-3)

B. (0,-1)

C. (0,1)

D. (1,3)

Answer: B

 [View Text Solution](#)

67. The angle subtended by the double ordinate of length $8a$ of the parabola $y^2 = 4ax$ at its vertex is

A. $\pi / 3$

B. $\pi / 4$

C. $\pi / 2$

D. $\pi / 6$

Answer: C



Watch Video Solution

68. If $2x+y+a=0$ is a focal chord of the parabola $y^2 + 8x = 0$

then $a =$

A. -4

B. 4

C. -2

D. 2

Answer: B



Watch Video Solution

69. PQ is a double ordinate of the parabola $y^2 + 4x$. The locus of its point of trisection is

A. $9y^2 + 4x = 0$

B. $4y^2 = 9x$

C. $9x^2 + 4y = 0$

D. $9y^2 = 4x$

Answer: D



Watch Video Solution

70. The ratio in which the line segment joining the points (4,-6) and (3,1) is divided by the parabola $y^2 = 4x$ is

A. $\frac{-20 \pm \sqrt{155}}{11} : 1$

B. $\frac{-2 \pm 2\sqrt{155}}{11} : 1$

C. $-20 \pm 2\sqrt{155} : 11$

D. $-20 \pm \sqrt{155} : 11$

Answer: C

 [Watch Video Solution](#)

71. The number of point of intersection of the circle $x^2 + y^2 = 2ax$ with the parabola $y^2 = x$ is

A. 3

B. 1

C. 2

D. 4

Answer: C



Watch Video Solution

72. If $a \neq 0$ and the line $2bx+3cy+4d = 0$ passes through the points of intersection of the parabolas $y^2 = 4ax$ and $x^2 = 4ay$ then

A. $d^2 + (2b + 3c)^3 = 0$

B. $d^2 + (3b - 2c)^2 = 0$

$$C. d^2 + (2b - 3c)^3 = 0$$

$$D. d^2 + (3b + 2c)^2 = 0$$

Answer: A



Watch Video Solution

73. Which of the following equations represents a parabola

$$A. (x - y)^3 = 3$$

$$B. \frac{x}{y} - \frac{y}{x} = 0$$

$$C. \frac{x}{y} + \frac{4}{x} = 0$$

$$D. (x + y)^2 + 3 = 0$$

Answer: C

 [Watch Video Solution](#)

74. The equation of the tangent to the parabola $y^2 = 12x$ at (3,-6) is

A. $x+y+3=0$

B. $x+y+1=0$

C. $x-y+2a=0$

D. $x+y+1=0$

Answer: A

 [Watch Video Solution](#)

75. The equation of the tangent to the parabola

$$x^2 = 4ya \text{ at } (-2, 1) \text{ is}$$

A. $x+y+3=0$

B. $x+y+1=0$

C. $x-y+2a=0$

D. $x+y+1=0$

Answer: D

 [Watch Video Solution](#)

76. The equation of the tangent to the parabola $y^2 = 4x$ at

the end of the latus rectum in the fourth quadrant is

A. $x + y + 3 = 0$

B. $x + y + 1 = 0$

C. $x - y + 2a = 0$

D. $x + y - 1 = 0$

Answer: B



Watch Video Solution

77. The equation of the tangent to the parabola $y^2 = 8x$ inclined at 30° to the x axis is

A. $3x - \sqrt{3}y + 4 = 0$

B. $2x - 3y + 14 = 0$

C. $2x - \sqrt{2}y + 7 = 0$

$$D. x - \sqrt{3y} + 6 = 0$$

Answer: D

 [Watch Video Solution](#)

78. The equation to the normal to the parabola $y^2 = 4x$ at (1,2) is

A. $x+y-3=0$

B. $x - y + 6 = 0$

C. $x - y + 5 = 0$

D. $x-y+4=0$

Answer: A





Watch Video Solution

79. The equation of the normal to the curve $x^2 = 4y$ at $(1,2)$ is

A. $2x+y+4=0$

B. $2x+y-4=0$

C. $2x-y+4=0$

D. $2x-y-4=0$

Answer: B



Watch Video Solution

80. The line $x+y=6$ is a normal to the parabola $y^2 = 8x$ at the point

A. (18, - 12)

B. (4,2)

C. (2,4)

D. (3,3)

Answer: C



Watch Video Solution

81. The equation of the normal at the end of latusrectum in the fourth quadrant of the parabola $y^2 = 4ax$ is

A. $x+y+3a=0$

B. $x+y-3a=0$

C. $x-y+3a=0$

D. $x-y-3a=0$

Answer: D



Watch Video Solution

82. The equation of the tangent to the parabola $y^2 = 8x$ and which is parallel to the line $x-y+3=0$ is

A. $x-y+2=0$

B. $x+y-2=0$

C. $x-y-2=0$

D. $2x-y+4=0$

Answer: A

 [Watch Video Solution](#)

83. The equation of the tangent to the parabola $y^2 = 16x$ and perpendicular to the line $x-4y-7=0$ is

A. $4x+y+1=0$

B. $4x+y+7=0$

C. $4x+y-1=0$

D. $4x+y-7=0$

Answer: A





Watch Video Solution

84. If the line $x+y+2=0$ touches the parabola $y^2 = kx$ then

$k=$

A. 2

B. 8

C. 1

D. 0

Answer: B



Watch Video Solution

85. Find the value of k if the line $2y=5x+k$ is a tangent to the parabola $y^2 = 6x$

A. $2/3$

B. $4/5$

C. $3/5$

D. $6/5$

Answer: D

 [Watch Video Solution](#)

86. If $x+y+k=0$ is a tangent to the parabola $x^2 = 4y$ then $k=$

A. 1

B. 2

C. -1

D. 4

Answer: A



Watch Video Solution

87. The straight line $x+y$ touches the parabola $y = x - x^2$

then $k =$

A. 0

B. -1

C. 1

D. none

Answer: C



Watch Video Solution

88. The line $y=2x+k$ is a normal to the parabola $y^2 = 4x$

then $k=$

A. 12

B. -12

C. 10

D. -10

Answer: B



Watch Video Solution

89. The line $y = x\sqrt{2} + \lambda$ is a normal to the parabola $y^2 = 4x$ then $\lambda =$

A. $4\sqrt{2}$

B. $-4\sqrt{2}$

C. $2\sqrt{2}$

D. $-2\sqrt{2}$

Answer: B

 [Watch Video Solution](#)

90. Let $x+y=k$ be a normal to the parabola $y^2 = 12x$. If p is the length of the perpendicular from the focus of the parabola onto this normal then $4k-2p^2 =$

A. 1

B. 0

C. -1

D. 2

Answer: B



Watch Video Solution

91. The point of contact of $2x - y + 2 = 0$ to the parabola

$$y^2 = 4ax \text{ is}$$

A. (2,4)

B. (3,4)

C. (1,4)

D. (2,1)

Answer: C



Watch Video Solution

92. The point of contact of $x-2y+4a=0$ to the parabola

$$y^2 = 4ax \text{ is}$$

A. (4a,4a)

B. (a,4a)

C. (3a,4a)

D. (4a,2a)

Answer: A





Watch Video Solution

93. The line $4x+6y+9=0$ touches the parabola $y^2=4ax$ at the point

A. $(-3, 9/4)$

B. $(3, -9/4)$

C. $(9/4, -3)$

D. $(-9/4, -3)$

Answer: C



Watch Video Solution

94. The point on the curve $y^2 = x$ the tangent at which makes an angle of 45° with x-axis will be given by

A. (2,4)

B. (1/2,1/2)

C. (1/2,1/4)

D. (1/4,1/2)

Answer: D



[Watch Video Solution](#)

95. The tangent to $y^2 = ax$ makes an angle 45° with x-axis .

Then its point of contact is

A. $(a/2, a/4)$

B. $(-a/2, a/4)$

C. $(a/4, a/2)$

D. $(-a/4, a/2)$

Answer: C



Watch Video Solution

96. The condition that the line $lx+my+n=0$ to touch the parabola $y^2 = 4ax$ is

A. $am^2 = ln$

B. $an^2 = lm$

C. $a^2m = ln$

D. $am=ln$

Answer: A

 [Watch Video Solution](#)

97. The line $y = m(x+a) + a/m$ touch the parabola $y^2 = 4a(x + a)$ for m

- A. is equal to 0
- B. is any positive real number
- C. is any negative number
- D. is any real number

Answer: D





Watch Video Solution

98. The condition that the line $y=mx+c$ to be a tangent to the parabola $y^2 = 4a(x + a)$ is

A. $c = a\left(m + \frac{1}{a}\right)$

B. $c = a\left(m + \frac{1}{m}\right)$

C. $c = a\left(m - \frac{1}{m}\right)$

D. $a = c\left(m + \frac{1}{m}\right)$

Answer: B



Watch Video Solution

99. The line among the following that touches the parabola

$$y^2 = 4ax \text{ is}$$

A. $x + my + am^3 = 0$

B. $x - my + am^2 = 0$

C. $x + my - am^2 = 0$

D. $y + mx + am^2 = 0$

Answer: B

 [Watch Video Solution](#)

100. The equation of the common tangent to

$$x^2 + y^2 = 2a^2 \text{ and } y^2 = 8ax \text{ is}$$

A. $y = \pm (x + a)$

B. $y = \pm (x + 2a)$

C. $y = \pm (x + 3a)$

D. $y = \pm (x + 4a)$

Answer: B



Watch Video Solution

101. The equation of the common tangent to $x^2 + y^2 = 8$ and $y^2 = 16x$ is

A. $y = \pm (x + 2)$

B. $y = \pm (x + 4)$

C. $2x+3y+36=0$

D. $3x+2y+24=0$

Answer: B



Watch Video Solution

102. The equation of the common tangent to $y^2 = 8x$ and

$x^2 + y^2 - 12x + 4 = 0$ is

A. $y = \pm (x + 2)$

B. $y = \pm (x + 4)$

C. $2x + 3y + 36 = 0$

D. $3x + 2y + 24 = 0$

Answer: A





Watch Video Solution

103. The slope of the line touching both the parabolas $y^2 = 4x$ and $x^2 = 32y$ is

A. $\frac{1}{8}$

B. $\frac{2}{3}$

C. $\frac{1}{2}$

D. $\frac{3}{2}$

Answer: C



Watch Video Solution

104. The points of intersection of the parabolas $y^2 = 5x$ and $x^2 = 5y$ lie on the line

A. $x+y=10$

B. $x-2y=0$

C. $x-y=0$

D. $2x-y=0$

Answer: C



Watch Video Solution

105. The two parabolas $y^2 = 4x$ and $x^2 = 4y$ intersect at a point P whose abscissae is not zero such that

- A. they both touch each other at P
- B. they cut at right angles at P
- C. the tangents to each curve at P make complementary angles with the x-axis
- D. none

Answer: C

 [Watch Video Solution](#)

106. If the common tangent of the circle $x^2 + y^2 = c^2$ and the parabola $y^2 = 4ax$ subtends an angle θ with x-axis then $\tan^2 \theta =$

A. $\frac{\sqrt{c^2 + 4a^2} - c}{2c}$

B. $\frac{\sqrt{c^2 + 4a^2} - c}{2}$

C. $\frac{\sqrt{3c^2 + 4a^2} - c}{2c}$

D. $\frac{\sqrt{c^2 + a^2} + 4c}{2c}$

Answer: A



Watch Video Solution

107. The sum of the slopes of the tangents to the parabola $y^2 = 8x$ drawn from the point $(-2,3)$ is

A. -1

B. -2

C. $-3/2$

D. 2

Answer: C



Watch Video Solution

108. The product of the slopes of the tangents to the parabola $y^2 = 4x$ drawn from the point (2,3) is

A. -1

B. -2

C. $-3/2$

D. $1/2$

Answer: D



Watch Video Solution

109. The slope of tangents drawn from a point $(4,10)$ to the parabola $y^2 = 9x$ are

A. $1/4, 3/4$

B. $1/4, 9/4$

C. $1/4, 1/3$

D. none

Answer: B



Watch Video Solution

110. The point of intersection of the tangents at the ends of latusrectum of the parabola $y^2 = 4x$ is

A. (0,0)

B. (0,1)

C. (-1,0)

D. (1,0)

Answer: C



Watch Video Solution

111. The locus of the point of intersection of perpendicular tangents to the parabola $y^2 = 4ax$ is

A. $x=a$

B. $x+a=0$

C. $y=a$

D. $y+a=0$

Answer: B



Watch Video Solution

112. If two tangents drawn from a point P to the parabola $y^2 = 4x$ are at right angles then the locus of P is

A. $x=1$

B. $2x+1=0$

C. $x=-1$

D. $2x-1=0$

Answer: C





Watch Video Solution

113. The tangents at the ends of a focal chord of a parabola $y^2 = 4ax$ intersect on the directrix at an angle of

A. 30°

B. 45°

C. 60°

D. 90°

Answer: D



Watch Video Solution

114. The locus of the point of intersection of the perpendicular tangents to the parabola $x^2 = 4ay$ is

A. $y=a$

B. $y=-a$

C. $x=a$

D. $x=-a$

Answer: B

 [Watch Video Solution](#)

115. The locus of the point of intersection of two tangents to the parabola $y^2 = 4ax$ which make an angle 30° with one another is

A. $(x + a)^2 = 3(y^2 - 4ax)$

B. $(x - a)^2 = y^2 - 4ax$

C. $3(x + a)^2 = y^2 - 4ax$

D. $x+a=0$

Answer: A



[View Text Solution](#)

116. The locus of the point of intersection of two tangents to the parabola $y^2 = 4ax$ which make complementary angles with the axis of the parabola is

A. $x=a$

B. $x+a=0$

C. $y=a$

D. $y+a=0$

Answer: A



Watch Video Solution

117. The locus of the point of intersection of two tangents to the parabola $y^2 = 4ax$ which make the angles θ_1 and θ_2 with the axis so that $\tan \theta_1 \tan \theta_2 = k$ is

A. $kx-y=0$

B. $kx-a=0$

C. $y=ka=0$

D. $x-ka=0$

Answer: B



Watch Video Solution

118. The locus of the point of intersection of two tangents to the parabola $y^2 = 4ax$ which make the angles θ_1 and θ_2 with the axis so that $\cot \theta_1 + \cot \theta_2 = k$ is

A. $kx - y = 0$

B. $kx - a = 0$

C. $y - ka = 0$

D. $x - ka = 0$

Answer: C



Watch Video Solution

119. The locus of the point of intersection of two tangents to the parabola $y^2 = 4ax$ which make the angles θ_1 and θ_2 with the axis so that $\tan \theta_1 \tan \theta_2 = k$ is

A. $kx - y = 0$

B. $kx - a = 0$

C. $y - ka = 0$

D. $kx^2 + 2ax - y^2 = 0$

Answer: D



Watch Video Solution

120. The locus of point of intersection of tangents to

$y^2 = 4ax$ which includes an angle α is

A. $(y^2 - 4ax)(x + a)^2 = a^2x^2$

B. $(y^2 - 4ax)^2(x - a)^2 = d^2x^2$

C. $y^2 - 4ax = d^2x^2(x + a)^2$

D. none

Answer: A



[Watch Video Solution](#)

121. Two straight lines are perpendicular to each other. One

of them touches the parabola $y^2 = 4a(x + a)$, and the

other touches $y^2 = 4b(x + b)$. Then locus of point of intersection of two lines is

A. $x+a=0$

B. $x+b=0$

C. $x+a+b=0$

D. $x-a-b=0$

Answer: C



[Watch Video Solution](#)

122. The locus of the point of intersection of tangents to the parabola $y^2 = 4(x + 1)$ and $y^2 = 8(x + 2)$ which are perpendicular to each other is

A. $x+7=0$

B. $x-y=4$

C. $x+3=0$

D. $y-x=12$

Answer: C



[View Text Solution](#)

123. The equation of a tangent to the parabola $y^2 = 8x$ is $y = x + 2$. The point on this line from which the other tangent to the parabola is perpendicular to the given tangent is

A. $(-1,1)$

B. (0,2)

C. (2,4)

D. (-2,0)

Answer: D



Watch Video Solution

124. Through the vertex O of the parabola $y^2 = 4ax$ a perpendicular is drawn to any tangent meeting it at P and the parabola at Q. Then $OP \cdot OQ =$

A. a^2

B. $2a^2$

C. $3a^2$

D. $4a^2$

Answer: D



View Text Solution

125. If M is the foot of the perpendicular from a point P on a parabola to its directrix and SPM is an equilateral triangle where S is the focus then SP=

- A. a
- B. 2a
- C. 3a
- D. 4a

Answer: D



Watch Video Solution

126. If y_1, y_2 are the ordinates of two points P and Q on the parabola and y_3 is the ordinate of the point of intersection of tangents at P and Q then

A. y_1, y_2, y^3 are in A.P..

B. y_1, y_3, y_2 are in A.P

C. y_1, y_2, y_3 are in G.P

D. y_1, y_3, y_2 are in G.P

Answer: B



Watch Video Solution

127. The locus of foot of perpendicular from the focus upon any tangent to the parabola $y^2 = 4ax$ is

A. l_1, l_2, l_3 are in G.P

B. l_2, l_1, l_3 are in G.P

C. l_3, l_1, l_2 are in A.P

D. l_3, l_2, l_1 are in A.P

Answer: B



Watch Video Solution

128. The length of the perpendicular from the focus S of the parabola $y^2 = 4ax$ on the tangent at P is

A. $\sqrt{OS \cdot SP}$

B. OS.SP

C. OS+OP

D. none

Answer: A



View Text Solution

129. The number of tangents to $y^2 = 2x$ through (1,2) is

A. 0

B. 1

C. 2

D. 3

Answer: C



Watch Video Solution

130. The number of tangents to $y^2 = 6x$ through $(-1,-1)$ is

A. 0

B. 1

C. 2

D. 3

Answer: C



Watch Video Solution

131. If the ends of a focal chord of the parabola $y^2 = 4ax$ are (x_1, y_1) and (x_2, y_2) then $x_1x_2 + y_1y_2 =$

A. a^2

B. $-3a^2$

C. $5a^2$

D. $-5a^2$

Answer: B



[Watch Video Solution](#)

132. If (x_1, y_1) and (x_2, y_2) are the end points of a focal chord of the parabola $y^2 = 5x$, then $4x_1x_2 + y_1y_2 =$

A. 25

B. 5

C. 0

D. $\frac{5}{4}$

Answer: C



Watch Video Solution

133. The point of intersection of the tangents at t_1 and t_2 to the parabola $y^2 = 12x$ is

A. $(2t_1, t_2, 2[t_1 - t_2])$

B. $(3t_1, 3[t_1 - t_2])$

C. $(3t_1, t_2, 3[t_1 - t_2])$

D. $(2t_1, t_2, 3[t_1 - t_2])$

Answer: C



Watch Video Solution

134. The slope of a chord of the parabola $y^2 = 4ax$ which is normal at one end and which subtends a right angle at the origin is

A. $1/\sqrt{2}$

B. $\sqrt{2}$

C. 2

D. none

Answer: B



[View Text Solution](#)

135. On the parabola $y^2 = 8x$ if one extremity of a focal chord is $(\frac{1}{2}, -2)$ then its other extremity is

A. (2,2)

B. $(\frac{1}{8}, -8)$

C. $(8, \frac{1}{8})$

D. (8,8)

Answer: D



[Watch Video Solution](#)

136. A focal chord of the parabola $y^2 = 4ax$ meets it at P and Q . If S is the focus then $\frac{1}{SP} + \frac{1}{SQ} =$

A. a

B. $1/a$

C. $2a$

D. $2/a$

Answer: B



[View Text Solution](#)

137. The latusrectum of a parabola whose focal chord is PSQ such that $SP = 3$ and $SQ = 2$ is given by

A. $24/5$

B. $12/5$

C. $6/5$

D. none

Answer: A



Watch Video Solution

138. The circle described on any focal chord of a parabola as diameter touches the

A. axes

B. directrix

C. parabola

D. none

Answer: B



[View Text Solution](#)

139. The circle on a focal radius as diameter of a parabola

$$y^2 = 4ax \text{ touches}$$

A. directrix

B. axis

C. tangent at the vertex

D. none

Answer: C





[View Text Solution](#)

140. A circle of radius 4 drawn on a chord of the parabola $y^2 = 8x$ as diameter touches the axis of the parabola. Then the slope of the chord is

A. $1/2$

B. $3/4$

C. 1

D. 2

Answer: C



[Watch Video Solution](#)

141. The slopes of the focal chords of the parabola $y^2 = 32x$ which are tangents to the circle $x^2 + y^2 - 4$ are

A. $\frac{1}{\sqrt{3}}, \frac{-1}{\sqrt{3}}$

B. $\frac{1}{\sqrt{15}}, \frac{-1}{\sqrt{15}}$

C. $\frac{2}{\sqrt{5}}, \frac{-2}{\sqrt{5}}$

D. $\frac{1}{\sqrt{2}}, \frac{-1}{\sqrt{2}}$

Answer: B



Watch Video Solution

142. The locus of the midpoints of the focal chords of the parabola $y^2 = 4ax$ is

A. $4a \sin^2 \theta$

B. $4a \cos^2 \theta$

C. $4a \sec^2 \theta$

D. $4a \csc^2 \theta$

Answer: C



Watch Video Solution

143. If a chord of the parabola $y^2 = 4x$ passes through its focus and makes an angle θ with the X-axis then its length is

A. $4 \cos^2 \theta$

B. $4 \sin^2 \theta$

C. $4 \cos e c^2 \theta$

D. $\sec^2 \theta$

Answer: C



Watch Video Solution

144. The length of the chord of the parabola $x^2 = 4ay$ passing through the vertex and having slope $\tan \alpha$ is

A. $4a \cos e c \alpha \cot \alpha$

B. $4a \tan \alpha \sec \alpha$

C. $4a \cos \alpha \cot \alpha$

D. $4a \sin \alpha \tan \alpha$

Answer: B



Watch Video Solution

145. In the parabola $y^2 = 4ax$ the length of the chord passing through the vertex and inclined to the axis at $\pi/4$

A. $4a\sqrt{2}$

B. $2a\sqrt{2}$

C. $a\sqrt{2}$

D. none

Answer: A



Watch Video Solution

146. The length of chord intercepted by the parabola $y = x^2 + 3x$ on the line $x+y=5$ is

A. $3\sqrt{26}$

B. $2\sqrt{26}$

C. $6\sqrt{2}$

D. none

Answer: C

 [Watch Video Solution](#)

147. If the line $y = mx + a$ meets the parabola $x^2 = 4ay$ in two points whose abscissa are x_1 and x_2 then $x_1 + x_2 = 0$ If

A. $m=-1$

B. $m=1$

C. $m=2$

D. $m=-1/2$

Answer: C



Watch Video Solution

148. Prove that the portion of the tangent intercepted between the point of contact and the directrix of the parabola $y^2 = 4ax$ subtends a right angle at its focus.

A. 30°

B. 45°

C. 60°

D. 90°

Answer: D



Watch Video Solution

149. The subnormal of the parabola $y^2 = 4ax$ is equal to

A. focus

B. vertex

C. end of the latusrectum

D. none

Answer: A



Watch Video Solution

150. If a normal chord of a point on the parabola $y^2 = 4ax$, subtends a right angle at the vertex, then $t =$

A. $4at + n = 0$

B. $4at + 4am + n = 0$

C. $4am + n = 0$

D. $at + n = 0$

Answer: A



Watch Video Solution

151. If the chord $y = mx + c$ subtends a right angle at the vertex of the parabola $y^2 = 4ax$ then the value of c is

A. $-4am$

B. $4am$

C. $-2am$

D. $2am$

Answer: A

 [Watch Video Solution](#)

152. If P is a point on the parabola $y^2 = 8x$ and A is the point $(1,0)$ then the locus of the midpoint of the line segment AP is

A. $y^2 = 4\left(x - \frac{1}{2}\right)$

B. $y^2 = 2(2x + 1)$

C. $y^2 = x - \frac{1}{2}$

D. $y^2 = 2x + 1$

Answer: A



Watch Video Solution

153. The locus of the point of intersection of two tangents to the parabola $y^2 = 4ax$ which make complementary angles with the axis of the parabola is

A. $(y^2 - 4ax)(y^2 + 4a^2) + 4a^2l^2 = 0$

B. $(y^2 - 4ax)(y^2 + 4a^2) - 4a^2l^2 = 0$

$$C. (y^2 - 4ax)(y^2 - 4a^2) + 4a^2l^2 = 0$$

$$D. (y^2 - 4ax)(y^2 - 4a^2) - 4a^2l^2 = 0$$

Answer: A



Watch Video Solution

154. The tangents at the points $(at_1^2, 2at_1)$, $(at_2^2, 2at_2)$ on the parabola $y^2 = 4ax$ are at right angles if

A. $t_1 = t_2$

B. $t_1 = -t_2$

C. $t_1t_2 = 2$

D. $t_1t_2 = -1$

Answer: B



Watch Video Solution

155. The tangent to $y^2 = ax$ makes an angle 45° with x-axis

. Then its point of contact is

A. A.P.

B. G.P.

C. H.P.

D. none

Answer: B



Watch Video Solution

156. The tangents at the points $(at_1^2, 2at_1)$, $(at_2^2, 2at_2)$ on the parabola $y^2 = 4ax$ are at right angles if

A. $t_1t_2 = -1$

B. $t_1t_2 = 1$

C. $t_1t_2 = 2$

D. $t_1t_2 = -2$

Answer: A

 [Watch Video Solution](#)

157. If P $(at_1^2, 2at_1)$ and Q $(at_2^2, 2at_2)$ are two variable points on the curve $y^2 = 4ax$ and PQ subtends a right angle at the vertex then $t_1t_2 =$

A. -1

B. -2

C. -3

D. -4

Answer: D



Watch Video Solution

158. The length of the chord of contact of tangents drawn from (x_1, y_1) to the parabola $y^2 = 4ax$ is

A. $\sqrt{(y_1^2 - 4ax_1)(y_1^2 + 4a^2)} / a$

B. $\sqrt{(y_1^2 - 4ax_1)} / a$

C. $\sqrt{(y_1^2 + 4ax_1)(y_1^2 - 4a^2)} / a$

D. $\sqrt{(y_1^2 - 4ax_1)(y_1^2 - 4a^2)} / a$

Answer: A



View Text Solution

159. The area of the triangle formed by the tangents and chord of contact from (x_1, y_2) to the parabola $y^2 = 4ax$ is

A. $(y_1^2 - 4ax_1)^{3/2}$

B. $2a(y_1^2 - 4ax_1)^{3/2}$

C. $(y_1^2 - 4ax_1)^{3/2}$

D. none

Answer: C



[View Text Solution](#)

160. If y_1, y_2 and y_3 are the ordinates of the vertices of a triangle inscribed in the parabola $y^2 = 4ax$ then its area is

A. $\frac{1}{2a}(y_1 - y_2)(y_2 - y_3)(y_3 - y_1)$

B. $\frac{1}{4a}(y_1 - y_2)(y_2 - y_3)(y_3 - y_1)$

C. $\frac{1}{8a}(y_1 - y_2)(y_2 - y_3)(y_3 - y_1)$

D. none

Answer: C

[Watch Video Solution](#)

161. The area of the triangle inscribed in the parabola $y^2 = 4x$ the ordinates of whose vertices are 1, 2 and 4 is

A. $\frac{7}{2}$ sq.unit

B. $\frac{5}{2}$ sq.unit

C. $\frac{3}{2}$ sq.unit

D. $\frac{3}{4}$ sq.unit

Answer: D

 [Watch Video Solution](#)

162. The tangents to the parabola $y^2 = 4ax$ at $P(t_1)$ and $Q(t_2)$ intersect at R. The area of ΔPQR is

A. $\frac{1}{2}a^2(t_1 - t_2)^2$

B. $\frac{1}{2}a^2(t_1 - t_2)$

C. $\frac{1}{2}a^2(t_1 - t_2)^3$

D. none

Answer: C



Watch Video Solution

163. The orthocentre of the triangle formed by three tangents to the parabola $y^2 = 4ax$ lies on the

A. axis

B. directrix

C. parabola

D. latus rectum

Answer: B



View Text Solution

164. Prove that the orthocentre of the triangle formed by any three tangents to a parabola lies on the directrix of the parabola

A. vertex

B. focus

C. foot of the directrix

D. none

Answer: B



Watch Video Solution

165. The feet of the perpendiculars drawn from the focus of a parabola to the sides of the triangle formed by its tangents lie on

A. x-axis

B. y-axis

C. directrix

D. tangent at the vertex

Answer: D



View Text Solution

166. If the distances of two points P and Q on the parabola $y^2 = 4ax$ from the focus of a parabola are 4 and 9 respectively then the distance of the point of intersection of tangents at P and Q from the focus is

A. ST

B. 2ST

C. ST^2

D. $2ST^2$

Answer: C



Watch Video Solution

167. If the distances of two points P and Q on the parabola $y^2 = 4ax$ from the focus of a parabola are 4 and 9 respectively then the distance of the point of intersection of tangents at P and Q from the focus is

A. 8

B. 6

C. 5

D. 13

Answer: B



Watch Video Solution

168. If the distances of two points P and Q on the parabola $y^2 = 4ax$ from the focus of a parabola are 4 and 9 respectively then the distance of the point of intersection of tangents at P and Q from the focus is

A. $\angle TSP = \angle TSQ$

B. $\angle TSP < \angle TSQ$

C. $\angle TSP > \angle TSQ$

D. none

Answer: A



Watch Video Solution

169. PSQ is a focal chord of a parabola whose focus is S and vertex A . PA and QA are produced to meet the directrix in R and T respectively . Then $\angle RST =$

A. 90°

B. 60°

C. 45°

D. 30°

Answer: A



[View Text Solution](#)

170. If L,M,N are the three points on the parabola $y^2 = 4ax$ whose ordinates are in G.P then the tangents at L and N

meet on the

- A. parabola
- B. abscissa of M
- C. ordinate of M
- D. none

Answer: B



[View Text Solution](#)

171. The equation of the normal to the parabola $y^2 = 8x$ at the point t is

A. $y - x = t + 2t^2$

B. $y + tx = 4t + 2t^3$

C. $x + ty = t + 2t^2$

D. $y - x = 2t - 3t^3$

Answer: B



Watch Video Solution

172. The slope of the normal at $(at^2, 2at)$ of the parabola $y^2 = 4ax$ is

A. $1/t$

B. t

C. $-t$

D. $-1/t$

Answer: C



Watch Video Solution

173. If the normal at t_1 on the parabola $y^2 = 4ax$ meet it again at t_2 on the curve then $t_1(t_1 + t_2) + 2 =$

A. t

B. $-t - 1/t$

C. $-t - 2/t$

D. none

Answer: C



Watch Video Solution

174. If the normal at t_1 on the parabola $y^2 = 4ax$ meet it again at t_2 on the curve then $t_1(t_1 + t_2) + 2 =$

A. 0

B. 1

C. t_1

D. t_2

Answer: A

 [Watch Video Solution](#)

175. If the normal at (1,2) on the parabola $y^2 = 4x$ meets the parabola again at the point $(t^2, 2t)$ then the value of t is

A. 1

B. 3

C. -3

D. 1

Answer: C



Watch Video Solution

176. If the normal to the parabola $y^2 = 4x$ at $P(1,2)$ meets the parabola again in Q then $Q=$

A. (-6,9)

B. (9,-6)

C. (-9,-6)

D. (-6,-9)

Answer: B



Watch Video Solution

177. If the normals at the point t_1 and t_2 on $y^2 = 4ax$ intersect at the point t_3 on the parabola then $t_1 t_2 =$

A. 1

B. 2

C. t_3

D. $2t_3$

Answer: B





[View Text Solution](#)

178. The number of normals drawn to the parabola $y^2 = 4x$ from the point $(1,0)$ is

A. 0

B. 1

C. 2

D. 3

Answer: B



[Watch Video Solution](#)

179. The number of normals that can be drawn through $(-1,4)$ to the parabola $y^2 - 4x + 6y = 0$ are

A. 4

B. 3

C. 2

D. 1

Answer: D



Watch Video Solution

180. From a point $(C, 0)$ three normals are drawn to the parabola $y^2 = x$. Then

A. $C < \frac{1}{2}$

B. $C = \frac{1}{2}$

C. $C > \frac{1}{2}$

D. $\frac{1}{2} > C > \frac{1}{4}$

Answer: C



Watch Video Solution

181. If the tangents and normals at the extremities of a focal chord of a parabola intersect at (x_1, y_1) and (x_2, y_2) respectively then

A. $x_1 = x_2$

B. $x_1 = y_2$

C. $y_1 = y_2$

D. $x_2 = y_1$

Answer: C



View Text Solution

182. The normals at three points P,Q,R of the parabola $y^2 = 4ax$ meet in (h,k) . The centroid of triangle PQR lies on

A. $x=0$

B. $y=0$

C. $x=-0$

D. $y=a$

Answer: B



[View Text Solution](#)

183. The ordinate of the centroid of the triangle formed by conormal points on the parabola $y^2 = 4ax$ is

A. 4

B. 0

C. 2

D. 1

Answer: B



[View Text Solution](#)

184. The normals at two points P and Q of a parabola $y^2 = 4ax$ meet at (x_1, y_1) on the parabola. Then $PQ^2 =$

A. $(x_1 + 4a)(x_1 + 8a)$

B. $(x_1 + 4a)(x_1 - 8a)$

C. $(x_1 - 4a)(x_1 + 8a)$

D. $(x_1 - 4a)(x_1 - 8a)$

Answer: B



[View Text Solution](#)

185. If a normal subtends a right angle at the vertex of a parabola $y^2 = 4ax$ then its length is

A. $\sqrt{5} a$

B. $3\sqrt{5}a$

C. $6\sqrt{3}a$

D. $7\sqrt{5}a$

Answer: C



Watch Video Solution

186. If α is the inclination of a tangent to the parabola $y^2 = 4ax$ then the distance between the tangent and a parallel normal is

A. $a \operatorname{cosec} \alpha \sec \alpha$

B. $a \operatorname{cosec} \alpha \sec^2 \alpha$

C. $a \cos e c^2 \alpha \sec \alpha$

D. $a \cos e c^2 \alpha \sec^2 \alpha$

Answer: D



View Text Solution

187. The length of the normal chord drawn at one end of the latus rectum of $y^2 = 4ax$ is

A. $2\sqrt{2} a$

B. $4\sqrt{2}a$

C. $8\sqrt{2}a$

D. $10\sqrt{2}a$

Answer: C



[View Text Solution](#)

188. The locus of a point that divides chords of slope 2 of the parabola $y^2 = 4x$ internally in the ratio 1:2 is a parabola .

Then the vertex is

A. $(2/9, 8/9)$

B. $(3/7, 5/7)$

C. $(-2/9, 8/9)$

D. $(1/9, 4/9)$

Answer: A



[View Text Solution](#)

189. Let O be the vertex and Q be any point on the parabola $x^2 = 8y$. If the point P divides the line segment OQ internally in the ratio 1: 3 then the locus of P is:

A. $x^2 = y$

B. $y^2 = x$

C. $y^2 = 2x$

D. $x^2 = 2y$

Answer: D



Watch Video Solution

190. If a normal chord of a point on the parabola $y^2 = 4ax$, subtends a right angle at the vertex, then $t =$

A. 4

B. 2

C. 1

D. 3

Answer: B

 [Watch Video Solution](#)

191. If a normal subtends a right angle at the vertex of a parabola $y^2 = 4ax$ then its length is

A. $\sqrt{2}$

B. 2

C. $\sqrt{3}$

D. 3

Answer: A



Watch Video Solution

192. The normal at 'P' cuts the axis of the parabola $y^2 = 4ax$ in G and S is the focus of the parabola. If $\triangle SPG$ is equilateral then each side is of length.

A. SP

B. 2SP

C. $\frac{1}{2}$ SP

D. none

Answer: A



Watch Video Solution

193. The circle passing through three conormal points also passes through

A. vertex

B. foot of the directrix

C. focus

D. none

Answer: A



[View Text Solution](#)

194. The normal at 'P' cuts the axis of the parabola $y^2 = 4ax$ in G and S is the focus of the parabola. If $\triangle SPG$ is equilateral then each side is of length.

- A. a
- B. 2a
- C. 3a
- D. 4a

Answer: D



[Watch Video Solution](#)

195. If the normals at two points on the parabola $y^2 = 4ax$ intersect on the parabola then the product of the abscissae is

A. $4a^2$

B. $-4a^2$

C. $2a$

D. $4a^4$

Answer: A



Watch Video Solution

196. If the normals at two points on the parabola intersect on the curve then the product of the ordinates of the points is

A. $8a$

B. $8a^2$

C. $8a^3$

D. $8a^4$

Answer: B



[View Text Solution](#)

197. The locus of the point of intersection of perpendicular tangents to the parabola $y^2 = 4ax$ is

A. $y^2 = a(x - 3a)$

B. $y^2 = a(x + 3a)$

C. $y^2 = a(x + 2a)$

D. $y^2 = a(x - 2a)$

Answer: A



Watch Video Solution

198. The three normals from a point to the parabola $y^2 = 4ax$ cut the axes in points whose distance from vertex are in in A.R then the loous of the point is

A. $27ay^2 = 2(x - 2a)^3$

B. $27ay^3 = 2(x - 2a)^2$

C. $9ay^2 = 2(x - 2a)^3$

D. $9ay^3 = 2(x - 2a)^2$

Answer: A



[View Text Solution](#)

199. If the normals from any point to the parabola $x^2 = 4y$ cuts the line $y=2$ in points whose abscissae are in A.P., then the slope of the tangents at the 3 conormal points are in

A. AP

B. GP

C. HP

D. none

Answer: A



[View Text Solution](#)

200. If a circle cuts the parabola $y^2 = 4ax$ in four points then the algebraic sum of ordinates of the four points is

A. 0

B. 1

C. -1

D. none

Answer: A



[View Text Solution](#)

201. The feet of the normals to $y^2 = 4ax$ from the point $(6a,0)$ are

A. $(0,0)$

B. $(4a,4a)$

C. $(4a,-4a)$

D. $(0,0),(4a,4a),(4a,-4a)$

Answer: D

 [Watch Video Solution](#)

202. If $P(-3, 2)$ is one end of focal chord PQ of the parabola

$y^2 + 4x + 4y = 0$ then slope of the normal at Q is

A. $(-1/2)$

B. 2

C. $1/2$

D. -2

Answer: A



Watch Video Solution

203. The normal at a point P on the parabola $y^2 = 4ax$ cuts the curve again at Q . If M is the midpoint of PQ then the product of the ordinates of P and M is

A. a^2

B. $2a^2$

C. $4a^2$

D. $-4a^2$

Answer: D



[View Text Solution](#)

204. The subnormal of the parabola $y^2 = 4ax$ is equal to

A. latus rectum

B. semi latus rectum

C. 2(latus rectum)

D. none

Answer: B

 [Watch Video Solution](#)

205. The length of the subnormal to the curve $y^2 = 2px$ is

A. p

B. $p/2$

C. $2p$

D. $4p$

Answer: A

 [Watch Video Solution](#)

206. If P is a point on the parabola $y^2 = 4ax$ such that the subtangent and subnormal at p are equal then the

coordinates of P are

A. $(a, 2a)$ or $(a, -2a)$

B. $(2a, 2\sqrt{2}a)$

C. $(4a, -4a)$ or $(4a, 4a)$

D. none

Answer: A



[View Text Solution](#)

207. If PSP^1 is a focal chord of a parabola $y^2 = 4ax$ and SL is its semi latusrectum then SP SL and SP^1 are in

A. A.P.

B. H.P

C. G.P

D. none of these

Answer: B



Watch Video Solution

208. An arch is in the shape of a parabola whose axis is vertically downwards and measures 24 mts across its bottom on the ground. Its highest point is 24 mts. The measure of the horizontal beam across its cross section at a height of 18 mts is

A. 50 mt

B. 40 mt

C. 45 mt

D. 55 mt

Answer: B



Watch Video Solution

209. The points on $y = x^2 + 7x + 2$ which is closest to the line $y=3x-3$ is

A. (2,8)

B. (2,-8)

C. (-2,8)

D. (-2,-8)

Answer: D



Watch Video Solution

210. Let P be the point on the parabola $y^2 = 8x$ which is at a minimum distance from the centre C of the circle $x^2 + (y + 6)^2 = 1$. Then the equation of the circle passing through C and having its centre at P is

A. $x^2 + y^2 - 4x + 8y + 12 = 0$

B. $x^2 + y^2 - x + 4y - 12 = 0$

C. $x^2 + y^2 - \frac{x}{4} + 2y - 24 = 0$

D. $x^2 + y^2 - 4x + 9y + 18 = 0$

Answer: A

 [View Text Solution](#)

Exercise 1 B

1. The chord of contact of $(2,1)$ w.r.t the parabola $x^2 = y$ is

A. $x+4y+3=0$

B. $2x-3y+4=0$

C. $3x+2y+4=0$

D. $4x-y-1=0$

Answer: D

 [Watch Video Solution](#)

2. The polar of $(-2,3)$ w.r.t the parabola $y^2 = 4x$ is

A. $2x-3y-4=0$

B. $2x-y-2=0$

C. $3x-y+4=0$

D. $5x-4y+24=0$

Answer: A



Watch Video Solution

3. The polar of $(a,0)$ w.r.t the parabola $y^2 = 4ax$ is

A. $x=a$

B. $x+a=0$

C. $y=a$

D. $y+a=0$

Answer: B



Watch Video Solution

4. The pole of the line $2x+3y-4 = 0$ with respect to the parabola $y^2 = 4x$ is

A. (2,3)

B. (-2,-3)

C. (1,1)

D. (2,-3)

Answer: B



Watch Video Solution

5. The pole of the straight line $x - 2y + 4 = 0$ with respect to the parabola $y^2 = 6x$ is

A. (4,6)

B. (-4,6)

C. (4,-6)

D. (-4,-6)

Answer: A



Watch Video Solution

6. The pole of the line $3x+4y-4=0$ w.r.t parabola $x^2 = 4y$ is

A. $(3/2,1)$

B. $(3/2,-1)$

C. $(-3/2,1)$

D. $(-3/2,-1)$

Answer: D



Watch Video Solution

7. The points $(3,-2)$, $(1,-2)$ are conjugate w.r.t. the parabola

A. $y^2 = 2x$

B. $y^2 = 4x$

C. $y^2 = 8x$

D. none

Answer: A



Watch Video Solution

8. The lines $3x+2y-1=0$, $2x-y-2 =0$ are conjugate w.r.t the parabola

A. $y^2 = 8x$

B. $y^2 = x$

C. $y^2 = 2x$

D. none

Answer: A



Watch Video Solution

9. If the points $(2,4)$, $(k,6)$ are conjugate with respect to the parabola $y^2 = 4x$ then $k =$

A. 10

B. $7/2$

C. -12

D. -2

Answer: A



Watch Video Solution

10. If the lines $2x+3y+12=0$ and $x-y+k=0$ are conjugate with respect to the parabola $y^2 = 8x$ then $k=$

A. 10

B. $7/2$

C. -12

D. -2

Answer: C



[Watch Video Solution](#)

11. If the lines $2x+3y+12=0$ and $x-y+4k=0$ are conjugate with respect to the parabola $y^2 = 8x$ then the value of k is

A. -3

B. 3

C. 2

D. -2

Answer: A



Watch Video Solution

12. The polar of $(-a, -2a)$ w.r.t the circle

$x^2 + y^2 - 2ax - 3a^2 = 0$ touches the parabola

A. $y^2 = 4ax$

B. $y^2 = 6ax$

C. $x^2 = 4ax$

D. $y^2 = ax$

Answer: A



Watch Video Solution

13. The locus of poles of tangents of the parabola $y^2 = 4ax$ w.r.t the parabola $y^2 = 4bx$ is

A. $ax^2 = 4b^2y$

B. $ax = 4b^2y^2$

C. $ay^2 = 4b^2x$

D. none

Answer: C



[View Text Solution](#)

14. If the polar of a point P w.r.t the circle $x^2 + y^2 = a^2$ touches the parabola $y^2 = 4ax$ then the locus of P is

A. $y^2 = ax$

B. $y^2 + ax = 0$

C. $y^2 = 2ax$

D. $y^2 + 2ax = 0$

Answer: B

[Watch Video Solution](#)

15. The locus of the midpoints of the focal chords of the parabola $y^2 = 4ax$ is

A. $x+a=0$

B. $x+2a=0$

C. $x+3a=0$

D. $x+4a=0$

Answer: D

 [Watch Video Solution](#)

16. If the polar of a point P w.r.t the circle $x^2 + y^2 = a^2$ touches the parabola $y^2 = 4ax$ then the locus of P is

A. $x^2 - y^2 = 4a^2$

B. $x^2 - y^2 = 2a^2$

C. $x^2 - y^2 = a^2$

D. none

Answer: A



Watch Video Solution

17. The chord of contact of a point P to the parabola

$y^2 = 4ax$ touch the circle $x^2 + y^2 = r^2$. The locus of P is

A. $4a^2x^2 = r^2(y^2 + 4a^2)$

B. $a^2x^2 = 2r^2(y^2 - 4a^2)$

C. $2a^2x^2 = 2r^2(y^2 + 2a^2)$

$$D. 4a^2x^2 = r^2(y^2 + 4a^2)$$

Answer: A



[View Text Solution](#)

18. The locus of the point for which the chord of contact w.r.t $y^2 = 4ax$ subtends a right angle at the vertex of the parabola is

A. $x+2a=0$

B. $x+4a=0$

C. $y+2a=0$

D. $y+4a=0$

Answer: B



View Text Solution

19. The locus of poles of chords of the parabola $y^2 = 4ax$ which are at a constant distance d from the vertex is

A. $d^2x^2 + 4a^2(d^2 - y^2) = 0$

B. $d^2y^2 + 4a^2(d^2 - x^2) = 0$

C. $d^2y^2 + 2a^2(3d^2 - 2x^2) = 0$

D. $x^2 + 2a^2(d^2 + 2x^2) = 0$

Answer: C



View Text Solution

20. A chord of the parabola $y^2 = 4ax$ subtends a right angle at the vertex. The tangents at the extremities of the chord intersect on

A. $x+a=0$

B. $x+2a=0$

C. $x+4a=0$

D. none

Answer: C



[View Text Solution](#)

21. The equation of the chord of the parabola $y^2 = 2x$ having (1,1) as its midpoint is

A. $x+y=0$

B. $x-y=0$

C. $x-y+1=0$

D. $2x-y=0$

Answer: B



Watch Video Solution

22. The midpoint of the chord $2x-y-2=0$ of the parabola $y^2 = 8x$ is

A. (1,0)

B. (2,2)

C. (3,4)

D. (0,-2)

Answer: B



Watch Video Solution

23. The tangent at the point P (x_1, y_1) to the parabola $y^2 = 4ax$ meets the parabola $y^2 = 4a(x + b)$ at Q and R then the midpoint of QR is

A. (2,4)

B. (4,2)

C. (7,9)

D. none

Answer: A



[View Text Solution](#)

24. If the tangent at the point P (2,4) to the parabola $y^2 = 8x$ meets the parabola $y^2 = 8x + 5$ at Q and R then the midpoint of QR is

A. (2,4)

B. (4,2)

C. (7,9)

D. none

Answer: A



[View Text Solution](#)

25. The locus of the midpoints of the focal chords of the parabola $y^2 = 4ax$ is

A. $y^2 = 8ax$

B. $y^2 = 4ax$

C. $y^2 = 2ax$

D. $y^2 = ax$

Answer: C



Watch Video Solution

26. Let O be the origin and A be a point on the curve $y^2 = 4x$. Then the locus of the midpoint of OA is

A. $x^2 = 4y$

B. $x^2 = 2y$

C. $y^2 = 16x$

D. $y^2 = 2x$

Answer: D



Watch Video Solution

27. The locus of the midpoints of the focal chords of the parabola $y^2 = 4ax$ is

A. $y^2 = 2a(x + a)$

B. $y^2 = 2a(x - a)$

C. $y^2 = a(2x + a)$

$$D. y^2 = a(2x - a)$$

Answer: B

 [Watch Video Solution](#)

28. The locus of the midpoints of the focal chords of the parabola $y^2 = 6x$ which pass through a fixed point (9,5) is

A. $y^2 + 5y + 3x + 27 = 0$

B. $y^2 + 5y - 3x + 27 = 0$

C. $y^2 - 5y - 3x + 27 = 0$

D. $y^2 - 5y - 3x - 27 = 0$

Answer: C



[Watch Video Solution](#)

29. The locus of middle points of normal chords of the parabola $y^2 = 4ax$ is

A. $y(y + k) = 2a(x + h)$

B. $y(y - k) = 2a(x - h)$

C. $y(y - h) = 2a(x + h)$

D. $y(y + k) = 2a(x - h)$

Answer: B

[Watch Video Solution](#)

30. The point of intersection of the tangents of the parabola $y^2 = 16x$ at the extremities of the chord having (3,4) as its midpoint is

A. (1,4)

B. (-1,4)

C. (1,-4)

D. (-1,-4)

Answer: B



[View Text Solution](#)

31. The locus of the midpoints of the focal chords of the parabola $y^2 = 4ax$ is

A. $y^2 = 2a(x + 4a)$

B. $y^2 = 2a(x - 4a)$

C. $y^2 = a(x + 2a)$

D. $y^2 = a(x - 2a)$

Answer: B



Watch Video Solution

32. A variable tangent to the parabola $y^2 = 4ax$ meets the parabola $y^2 + 4ax = 0$ at the points P, Q . The locus of the middle point of PQ is

A. $y^2 + 4ax = 0$

B. $y^2 + 2ax = 0$

$$C. y^2 + ax = 0$$

$$D. 3y^2 + 4ax = 0$$

Answer: D



View Text Solution

33. A tangent to the parabola $y^2 + 4bx = 0$ meets the parabola $y^2 = 4ax$ in P and Q . The locus of the middle point of PQ is

$$A. y^2(2a + b) = 4a^2x$$

$$B. y^2(2a - b) = 4a^2x$$

$$C. y^2(2a + b) = 4ax$$

$$D. y^2(2a - b) = 4ax$$

Answer: A



View Text Solution

34. The locus of midpoints of chords of the parabola

$y^2 = 4ax$ which touch the circle $x^2 + y^2 = a^2$ is

A. $(y^2 - 2ax)^2 = a^2(y^2 + 4a^2)$

B. $(y^2 + 2ax)^2 = a^2(y^2 + 4a^2)$

C. $(y^2 + 2ax)^2 = a^2(y^2 - 4a^2)$

D. $(y^2 - 2ax)^2 = a^2(y^2 - 4a^2)$

Answer: A



View Text Solution

35. The locus of the midpoints of the chords of the parabola

$y^2 = 6x$ which touch the circle $x^2 + y^2 + 4x - 12 = 0$ is

A. $(y^2 - 3x - 6)^2 = 16(y^2 + 9)$

B. $(x^2 - 3y - 16)^2 = 16(y^2 + 19)$

C. $8(y^2 - 3x - 6)^2 = 16(y^2 + 9)$

D. $2(y^2 - 3x - 6)^2 = 16(y^2 - 9)$

Answer: A



[View Text Solution](#)

36. The locus of midpoints of chords of the parabola

$y^2 = 4ax$ which are parallel to line $y = mx + c$ is

A. $x=2a$

B. $x=2a/m$

C. $y=2a$

D. $y=2a/m$

Answer: D



Watch Video Solution

37. The tangent at 't' on the parabola $y^2 = 4ax$ is parallel to a normal chord then distance between them is

A. a

B. $2a$

C. $4a$

D. 8a

Answer: C



Watch Video Solution

38. An equilateral triangle is inscribed in the parabola $y^2 = 4ax$ whose vertex is at the vertex of the parabola . The length of its side is

A. $2\sqrt{3}a$

B. $4\sqrt{3}a$

C. $8\sqrt{3}a$

D. $16\sqrt{3}a$

Answer: C



[View Text Solution](#)

39. The equation to the pair of tangents drawn from (3,-2) to the parabola $y^2 = x$ is

A. $x^2 + 8xy + 12y^2 + 10x + 24y + 9 = 0$

B. $2x^2 + 3xy - 22y^2 + 15x + 4y + 9 = 0$

C. $3x^2 + 18xy + 22y^2 + 50x + 64y + 19 = 0$

D. $x^2 - 8xy - 12y^2 - 10x - 24y + 9 = 0$

Answer: A



[View Text Solution](#)

40. The combined equation to the tangents to the parabola

$y^2 = 4ax$ from an external point A (x_1, y_1) is

A. $(y^2 - 4ax)(y_1^2 - 4ax_1) = (yy_1 - 2ax - 2ax_1)^2$

B. $y^2 - 4ax = (yy_1 - 2ax - 2ax_1)^2$

C. $y^2 - 4ax = (yy_1 - 2ax)^2$

D. none of these

Answer: A

 [View Text Solution](#)

41. Two tangents are drawn from a point $(-2, -1)$ to the

curve $y^2 = 4x$, If α is the angle between them, then

$|\tan \alpha|$ is equal to

A. 3

B. $1/3$

C. 1

D. $1/2$

Answer: A



Watch Video Solution

42. The angle between the tangents to the parabola $y^2 = 4ax$ at the points where it intersects with the line $x - y - a = 0$ is

A. $\pi/4$

B. $\pi/2$

C. $\pi / 3$

D. $\pi / 6$

Answer: B



Watch Video Solution

43. The angle between the tangents to the parabola $y^2 = 4ax$ at the points where it intersects with the line $x - y - a = 0$ is

A. $\pi / 3$

B. $\pi / 4$

C. $\pi / 6$

D. $\pi / 2$

Answer: D



Watch Video Solution

44. The angle between the tangents drawn from the origin to the parabola $y^2 = 4a(x - a)$ is

A. $\pi / 6$

B. $\pi / 4$

C. $\pi / 3$

D. $\pi / 2$

Answer: D



View Text Solution

45. The slope of tangents drawn from a point (4,10) to the parabola $y^2 = 9x$ are

A. $x - y + 1 = 0, x - 2y + 4 = 0$

B. $x - y - 1 = 0, 2x - y - 4 = 0$

C. $x - 4y + 36 = 0, 9x - 4y + 4 = 0$

D. $x + y + 5 = 0, 2x - 2y - 14 = 0$

Answer: C

 [Watch Video Solution](#)

46. Find the angle between tangents drawn from P(2, 3) to the parabola $y^2 = 4x$

A. $x + y + 1 = 0, x + 2y + 4 = 0$

B. $x - y + 1 = 0, x - 2y + 4 = 0$

C. $x + y - 1 = 0, x - 2y + 4 = 0$

D. $x - y - 1 = 0, 2x - y - 4 = 0$

Answer: D



Watch Video Solution

47. The locus of middle points of normal chords of the parabola $y^2 = 4ax$ is

A. $\frac{y^2}{2a} + \frac{4a^3}{y^2} - x = 2a$

B. $\frac{y^2}{2a} - \frac{4a^3}{y^2} - x = 2a$

C. $\frac{y^2}{2a} + \frac{4a^3}{y^2} - x + 2a$

D. $\frac{y^2}{2a} - \frac{4a^3}{y^2} - x + 2a$

Answer: A



Watch Video Solution

48. The locus of the middle points of chords of the parabola which are such that the normals at their extremities meet on the parabola is

A. $y^2 = 2a(x + 2a)$

B. $y^2 = 2a(x - 2a)$

C. $y^2 = a(x + 2a)$

D. $y^2 = a(x - 2a)$

Answer: A



View Text Solution

49. The locus of middle points of normal chords of the parabola $y^2 = 4ax$ is

A. $(x + 2a)y^2 = 4a^3$

B. $(x + 2a)y^2 + 4a^3 = 0$

C. $(x - 2a)y^2 = 4a^3$

D. $(x - 2a)y^2 + 4a^3 = 0$

Answer: B



Watch Video Solution

50. The locus of the midpoints of the focal chords of the parabola $y^2 = 4ax$ is

A. $y^2 = a(x - a)$

B. $y^2 = a(x + a)$

C. $y^2 = 2a(x - a)$

D. $y^2 = 2a(x + a)$

Answer: A

 [Watch Video Solution](#)

Exercise 2 Special Type Questions Set 1

1. I : The length of the latus rectum of the parabola

$$y^2 + 8x - 2y + 17 = 0 \text{ is } 8.$$

II: The focal distance of the point (9,6) on the parabola

$$y^2 = 4x \text{ is } 12$$

A. only I is true

B. only II is true

C. both I and II are true

D. neither I nor II true

Answer: A



Watch Video Solution

2. For the parabola $y^2 + 6y - 2x + 5 = 0$

I) The vertex is $(-2, -3)$ II) The directrix is $y + 3 = 0$

Which of the following is correct ?

A. Both I and II are true

B. I is true II is false

C. I is false ,II is true

D. Both I and II are false

Answer: B

 [Watch Video Solution](#)

3. I: If the points $(2, -1)$, $(5, k)$ are conjugate with respect to the parabola $x^2 = 8y$ then $k=7$

II : If the lines $2x+3y+12=0$, $x-y+k=0$ are conjugate with respect to the parabola $y^2 = 8x$ then $k = -12$

- A. only I is true
- B. only II is true
- C. both I and II are true
- D. neither I nor II true

Answer: B

 [Watch Video Solution](#)

4. I : The locus of the midpoint of chords of the parabola $y^2 = 4ax$ which subtends a right angle at the vertex is $y^2 = 2a(x - 4a)$

II : The locus of midpoint of chords of the parabola $y^2 = 4ax$ which touch the circle $x^2 + y^2 = a^2$ is $(y^2 - 2ax)^2 = a^2(y^2 + 4a^2)$.

- A. only I is true
- B. only II is true
- C. both I and II are true
- D. neither I nor II true

Answer: C



[View Text Solution](#)

Exercise 2 Special Type Questions Set 2

1. If the equation of the parabola whose axis is parallel to x-axis and passing through $(2,-1), (6,1), (3,-2)$ is $ay^2 + bx + cy + d = 0$ then the ascending order of a,b,c,d is

A. a,b,c,d

B. b,c,a,d

C. c,a,b,d

D. b,a,c,d

Answer: D



View Text Solution

2. The equation of the directrix of the parabola whose vertex (3,2) and focus (2,-1) is

A. a,b,c

B. b,c,a

C. c,a,b

D. b,a,c

Answer: C

 [Watch Video Solution](#)

3. If the line $ax+by+c=0$ touches both the parabolas $y^2 = -32y$ then the ascending order of a,b,c is

A. a,b,c

B. b,c,a

C. c,a,b

D. b,a,c

Answer: D



View Text Solution

4. If the chord of contact of $(3,-2)$ with respect to the parabola $y^2 = x$ is $ax+by+c = 0$ then the ascending order of a,b,c is

A. a,c,b

B. b,c,a

C. c,a,b

D. b,a,c

Answer: B



Watch Video Solution

Exercise 2 Special Type Questions Set 3

1. Match the following .

Parabola

I. $y^2 - x - 3y + 2 = 0$

II. $y^2 - 8x - 4y - 4 = 0$

III. $x^2 + 4x - 8y + 28 = 0$

IV. $x^2 - 2x - 8y - 23 = 0$

Focus

(a) (1, 2)

(b) (- 2, 5)

(c) (1, - 1)

(d) (5/4, 1)

A. a,b,c,d

B. b,c,a,d

C. d,a,b,c

D. b,d,a,c

Answer: C



[View Text Solution](#)

2. Match the following .

Point, parabola

tangent

I. $(3, -6)y^2 = 12x$

$(a)x + y + 1 = 0$

II. $(2, 4)y^2 = 8x$

$(b)x + y + 3 = 0$

III. $(-2, 1)x^2 = 4y$

$(c)x - y + 2 = 0$

A. a,b,c

B. b,c,a

C. c,a,b

D. b,a,c

Answer: B



Watch Video Solution

3. Match the following

Point, parabola

I. $(3, -2)y^2 = x$

II. $(2, 1)x^2 = y$

III. $(-2, 3)y^2 = 4x$

IV. $(5, -6)x^2 = 8y$

Polar

(a) $5x - 4y + 24 = 0$

(b) $2x - 3y - 4 = 0$

(c) $4x - y - 1 = 0$

(c) $x + 4y + 3 = 0$

A. a,b,c,d

B. b,c,a,d

C. d,c,b,a

D. b,d,a,c

Answer: C



Watch Video Solution

4. Match the following.

Line, parabola

Pole

I. $2x - 3y + 4 = 0$ $y^2 = 4x$

(a) (4, 6)

II. $2x + 3y - 4 = 0$ $y^2 = 4x$

(b) (2, 3)

III. $x - 2y + 4 = 0$ $y^2 = 6x$

(c) (-2, -3)

IV. $3x + 4y - 4 = 0$ $x^2 = 4y$

(d) (-3/2, -1)

A. a,b,c,d

B. b,c,a,d

C. d,c,b,a

D. b,d,a,c

Answer: B



Watch Video Solution

Exercise 2 Special Type Questions Set 4

1. A: The focus of the parabola $(y - 3)^2 = 6(x + 3)$ is $(-3//2, 2)$.

R : The focus of the parabola $(y - \beta)^2 = \pm 4a(x - \alpha)$ is $(\alpha \pm a, \beta)$.

A. Both A and R are true and R is the correct explanation of A

B. Both A and R are true but R is not correct explanation of A

C. A is true but R is false

D. A is false but R is true

Answer: D



Watch Video Solution

2. A : The condition that the line $x/p+y/q=1$ to be a tangent to the parabola $y^2 = 4ax$ is $ap+q^2 = 0$.

R: The condition that the line $lx+my+n=0$ may touch the parabola $y^2 = 4ax$ is $am^2 = ln$

A. Both A and R are true and R is the correct explanation of A

B. Both A and R are true but R is not correct explanation of A

C. A is true but R is false

D. A is false but R is true

Answer: A

 [Watch Video Solution](#)

3. A : The sum and product of the slopes of the tangents to the parabola $y^2 = 8x$ drawn from the point $(-2, 3)$ are $-3/2, -1$.

R : If m_1, m_2 are the slopes of the tangents of the parabola $y^2 = 4ax$ through $P(x_1, y_1)$ then $m_1 + m_2 = y_1/x_1, m_1 m_2 = a/x_1$.

- A. Both A and R are true and R is the correct explanation of A
- B. Both A and R are true but R is not correct explanation of A
- C. A is true but R is false
- D. A is false but R is true

Answer: A



Watch Video Solution

4. Given :A circle $2x^2 + 2y^2 = 5$ and a parabola $y^2 = 4\sqrt{5}x$.

Statement -I : an equation of a common tangent to these

curves is $y = x + \sqrt{5}$.

Statement -I - If the line, $y = mx + \frac{\sqrt{5}}{m}$ ($m \neq 0$) is their common tangent ,then m satisfies $m^4 - 3m^2 + 2 = 0$

A. Statement -I is true , statement -II is false

B. Statement -I is false , Statement -II is true

C. Statement -I is true , Statement -II is true ,Statement -

II is a correct explanation for Statement -I

D. Statement -I is true , Statement -II is true ,Statement -

II is not a correct explanation for Statement -I

Answer: D



Watch Video Solution