



MATHS

BOOKS - DEEPTI MATHS (TELUGU ENGLISH)

QUADRATIC EXPRESSIONS

SOLVED EXAMPLES

1. The roots of the equation $6\sqrt{5}x^2 - 9x - 3\sqrt{5} = 0$ is

A. $\sqrt{5}/2\sqrt{5}/5$

B. $-\sqrt{5}/2\sqrt{5}/5$

C. $\sqrt{5}/2, -\sqrt{5}/5$

D. $-\sqrt{5}/2, -\sqrt{5}/5$

Answer: C



Watch Video Solution

2. IF the product of the roots of the equation $x^2 - 3kx + 2e^{2\log k} - 1 = 0$ is 17 then K=

A. 5

B. 3

C. 2

D. 9

Answer: B



Watch Video Solution

3. IF the different between the roots of $x^2 - px + q = 0$ is 2, then the relation between p , and q is

A. $p = 4(q + 1)^2$

B. $p^2 = (q + 1)$

C. $p^2 = 4(q + 1)$

D. $p = 4(q + 1)$

Answer: C



Watch Video Solution

4. if the equation $(3x)^2 + (27 \times 3^{1/p} - 15)x + 4 = 0$ has equal roots then $p =$

A. 0

B. 2

C. $-1/2$

D. none

Answer: C



Watch Video Solution

5. IF $2 + \sqrt{3}$ is a root of the equation $x^2 + px + q = 0$ then

A. $p = -6, q = 13$

B. $p = -6, q = 14$

C. $p = -4, q = 1$

D. $p = -8, q = 25$

Answer: C



[Watch Video Solution](#)

6. The number of solutions of the equation $|x^2| - 3|x| + 2 = 0$ is

A. 4

B. 1

C. 3

D. 2

Answer: A



Watch Video Solution

7. Solve $4^{x-1} - 3 \cdot 2^{x-1} + 2 = 0$

- A. $\{1, 2\}$
- B. $\{1, -1\}$
- C. $\{1, 5\}$
- D. $\{4, -1\}$

Answer: A



Watch Video Solution

8. The minimum value of $3x^2 + 2x + 11$ is

- A. 32

B. $32/3$

C. 2

D. 3

Answer: B



[Watch Video Solution](#)

9. The solution set of $x^2 - 8x + 15 > 0$ is

A. $(-1, 4)$

B. $(-\infty, -3] \cup [7, \infty)$

C. $(-\infty, 3) \cup (5, \infty)$

D. $[-4, 1]$

Answer: C



[Watch Video Solution](#)

10. The range of $\frac{x^2 - 2x + 3}{x^2 - 2x - 8}$ is

- A. $(-\infty, 0] \cup (1, \infty)$
- B. $[1/2, 2]$
- C. $(-\infty, -2/9] \cup (1, \infty)$
- D. $(-\infty, -6] \cup [-2, \infty)$

Answer: C



Watch Video Solution

11. IF $p, q, \in \{1, 2, 3, 4\}$ the number of equation of the form $px^2 + qx + 1 = 0$ having real roots is

- A. 15
- B. 9
- C. 7
- D. 8

Answer: C

 [Watch Video Solution](#)

12. The set of values of p for which roots of the equation $3x^2 + 2x + p(p - 1) = 0$ are of opposite sign is

A. $(0, \infty)$

B. $(-\infty, 0)$

C. $(0, 1)$

D. $(1, \infty)$

Answer: C

 [Watch Video Solution](#)

13. IF α and β be the roots of the equation $x^2 + px - \frac{1}{2p^2} = 0$ where $p \in R$, then the minimum value of $\alpha^4 + \beta^4 =$

A. $2 + \sqrt{2}$

B. 2

C. $2\sqrt{2}$

D. $2 - \sqrt{2}$

Answer: A



Watch Video Solution

14. Sum of the non real roots of $(X^2 + x - 2)(x^2 + x - 3) = 12$ is

A. -1

B. 1

C. -6

D. 6

Answer: A



Watch Video Solution

15. if $x^2 + 3x + 5 = 0$ and $ax^2 + bx + c = 0$ have common roots / roots and $a, b, c \in \mathbb{N}$, then the minimum value of $a + b + c$ is

A. 9

B. -9

C. 3

D. -3

Answer: A



[Watch Video Solution](#)

16. if $c > 0$ and $4a + c < 2b$ then $ax^2 - bx + c = 0$ has a root in the interval

A. (2, 4)

B. (0, 2)

C. $(0, 1)$

D. $(-2, 0)$

Answer: B



Watch Video Solution

17. If $a=c=4b$, then the roots of $ax^2 + 4bx + c = 0$ are

A. $1, \frac{c}{a}$

B. $-1, \frac{c}{a}$

C. $-1, -\frac{c}{a}$

D. $1, -\frac{c}{a}$

Answer: C



Watch Video Solution

18. IF the equation $ax^2 + 2bx + 3c = 0$ and $3x^2 + 8x + 15 = 0$ have a common root , where a,b,c are the length of the sides of a ΔABC , then $\sin^2 A + \sin^2 B + \sin^2 C =$

A. 1

B. $\frac{3}{2}$

C. $\sqrt{2}$

D. 2

Answer: D



[Watch Video Solution](#)

19. IF both the roots of equation $x^2 - 2ax + a^2 - 1 = 0$ lie in the interval $(-3, 4)$, then sum of the integral of a is

A. 0

B. 2

C. 4

D. 1

Answer: A



[Watch Video Solution](#)

EXERCISE 1A (QUADRATIC EQUATIONS)

1. If a, b are the roots of $x^2 + x + 1 = 0$ then $a^2 + b^2 =$

A. 1

B. 2

C. -1

D. 4

Answer: C



[Watch Video Solution](#)

2. IF α, β are the roots of $x^2 - x + 2 = 0$ then $\alpha^2\beta + \alpha\beta^2 =$

A. 5

B. 3

C. -2

D. 2

Answer: D



[Watch Video Solution](#)

3. If α, β are the roots of $3x^2 - 5x + 7 = 0$ then $\alpha^3 + \beta^3 =$

A. $90/7$

B. $-120/17$

C. $170/23$

D. $-190/27$

Answer: D



Watch Video Solution

4. If α, β are the roots of $ax^2 + bx + c = 0$ then $\alpha^2 + \beta^2 =$

A. $\frac{(b^2 - 2ac)}{a^2}$

B. $-\frac{b}{c}$

C. $\frac{(b^2 - 2ac)}{c^2}$

D. $\frac{(b^2 - 2ac)}{ac}$

Answer: A



Watch Video Solution

5. If α, β are the roots of $ax^2 + bx + c = 0$ then $\alpha\beta^2 + \alpha^2\beta + \alpha\beta =$

A. $\frac{ac - bc}{a^2}$

B. $\frac{bc - ac}{a^2}$

C. $\frac{ac - bc}{b^2}$

D. $\frac{bc - ac}{b^2}$

Answer: A



Watch Video Solution

6. If α, β are the roots of $x^2 + x + 1 = 0$ then $\alpha/\beta + \beta/\alpha =$

A. -1

B. 1

C. 2

D. none

Answer: A



Watch Video Solution

7. IF α, β are the roots of $ax^2 + bx + c = 0$ then $\frac{1}{\alpha^2} + \frac{1}{\beta^2} =$

A. 1

B. -1

C. 2

D. $p/q - 2$

Answer: B



Watch Video Solution

8. IF α, β are the roots of $ax^2 + bx + c = 0$ then $\frac{1}{\alpha^2} + \frac{1}{\beta^2} =$

A. $\frac{(b^2 - 2ac)}{a^2}$

B. $-\frac{b}{c}$

C. $\frac{(b^2 - 2ac)}{c^2}$

D. $\frac{(b^2 - 2ac)}{ac}$

Answer: C



Watch Video Solution

9. IF α, β are the roots of $ax^2 + bx + c = 0$ then $\alpha^3 + \beta^3 =$

A. $\frac{3abc - b^3}{a^3}$

B. $-\frac{b}{c}$

C. $\frac{(b^2 - 2ac)}{c^2}$

D. $\frac{(b^2 - 2ac)}{ac}$

Answer: A



Watch Video Solution

10. IF α, β are the roots of $ax^2 + bx + c = 0$ then $\frac{\alpha^2}{\beta} + \frac{\beta^2}{\alpha}$

A. $\frac{3abc - b^3}{a^3}$

B. $\frac{3abc - b^3}{a^2c}$

C. $\frac{b^2 - 2ac}{c^2}$

D. $\frac{b^2 - 2ac}{ac}$

Answer: B



Watch Video Solution

11. IF α, β are the roots of $ax^2 + bx + c = 0$ then $\frac{1}{\alpha^3} + \frac{1}{\beta^3} =$

A. $\frac{3abc - b^3}{a^3}$

B. $\frac{3abc - b^3}{a^2c}$

C. $\frac{3abc - b^3}{c^3}$

D. $\frac{b^2 - 2ac}{ac}$

Answer: C



Watch Video Solution

12. If α, β are the roots of $ax^2 + bx + c = 0$ then $\alpha^5\beta^8 + \alpha\beta^5 =$

A. $\frac{c^5(3abc - b^3)}{a^8}$

B. $\frac{c^3(3abc - b^3)}{a^3}$

C. $\frac{a^2(3abc - b^3)}{c^3}$

D. $\frac{a^3(3abc - b^3)}{c^3}$

Answer: A



Watch Video Solution

13. IF α, β are the roots of $ax^2 + bx + c = 0$ then $\frac{\alpha^2 + \beta^2}{\alpha^{-2} + \beta^{-2}} =$

A. $\frac{c^2}{a^2}$

B. $\frac{3abc - b^3}{a^2c}$

C. $\frac{3abc - b^3}{c^3}$

D. $\frac{b^2 - 2ac}{ac}$

Answer: A



Watch Video Solution

14. IF α, β are the roots of $ax^2 + bx + c = 0$ then $\frac{\alpha^3 + \beta^3}{\alpha^{-3} + \beta^{-3}} =$

A. $\frac{c^2}{d^2}$

B. $\frac{c^3}{a^3}$

C. $\frac{3abc - b^3}{c^3}$

D. $\frac{b^2 - 2ac}{ac}$

Answer: B



Watch Video Solution

15. IF α, β are the roots of $ax^2 + bx + c = 0$ then $\left(\frac{\alpha}{\beta} - \frac{\beta}{\alpha}\right)^2 =$

A. $\frac{b^2(b^2 - 4ac)}{c^2a^2}$

- B. $\frac{b^2(b^2 - 4ac)}{ca^3}$
- C. $\frac{b^2(b^2 - 4ac)}{a^4}$
- D. $\frac{b^2(b^2 - 4ac)}{c^4}$

Answer: A

 [Watch Video Solution](#)

16. IF α, β are the roots of $ax^2 + bx + c = 0$ then $\left(\frac{1}{\alpha^2} - \frac{1}{\beta^2}\right)^2 =$

- A. $\frac{b^2(b^2 - 4ac)}{c^2a^2}$
- B. $\frac{b^2(b^2 - 4ac)}{ca^3}$
- C. $\frac{b^2(b^2 - 4ac)}{a^4}$
- D. $\frac{b^2(b^2 - 4ac)}{c^4}$

Answer: D

 [Watch Video Solution](#)

17. IF α, β are the roots of $ax^2 + bx + c = 0$ then $(a\alpha + b)^{-2} + (a\beta + b)^{-2} =$

A. $\frac{b^2 - 2ac}{a^4}$

B. $\frac{b^2 - 2ac}{a^3c}$

C. $\frac{b^2 - 2ac}{a^2c^2}$

D. $\frac{b^2 - 2ac}{c^4}$

Answer: C



Watch Video Solution

18. IF α, β are the roots of $ax^2 + bx + c = 0$ then $(a\alpha + b)^{-3} + (a\beta + b)^{-3} =$

A. $a^3 - 2abc$

B. $b^3 - 3abc$

C. $\frac{c^3 - 3abc}{b^3c^3}$

D. $\frac{b^3 - 3abc}{a^3c^3}$

Answer: D



Watch Video Solution

19. IF α, β are the roots of $ax^2 + bx + c = 0$ then

$$\left(\frac{\alpha}{a\beta + b}\right)^3 - \left(\frac{\beta}{a\alpha + b}\right)^3 =$$

A. 0

B. 1

C. $(a + b)^2$

D. $(a - b)^2$

Answer: A



Watch Video Solution

20. IF α, β are the roots of the equation $8x^2 - 3x + 27 = 0$ then the value of $\left(\frac{\alpha^2}{\beta}\right)^{1/3} + \left(\frac{\beta^2}{\alpha}\right)^{1/3}$ is

A. $1/3$

B. $1/4$

C. $7/2$

D. 4

Answer: B



[Watch Video Solution](#)

21. If α, β are the roots of $ax^2 + bx + c = 0$ then $(1 + \alpha + \alpha^2)(1 + \beta + \beta^2)$ is

A. zero

B. positive

C. negative

D. not determined

Answer: B



[Watch Video Solution](#)

22. If α, β are the roots of $ax^2 - 2bx + c = 0$ then $\alpha^3\beta^3 + \alpha^2\beta^3 + \alpha^3\beta^2 =$

A. $\frac{c^2(2b + c)}{a^3}$

B. $\frac{bc^2}{a^3}$

C. $\frac{c^3}{a^3}$

D. $\frac{c^2(b + 2ac)}{a^3}$

Answer: A



[Watch Video Solution](#)

23. IF α, β are real and $\alpha^2, -\beta^2$ are the roots of $a^2x^2 + x + 1 - a^2 = 0 (A > 1)$ then $\beta^2 =$

A. a^2

B. 1

C. $1 - a^2$

D. $1 + a^2$

Answer: B



[Watch Video Solution](#)

24. If α, β are the roots of $x^2 - 2x + 4 = -0$ then $\alpha^5 + \beta^5 =$

A. 8

B. 16

C. 32

D. 64

Answer: C



Watch Video Solution

25. If α and β are the roots of the equation $x^2 - 2x + 4 = 0$, then $\alpha^9 + \beta^9 =$

A. -2^8

B. 2^9

C. -2^{10}

D. 2^{10}

Answer: C



Watch Video Solution

26. Let α and β be the roots of equation $px^2 + qx + r = 0$, $p \neq 0$, if p, q, r are in A.P and $\frac{1}{\alpha} + \frac{1}{\beta} = 4$ then the value of $|\alpha - \beta|$ is

A. $\frac{\sqrt{34}}{9}$

B. $\frac{2\sqrt{13}}{9}$

C. $\frac{\sqrt{61}}{9}$

D. $\frac{2\sqrt{17}}{9}$

Answer: B



Watch Video Solution

27. IF α, β be the roots of $6x^2 - 6x + 1 = 0$ then

$$\frac{1}{2}(a + b\alpha + c\alpha^2 + d\alpha^3) + \frac{1}{2}(a + b\beta + c\beta^2 + d\beta^3) =$$

A. $a + \beta + c + d$

B. $a + 2b + 3c + 4d$

C. $a + b/2 + c/3 + d/4$

D. none

Answer: C

 [View Text Solution](#)

28. If α, β are the roots of $x^2 - p(x + 1) + c = 0$ then $(1 + \alpha)(1 + \beta) =$

A. 1

B. c

C. $1 - c$

D. $1 + c$

Answer: D

 [Watch Video Solution](#)

29. If α, β are the roots of $x^2 - p(x + 1) - c = 0$ then

$$\frac{\alpha^2 + 2\alpha + 1}{\alpha^2 + 2\alpha + c} + \frac{\beta^2 + 2\beta + 1}{\beta^2 + 2\beta + c} =$$

A. 3

B. 2

C. 1

D. 0

Answer: C



[View Text Solution](#)

30. IF α, β are the roots of $ax^2 + bx + c = 0$ and $s_n = \alpha^n + \beta^n$ then

$$\begin{vmatrix} 3 & 1 + S_1 & 1 + S_2 \\ 1 + S_1 & 1 + S_2 & 1 + S_3 \\ 1 + s_2 & 1 + s_3 & 1 + S_4 \end{vmatrix} =$$

A. 0

B. $\frac{(b^2 - 4ac)(a + b + c)^2}{a^4}$

C. $\frac{b^2 - 4ac}{a^4}$

D. $\frac{(a + b + c)^2}{a^4}$

Answer: B



[View Text Solution](#)

31. Let α and β be the roots of equation $x^2 - 6x - 2 = 0$. If $a_n = \alpha^n - \beta^n$, for $n \geq 1$ then the value of $\frac{a_{10} - 2a_8}{2a_9}$ is equal to

A. 6

B. -6

C. 3

D. -3

Answer: C

[Watch Video Solution](#)

32. IF α, β are the roots of the equation $ax^2 + bx + c = 0$, then the

value of the determinant
$$\begin{vmatrix} 1 & \cos(\beta - \alpha) & \cos \alpha \\ \cos(\alpha - \beta) & 1 & \cos \beta \\ \cos \alpha & \cos \beta & 1 \end{vmatrix}$$

A. $\sin(\alpha + \beta)$

B. $\sin \alpha \sin \beta$

C. $1 + \cos(\alpha + \beta)$

D. 0

Answer: D



Watch Video Solution

33. IF the roots of the quadratic equation $x^2 + px + q = 0$ are $\tan 30^\circ$ and $\tan 15^\circ$, respectively then the value of $2 + q - p$ is

A. 0

B. 1

C. 2

D. 3

Answer: D



Watch Video Solution

34. IF $\tan A, \tan B$ are the roots of $x^2 - px + q = 0$, the value of $\sin^2(A + B)$ is

A. $\frac{p^2}{p^2 + (1 - q)^2}$

B. $\frac{p^2}{p^2 + q^2}$

C. $\frac{q^2}{p^2 + (1 - q)^2}$

D. $\frac{p^2}{(p + q)^2}$

Answer: A



Watch Video Solution

35. If α, β are the roots of $x^2 + ax - b = 0$ and γ, σ are the roots of $x^2 + ax + b = 0$ then $(\alpha - \gamma)(\beta - \gamma)(\alpha - \sigma)(\beta - \sigma) =$

A. b^2

B. $2b^2$

C. $3b^2$

D. $4b^2$

Answer: D



Watch Video Solution

36. If α, β are the roots of $x^2 + px - q = 0$ and γ, δ are the roots of $x^2 + px + r = 0$ then $(\alpha - \gamma)(\beta - \gamma)(\alpha - \delta)(\beta - \delta) =$

A. $2q^2$

B. $2q^2$

C. $(q + r)^2$

D. $(q - r)^2$

Answer: C



Watch Video Solution

37. If α, β are the roots of $x^2 + px + 1 = 0$ and γ, δ are the roots of $x^2 + qx + 1 = 0$ then $(\alpha - \gamma)(\beta - \gamma)(\alpha + \delta)(\beta + \delta) =$

A. $2p^2$

B. $2q^2$

C. $p^2 - q^2$

D. $q^2 - p^2$

Answer: D



Watch Video Solution

38. If α, β are the roots of $ax^2 + 2bx + c = 0$ and $\alpha + \sigma, \beta + \sigma$ are the roots of $Ax^2 + 2Bx + c = 0$ then $\frac{b^2 - ac}{B^2 - AC} =$

A. a/A

B. A/a

C. $(a/A)^2$

D. $(A/a)^2$

Answer: C



Watch Video Solution

39. α, β are the roots of $ax^2 + bx + c = 0$ and γ, σ are the roots of $px^3 + qx + r = 0$ and $D_1 : D_2$ be the respective discrimination of these equations .If $\alpha, \beta\gamma$ and δ are in A.P then $D_1 : D_2 =$

A. $a^2 : p^2$

B. $b^2 : q^2$

C. $c^2 : r^2$

D. none

Answer: A



Watch Video Solution

40. The ratio of the roots of the equation $ax^2 + bx + c = 0$ is same as the ratio of the roots of the equation $px^2 + qx + r = 0$. If D_1 and D_2 are the discrimination of $ax^2 + bx + c = 0$ and $px^2 + qx + r = 0$ respectively, then $D_1 : D_2 =$

A. $a^2 : p^2$

B. $b^2 : q^2$

C. $c^2 : r^2$

D. none

Answer: B

 [Watch Video Solution](#)

41. α, β are the roots of $ax^2 + bx + c = 0$ and γ, σ are the roots of $px^3 + qx + r = 0$ and $D_1 : D_2$ be the respective discrimination of these equations. If $\alpha, \beta\gamma$ and δ are in A.P then $D_1 : D_2 =$

A. $a^2 : p^2$

B. $a^2 : b^2$

C. $a^2 : c^2$

D. $a^2 : d^2$

Answer: A



Watch Video Solution

42. Let p and q be the roots of $x^2 - 2x + A = 0$ and let r and s be the roots of $x^2 - 18x + B = 0$. If $p < q < r < s$ are in ordered pair $(A, B) =$

A. $(-3, 77)$

B. $(77, -3)$

C. $(-3, -77)$

D. none of these

Answer: A



Watch Video Solution

43. Let x_1, x_2 , be the roots of the equation $x^2 - 3x + p = 0$ and let x_3, x_4 be the roots of the equation $x^2 - 12x + q = 0$ if the numbers x_1, x_2, x_4 (in order) form an increasing G.P then

A. $p = 2, q = 16$

B. $p = 2, q = 32$

C. $p = 4, q = 16$

D. $p = 4, q = 32$

Answer: B



Watch Video Solution

44. Let α, β be the roots of $x^2 - x + p = 0$ and γ, δ be the roots of $x^2 - 4x + q = 0$. If $\alpha, \beta, \delta, \gamma$ are in G.P then the integral values of p and q respectively , are

A. $-2, -32$

B. $-2, 3$

C. $-6, 3$

D. $-6, -32$

Answer: A



Watch Video Solution

45. x_1 and x_3 are the roots of the equation $Ax^2 - 4x + 1 = 0$ and x_2 and $x_4 = 0$ are the roots of the equation $Bx^2 - 6x + 1 = 0$ if x_1, x_2, x_3, x_4 form a H.P, then $(A,B) =$

A. $(3, 3)$

B. $(8, 8)$

C. $(3, 8)$

D. $(8, 3)$

Answer: C



Watch Video Solution

46. The coefficient of x in a quadratic equation $x^2 + px + q = 0$ was taken as 17 in place of 13 and its roots found to be -2 and -15 . The roots of the original equation are

A. 2, 15

B. 10, 3

C. -10 , -3

D. -2 , -15

Answer: C



Watch Video Solution

47. The students while solving a quadratic equation in x , one copied the constant term incorrectly and got the roots 3 and 2, The other copied the constant term and coefficient of x^2 as -6 and 1 respectively. The correct roots are

A. 3, - 2

B. - 3, 2

C. - 6, - 1

D. 6, - 1

Answer: D



[Watch Video Solution](#)

48. If 8 and 2 are the roots of $x^2 + ax + \beta = 0$ and 3, 3 are the roots of $x^2 + \alpha x + b = 0$ then the roots of the equation $x^2 + ax + b = 0$ are

A. 1, - 1

B. $-9, 2$

C. $-8, -2$

D. $9, 1$

Answer: D



Watch Video Solution

49. IF x_1 and x_2 are the real roots of the equation $x^2 - kx + c = 0$ then the distance between the points A $(x_1, 0)$ and $(x_2, 0)$ is

A. $\sqrt{k^2 - c}$

B. $\sqrt{c - k^2}$

C. $\sqrt{k^2 - 4c}$

D. $\sqrt{k^3 + 4c}$

Answer: C



Watch Video Solution

50. If one root of the equation $x^2 - 2(1 + i)x + (2 - i) = 0$ is $2 - i$ then the root is

A. $-i$

B. $2 + i$

C. i

D. $2 - i$

Answer: A



Watch Video Solution

51. If one root of $x^2 + ax + 8 = 0$ is 4 and the equation $x^2 + ax + b = 0$ has equal roots then the value of $b =$

A. 7

B. 9

C. 1

D. 3

Answer: B



[Watch Video Solution](#)

52. IF the roots of the equation $x^2 - 5x + 16 = 0$ are α, β and the roots of the equation $x^2 + px + q = 0$ and $\alpha^2 + \beta^2, \frac{\alpha\beta}{2}$, then

A. $p = 1, q = -56$

B. $p = -1, q = -56$

C. $p = 1, q = 56$

D. $p = -1, q = 56$

Answer: B



[Watch Video Solution](#)

53. IF $x^2 - 3x + 2$ is a factor of $x^4 - px^2 + q = 0$ then $(p, q) = 0$

A. $(-4, -5)$

B. $(4, 5)$

C. $(-5, -4)$

D. $(5, 4)$

Answer: D



[Watch Video Solution](#)

54. IF the product of the roots of $5x^2 - 4x + 2 + m(4x^2 - 2x - 1) = 0$

is 3, then $m =$

A. 0

B. -1

C. 2

D. 3

Answer: B



[Watch Video Solution](#)

55. IF $k > 0$ and the product of the roots of the equation $x^2 - 3kx + 2e^2 \log k - 1 = 0$ is 7 then the sum of the roots is

A. 2

B. 4

C. 6

D. 8

Answer: C



[Watch Video Solution](#)

56. The value of K , so that the sum and product of the roots of $2x^2 + (k - 3)x + 3k - 5 = 0$ are equal is

A. 0

B. 10

C. 2

D. 9

Answer: C



Watch Video Solution

57. IF the sum of the square of the roots of $x^2 + px - 3 = 0$ is 10 then the values of p=

A. $+2$

B. ± 3

C. 3

D. -3

Answer: A



Watch Video Solution

58. α, β are the roots of $x^2 + kx + 2 = 0$. If $\alpha - \beta = 1$ then $K =$

A. ± 1

B. ± 2

C. ± 3

D. ± 4

Answer: C



Watch Video Solution

59. α, β are roots of the equation $\lambda(x^2 - x) + x + 5 = 0$ If λ_2 are the two values of λ for which the roots α, β are connected by the relation

$\frac{\alpha}{\beta} + \frac{\beta}{\alpha} = 4$, then the value of $\frac{\lambda_1}{\lambda_2} + \frac{\lambda_2}{\lambda_1}$ is

A. 150

B. 254

C. 180

D. 1022

Answer: D



[Watch Video Solution](#)

60. The harmonic mean of the roots of the equation

$$(5 + \sqrt{2})x^2 - (4 + \sqrt{5})x + (8 + 2\sqrt{5}) = 0 \text{ is}$$

A. 2

B. 4

C. 6

D. 8

Answer: B



[Watch Video Solution](#)

61. IF C and D are the roots of $(x - a)(x - b) - k = 0$ then the roots of $(x - c)(x - d) + k = 0$ are

A. b,c

B. a,b

C. a,c

D. a,d

Answer: B



[Watch Video Solution](#)

62. IF α, β be the roots of the equation $(x - a)(x - b) + c = 0 (c \neq 0)$ then the roots of the equation $(x - c - \alpha)(x - c - \beta) = c$ are

A. a and b+c

B. a+c and b

C. $a+c$ and $b+c$

D. $a-c$ and $b-c$

Answer: C



[View Text Solution](#)

63. The equation whose roots are $2\sqrt{3}-5$ and $-2\sqrt{3}-5$ is

A. $x^2 + 10x - 13 = 0$

B. $x^2 - 10x + 13 = 0$

C. $x^2 + 10x + 13 = 0$

D. $x^2 - 10 - 13 = 0$

Answer: C



[Watch Video Solution](#)

64. The equation whose roots are $3 + 2i$, $3 - 2i$ is

A. $x^2 - 8x + 15 = 0$

B. $15x^2 - 34x + 15 = 0$

C. $x^2 - 6x + 13 = 0$

D. $x^2 + 1 = 0$

Answer: C



Watch Video Solution

65. If α and β are the roots of $2x^2 + x + 3 = 0$, then the equation

whose roots are $\frac{1 - \alpha}{1 + \alpha}$ and $\frac{1 - \beta}{1 + \beta}$ is

A. $2x^2 + x + 3 = 0$

B. $2x^2 - x - 3 = 0$

C. $2x^2 + x - 3 = 0$

$$D. 2x^2 - x - 3 = 0$$

Answer: A



Watch Video Solution

66. IF α, β are the roots of $x^2 + 5x - 4 = 0$ then the equation whose roots are $\frac{\alpha + 2}{3}, \frac{\beta + 2}{3}$ is

A. $9x^2 + 3x + 10 = 0$

B. $9x^2 + 3x - 10 = 0$

C. $9x^2 + x = 0$

D. $2x^2 - 3x + 10 = 0$

Answer: B



Watch Video Solution

67. If α, β are the roots of $x^2 - x + 1 = 0$ then the quadratic equation whose roots are $\alpha^{2015}, \beta^{2015}$ is

A. $x^2 - x + 1 = 0$

B. $x^2 + x + 1 = 0$

C. $x^2 + x - 1 = 0$

D. $x^2 - x - 1 = 0$

Answer: A



[Watch Video Solution](#)

68. IF α, β are the roots of $x^2 - ax + b = 0$, then the whose roots are $\frac{\alpha + \beta}{\alpha}, \frac{\alpha + \beta}{\beta}$ is

A. $bx^2 + a^2x + a^2 = 0$

B. $bx^2 - a^2x + a^2 = 0$

C. $b^2x^2 - a^2x + a^2 = 0$

D. $ax^2 + b^2x + b^2 = 0$

Answer: B



Watch Video Solution

69. IF α, β are the roots of $ax^2 + bx + c = 0$ then the equation roots are $\alpha/\beta, \beta/\alpha$ is

A. $acx^2 - (b^2 - 2ac)x + ac = 0$

B. $a^3x^2 + (b^3 - 3abc)x + c^3 = 0$

C. $x^2 - 2qx + (q^2 - p^2) = 0$

D. $x^2 + 2qx + (q^2 + p^2) = 0$

Answer: A



Watch Video Solution

70. IF α, β are the roots of the equation $ax^2 + bx + c = 0$ then the quadratic equation whose roots are $\alpha + \beta, \alpha\beta$ is

A. $a^2x^2 + a(b - c)x + bc = 0$

B. $a^2x^2 + a(b - c)x - bc = 0$

C. $ax^2 + (b + c)x + bc = 0$

D. $ax^2 - (b + c)x - bc = 0$

Answer: B



[Watch Video Solution](#)

71. In a $\triangle ABC$, the value of $\angle A$ is obtained from the equation $3 \cos A + 2 = 0$. The quadratic equation, whose roots are $\sin A$ and $\tan A$ is

A. $3x^2 + \sqrt{5}x - 5 = 0$

B. $6x^2 - \sqrt{5}x - 5 = 0$

C. $6x^2 + \sqrt{5}x - 5 = 0$

D. $6x^2 + \sqrt{5}x + 5 = 0$

Answer: C

 [Watch Video Solution](#)

72. The equation whose roots are the cubes of the roots of the equation

$ax^2 + bx + c = 0$ is

A. $a^3x^2 + (b^2 - 3abc)x + c^3 = 0$

B. $a^3x^2 + (b^2 + 3abc)x + c^2 = 0$

C. $a^3x^2 - (b^3 + 3abc)x + c^2 = 0$

D. none

Answer: A

 [Watch Video Solution](#)

73. IF α, β are the roots of $ax^2 + bx + c = 0$ and γ, δ are the roots of $lx^2 + mx + n = 0$ then the equation whose roots are $\alpha\gamma + \beta\delta, \alpha\delta + \beta\gamma$ is

A. $a^2t^2x^2 - ablmx + b = 0$

B. $a^2t^2x^2 - ablmx + (b^2nl + m^2ac - 4acnl) = 0$

C. $a^3t^2x^2 - ablmx$

D. none

Answer: B



Watch Video Solution

74.

IF

$$a = \cos\left(\frac{2\pi}{7}\right) + i \sin\left(\frac{2\pi}{7}\right), \alpha = a + a^2 + a^4 \text{ and } \beta = a^3 + a^5 + a^6$$

then α, β are the roots of the equation

A. $x^2 + x + 1 = 0$

B. $x^2 + x + 2 = 0$

C. $x^2 + x + 2 = 0$

D. $x^2 + 2x + 3 = 0$

Answer: B



Watch Video Solution

75. Let $\alpha \neq \beta$ satisfy $\alpha^2 + 1 = 6\alpha$, $\beta^2 + 1 = 6\beta$. Then, the quadratic equation whose roots are $\frac{\alpha}{\alpha + 1}$, $\frac{\beta}{\beta + 1}$ is

A. $8x^2 + 8x + 1 = 0$

B. $8x^2 - 8x - 1 = 0$

C. $8x^2 - 8x + 1 = 0$

D. $8x^2 + 8x - 1 = 0$

Answer: C



Watch Video Solution

76. IF $3p^2 = 5p + 2$ and $3q^2 = 5q + 2$ where $p \neq q$ then the equation whose roots are $3p - 2q$ and $3q - 2p$ is

A. $3x^2 - 5x - 100 = 0$

B. $5x^2 + 3x + 100 = 0$

C. $3x^2 - 5x + 100 = 0$

D. $3x^2 + 5x - 100 = 0$

Answer: A



Watch Video Solution

77. IF α, β are the roots of the equation $x^2 + 2ax + b = 0$, then the quadratic equation with rational coefficient one of whose roots is

$\alpha + \beta + \sqrt{\alpha^2 + \beta^2}$ is

A. $x^2 - 4ax + 12 = 0$

B. $x^2 + 4ax - 2b = 0$

C. $x^2 - 4ax - 2b = 0$

D. $x^2 + 4ax + 2b = 0$

Answer: D



Watch Video Solution

78. The quadratic equation for which the sum of the roots is 7 and the sum of the squares of the roots is 25 is

A. $x^2 - 7x + 12 = 0$

B. $x^2 \pm 54x + 6 = 0$

C. $x^2 - 12x + 35 = 0$

D. $5x^2 + 2x + 11 = 0$

Answer: A



Watch Video Solution

79. The quadratic equation for which the sum of the roots is 12 and the sum of the cubes of the roots is 468 is

A. $x^2 - 7x + 12 = 0$

B. $x^2 \pm 54x + 6 = 0$

C. $x^2 - 12x + 35 = 0$

D. $5x^2 + 2x + 11 = 0$

Answer: C



Watch Video Solution

80. IF $\alpha + \beta = -2$ and $\alpha^3 + \beta^3 = -56$

, then the quadratic equation whose roots are α and β is

A. $x^2 + 2x - 16 = 0$

B. $x^2 + 2x - 15 = 0$

C. $x^2 + 2x - 12 = 0$

D. $x^2 + 2x - 8 = 0$

Answer: D



[Watch Video Solution](#)

81. Two complex numbers α and β are such that $\alpha + \beta = 2$ and $\alpha^4 + \beta^4 = 272$, then the quadratic equation whose roots are α and β is

A. $x^2 - 2x - 16 = 0$

B. $x^2 + 2x - 15 = 0$

C. $x^2 - 2x - 8 = 0$

D. none of these

Answer: C



[Watch Video Solution](#)

82. IF the arithmetic mean of the roots of a quadratic equation is $\frac{8}{5}$ and the arithmetic mean of their reciprocals is $\frac{8}{7}$ then the equation is

A. $5x^2 + 16x + 7 = 0$

B. $5x^2 - 16x + 7 = 0$

C. $7x^2 + 16x + 5 = 0$

D. $7x^2 - 16x + 5 = 0$

Answer: B



[Watch Video Solution](#)

83. Let two numbers have arithmetic mean 9 geometric mean 4. then these numbers are the roots of the quadratic equation

A. $x^2 + 18x + 16 = 0$

B. $x^2 - 18x - 16 = 0$

C. $x^2 + 18 - 16 = 0$

D. $x^2 - 18 + 16 = 0$

Answer: D



Watch Video Solution

84. The equation whose roots are the arithmetic mean and twice the H.M between the roots of the equation $x^2 + ax - b = 0$ is

A. $2ax^2 + (a^2 - 8b)x + 4ab = 0$

B. $2ax^2 + (a^2 - 8b)x - 4ab = 0$

C. $2ax^2 + (a^2 + 8b)x - 4ab = 0$

D. none

Answer: B



Watch Video Solution

85. IF α, β are the roots of $9x^2 + 6x + 1 = 0$ then the equation with the roots $1/\alpha, 1/\beta$ is

A. $2x^2 + 3x + 18 = 0$

B. $x^2 + 6x - 9 = 0$

C. $x^2 + 6x + 9 = 0$

D. $x^2 - 6x + 9 = 0$

Answer: C



Watch Video Solution

86. The equation whose roots are greater by 1 than those of $2x^2 - 3x + 1 = 0$ is

A. $3x^2 - 5x - 2 = 0$

B. $2x^2 - 7x + 6 = 0$

C. $2x^2 + 5x + 7 = 0$

D. $3x^2 + \% x - 7 = 0$

Answer: B



Watch Video Solution

87. IF α, β are the roots of $ax^2 + bx + c = 0$ then the equation whose roots are $2 + \alpha, 2 + \beta$ is

A. $ax^2 + x(4a - b) + 4a - 2b + C = 0$

B. $ax^2 + x(4a - b) + 4a + 2b + c = 0$

C. $ax^2 + x(b - 4a) + 4a + 2bc = 0$

D. $ax^2 + x(b - 4a) + 4a - 2b + c = 0$

Answer: D



Watch Video Solution

88. The equation whose roots are smaller by 1 than those of $2x^2 - 5x + 6 = 0$ is

A. $2x^2 - 9x + 13 = 0$

B. $2x^2 - x + 3 = 0$

C. $2x^2 + 9x + 13 = 0$

D. $2x^2 + x + 3 = 0$

Answer: B



[Watch Video Solution](#)

89. the equation formed by decreasing each root of $ax^2 + bx + C = 0$ by 1 is $2x^2 + 8x + 2 = 0$ then

A. $a = -b$

B. $b = -c$

C. $c = -a$

D. $b = a + c$

Answer: B



Watch Video Solution

90. If α, β are the roots of $a^2x + bx + c = 0$ and $\alpha + h + \beta + h$ are the roots of $px^2 + qx + r = 0$ then $h =$

A. $\left(\frac{b}{a} - \frac{q}{p}\right)$

B. $\frac{1}{2}\left(\frac{b}{a} - \frac{q}{p}\right)$

C. $-\frac{1}{2}\left(\frac{a}{b} - \frac{p}{q}\right)$

D. none

Answer: B



Watch Video Solution

91. IF α, β are the roots of $x^2 + bx + c = 0$ and $\alpha + h, \beta + h$ are the roots of $x^2 + qx + r = 0$ then $h =$

A. $b + q$

B. $b - q$

C. $\frac{1}{2}(b + q)$

D. $\frac{1}{2}(b - q)$

Answer: D



Watch Video Solution

92. IF α and β are the roots of the equation $ax^2 + bx + C = 0$ and if $px^2 + qx + r = 0$ has roots $\frac{1 - \alpha}{\alpha}$ and $\frac{1 - \beta}{\beta}$ then $r =$

A. $a + 2b$

B. $a + b + C$

C. $ab + bc + ca$

D. abc

Answer: B



Watch Video Solution

93. The equation whose roots are numerically equal but opposite in sign of the roots of $3x^2 - 5x - 7 = 0$ is

A. $3x^2 - 5x - 2 = 0$

B. $2x^2 - 7x + 6 = 0$

C. $2x^2 + 5x + 7 = 0$

D. $3x^2 + 5x - 7 = 0$

Answer: D



Watch Video Solution

94. The equation whose roots are multiplied by 3 of those of $2x^2 + 3x - 1 = 0$ is

A. $2x^2 + 9x - 9 = 0$

B. $2x^2 - 7x + 6 = 0$

C. $2x^2 + 5x + 7 = 0$

D. $3x^2 + 5x - 7 = 0$

Answer: A



[Watch Video Solution](#)

95. The condition that one root of $ax^2 + bx + c = 0$ may be n times the other root is

A. $nb^2 = ac(n + 1)$

B. $nb^2 = ac(n + 1)^2$

C. $nb = ac(n + 1)$

$$D. nb = ac(n + 1)^2$$

Answer: B



Watch Video Solution

96. The condition that one root of $ax^2 + bx + c = 0$ may be the double the other is

A. $b^2 = 2ac$

B. $b^2 = 2ac$

C. $2b^2 = 9ac$

D. $2b^2 = 3ac$

Answer: C



Watch Video Solution

97. If one root of $x^2 + kx + 12 = 0$ may be the triple the other , then k=

A. ± 8

B. 3

C. $\pm 5\sqrt{10}$

D. $2 \pm \sqrt{5}$

Answer: A



[Watch Video Solution](#)

98. IF one root of $px^2 - 14x + 8 = 0$ is six times the other then $p =$

A. 1

B. 2

C. 3

D. 4

Answer: C



Watch Video Solution

99. IF the roots of $ax^2 + cx + c = 0$ are in the ratio $p:q$ then

$$\sqrt{\left[\frac{p}{q}\right]} + \sqrt{\left[\frac{q}{p}\right]} =$$

A. $\sqrt{a/c}$

B. $\sqrt{c/a}$

C. $-\sqrt{a/c}$

D. $-\sqrt{c/a}$

Answer: D



Watch Video Solution

100. IF one root of $x^2 - x - k = 0$ is square that of the other, then

k=.....

A. $2 \pm \sqrt{3}$

B. $3 \pm \sqrt{2}$

C. $2 \pm \sqrt{5}$

D. $5 \pm \sqrt{2}$

Answer: C

 [Watch Video Solution](#)

101. If one root of $x^2 + px + 1 = 0$ is square that of the other then $p =$

A. 1, - 2

B. 3, - 1

C. 2, $-\sqrt{5}$

D. $4 + \sqrt{5}$

Answer: A

 [Watch Video Solution](#)

102. IF the harmonic mean between the roots of $(5 + \sqrt{2})x^2 - bx + (8 + 2\sqrt{5}) = 0$ is 4 then value of b is

A. 2

B. 3

C. $4 - \sqrt{5}$

D. $4 + \sqrt{5}$

Answer: D



[Watch Video Solution](#)

103. IF the harmonic mean of the roots of $\sqrt{2}x^2 - bx + (8 - 2\sqrt{5}) = 0$ is 4 then the value of b=

A. 2

B. 3

C. $4 - \sqrt{5}$

D. $4 + \sqrt{5}$

Answer: C

 [Watch Video Solution](#)

104. The condition that a root of $ax^2 + bx + c = 0$ may be the reciprocal of a root of $a_1x^2 + b_1x + c_1 = 0$ is

A. $(aa_1 - cc_1)^2(ab_1 + bc_1)(a_1b + b_1c)$

B. $(aa_1 - cc_1)^2 = (ab_1 - bc_1)(a_1b - b_1c)$

C. $(aa_1 - bb_1)^2 = (ac_1 - bc_1)$

D. none

Answer: B

 [Watch Video Solution](#)

105. IF the sum of the roots of the quadratic equation $ax^2 + bx + c = 0$ is equal to sum of the square of their reciprocals then $\frac{b^2}{ac} + \frac{bc}{a^2} =$

- A. 2
- B. -2
- C. 1
- D. -1

Answer: A



[Watch Video Solution](#)

106. IF the sum of the roots of the equation $x^2 + px + q = 0$ is 3 times their difference , then

- A. $2p^2 = q$
- B. $2p^2 = 5q$
- C. $p^2 = 3q$

D. $2p^2 = 9q$

Answer: D



Watch Video Solution

107. If the roots of the equation $ax^2 + bx + c = 0$ be the square roots of the equation $lx^2 + mx + n = 0$ then

A. $2alc = lb^2 + ma^2$

B. $2alc = lb^2 - ma^2$

C. $alc = lb^2 + ma^2$

D. $alc = lb^2 - ma^2$

Answer: A



Watch Video Solution

108. IF the ratio of the roots of $x^2 + bx + c = 0$ and $x^2 + qx + r = 0$ are the same , then

A. $r^2c = qb^2$

B. $r^2b = qc^2$

C. $rb^2 = cq^2$

D. $rc^2 = bq^2$

Answer: C



Watch Video Solution

109. If the ratio of the roots of $a(x)^2 + bx + c = 0$ is same as that of the roots of $px^2 + qx + r = 0$ then $a/p, b/q, c/r$ are in

A. A. P

B. G. P

C. H. P

D. none

Answer: B



[Watch Video Solution](#)

110. The value of m for which one of the roots of $x^2 - 3x + 2m = 0$ is double of one of the roots of $x^2 - x + m = 0$ is

A. 1

B. -2

C. 2

D. none

Answer: B



[Watch Video Solution](#)

111. IF the difference of the roots of $x^2 - bx + c = 0$ is equal to the difference of the roots of $x^2 - cx + b = 0$ and $b \neq 0$, then $b+c=$

- A. -1
- B. -2
- C. -3
- D. -4

Answer: D



[Watch Video Solution](#)

112. The condition that the roots of the equation $ax^2 + bx + c = 0$ may differ by 5 is

- A. $b^2 - 25a^2 = 4ac$
- B. $b^2 - 5a^2 = 4ac$
- C. $b^2 + 15a^2 = 4ac$

D. none

Answer: A



[Watch Video Solution](#)

113. The condition that $\sin \theta \cos \theta$ may be the roots of $ax^2 + bx + c = 0$ is

A. $a(a + 2b) = c^2$

B. $a(a + 2c) = b^2$

C. $b(b + 2c) = a^2$

D. $b(b + 2a) = c^2$

Answer: B



[Watch Video Solution](#)

114. If $\sin \alpha$ and $\cos \alpha$ are the roots of $25x^2 + 5x - 12 = 0$, then value of $\sin 2\alpha$ is

- A. $12/25$
- B. $-12/25$
- C. $-24/25$
- D. $4/5$

Answer: C



[Watch Video Solution](#)

115. If $\tan A$ and $\tan B$ are the roots of the quadratic $x^2 - px + q = 0$ then $\sin^2(A + B) =$

- A. $\frac{p^2}{p^2 + q^2}$
- B. $\frac{p^2}{(p + q)^2}$
- C. $1 - \frac{p}{(1 - q)^2}$

D. $\frac{p^2}{p^2 + (1 - q)^2}$

Answer: D



Watch Video Solution

116. In a triangle PQR angle $R = \pi/2$ if $\tan(p/2)$ and $\tan(Q/2)$ are the roots of the equation $ax^2 + bx + c = 0$ ($a \neq 0$) then

A. $a+b=c$

B. $b + c = 0$

C. $a+c=b$

D. $b=c$

Answer: A



Watch Video Solution

117. In a triangle PQR $\angle R = \frac{\pi}{4}$, if $\tan\left(\frac{p}{3}\right)$ and $\tan\left(\frac{q}{3}\right)$ are the roots of the equation $ax^2 + bx + c = 0$ then

A. $a+b=c$

B. $b+c=0$

C. $a+c=b$

D. $b=c$

Answer: A



[Watch Video Solution](#)

118. If one root of $x^2 + px + 1 = 0$ is the cube of the other root, then $p =$

A. 0

B. 1

C. $1, \pm 2$

D. $0, \pm 2$

Answer: D



[Watch Video Solution](#)

119. IF one root of $ax^2 + bx + c = 0$ is equal to n th power of the other then $(a^n c)^{\frac{1}{n+1}} + (ac^n)^{\frac{1}{n+1}}$

A. 0

B. 1

C. b

D. $-b$

Answer: D



[Watch Video Solution](#)

120. IF x, a, b, c are real and $(x - a + b^2) + (x - b + c)^2 = 0$ then a, b, c are in

A. H.P

B. G.P

C. A.P

D. none of these

Answer: C



[Watch Video Solution](#)

121. IF the roots of $\frac{1}{x+a} + \frac{1}{x+b} = \frac{1}{c}$ are equal in magnitude and opposite in sign then the product of the roots is

A. $-\frac{1}{2}(a^2 + b^2)$

B. $\frac{1}{2}(a^2 + b^2)$

C. $-\frac{3}{2}(a^2 + b^2)$

D. none

Answer: A



Watch Video Solution

122. IF the roots of the equation $\frac{1}{x+p} + \frac{1}{x+q} = \frac{1}{r}$ are equal in magnitude and opposite in sign , then $p+q+r=$

A. r

B. $3r$

C. r°

D. $2r^2$

Answer: B



Watch Video Solution

123. IF $\frac{1}{x} + \frac{1}{x+a} = \frac{1}{m} + \frac{1}{m+a}$ has roots equal in magnitude but opposite sign, then

A. $a^2 = m^2$

B. $a^2 = 2m^2$

C. $2a^2 = m^2$

D. none

Answer: B



[Watch Video Solution](#)

124. IF the roots of $a^2x^2 - abx + c = 0$ are consecutive $\int e \geq rs$, then $b^2 - a^2 =$

A. $14c$

B. $7c$

C. $5c$

D. $4c$

Answer: D



Watch Video Solution

125. If r is the ratio of the roots of $ax^2 + bx + C = 0$ then $\frac{(r + 1)^2}{r} =$

A. 1

B. $b^2 - ac$

C. b^2 / ac

D. $b^2 - 4ac$

Answer: C



Watch Video Solution

126. IF the roots of $ax^2 + bx + c = 0$ are of the form $\frac{m+1}{m}, \frac{m+2}{m+1}$ then $(a+b+c)^2 =$

A. 0

B. 1

C. $b^2 - 4ac$

D. $2abc$

Answer: C



[Watch Video Solution](#)

127. IF $(a\alpha + b)^{-2} + (a\beta + b)^{-2} = 1$, where α, β are the roots of $ax^2 + bx + c = 0$ then $ac(ac + 2) =$

A. $b^2/2$

B. b^3

C. $2b$

D. b^2

Answer: D



Watch Video Solution

128. IF the equation $x^2 + 2(k + 1)x + 9k - 5 = 0$ has only negative roots , then

A. $k \leq 0$

B. $k \geq 0$

C. $k \geq 6$

D. $k \leq 6$

Answer: C



Watch Video Solution

129. All the values of m for which both roots of the equation $x^2 - 2mx + m^2 - 1 = 0$ are greater than -2 but less than 4 , lie in the interval

A. $-1 < m < 3$

B. $1 < m < 4$

C. $-2 < m < 0$

D. $m > 3$



Watch Video Solution

130. The values of a for which of a for which $2x^2 - 2(2a + 1)x + a(a + 1) = 0$ may have one root less than a and other root greater than a are given by

A. $1 > a > 0$

B. $-1 < a < 0$

C. $a > 0$

D. $a > 0$ or $a < -1$

Answer: D



[View Text Solution](#)

131. The value of a for which the equation $(1 - a^2)x^2 + 2ax - 1$ has roots belonging to $(0,1)$ is

A. $a > \frac{1 + \sqrt{5}}{2}$

B. $a > 2$

C. $\frac{1 + \sqrt{5}}{2} < a < 2$

D. $a > \sqrt{2}$

Answer: B



[Watch Video Solution](#)

132. The value of a for which each one of the roots of $x^2 - 4ax + 2a^2 - 3a + 5 = 0$ is greater than 2, are

A. $a \in (1, \infty)$

B. $a = 1$

C. $a \in (-\infty, 1)$

D. $a \in (9/2, \infty)$

Answer: D



Watch Video Solution

133. The value of ' a ' for which one root of the quadratic equation $(a^2 - 5a + 3)x^2 + (3a - 1)x + 2 = 0$ is twice as large as the other is

A. $-2/3$

B. $1/3$

C. $-1/3$

Answer: D[Watch Video Solution](#)

134. IF both the roots of the quadratic equation $x^2 - 2kx + k^2 + k - 5 = 0$ are less than 5, then k lies in the interval

A. $(5, 6]$

B. $(6, \infty)$

C. $(-\infty, 4)$

D. $[4, 5]$

Answer: C[Watch Video Solution](#)

135. If roots of the equation $x^2 - bx + c = 0$ be two consecutive integers, then $b^2 - 4c$ equals

A. -2

B. 3

C. 2

D. 1

Answer: D



[Watch Video Solution](#)

136. p and q are distinct prime numbers and if the equation $x^2 - px + q = 0$ has positive integer as its roots then the roots the roots of the equation are

A. $2, 3$

B. $1, 2$

C. 3, 1

D. 1, - 1

Answer: B

 [Watch Video Solution](#)

137. IF both the roots of the equation $x^2 - 6ax + 2 - 2a + 9a^2 = 0$ exceed 3, then

A. $a > \frac{9}{11}$

B. $a > \frac{11}{9}$

C. $a > \frac{11}{9}$

D. $a < \frac{11}{9}$

Answer: C

 [Watch Video Solution](#)

138. If the roots of $x^2 + x + a = 0$ exceed , a then

A. $2 < a < 3$

B. $a > 3$

C. $-3 < a < 3$

D. $a < \frac{1}{-2}$

Answer: D



Watch Video Solution

139. The roots of $3x^2 + 4x - 7 = 0$ are

A. rational and equal

B. rational and not equal

C. irrational

D. imaginary

Answer: B



[Watch Video Solution](#)

140. The roots of $5x^2 - 3x + 2 = 0$ are

- A. rational and equal
- B. rational and not equal
- C. irrational
- D. imaginary

Answer: D



[Watch Video Solution](#)

141. The roots of the equation $(b - c)x^2 + (c - a)x + (a - b) = 0$ are

- A. real

B. real and equal

C. real and not equal

D. imaginary

Answer: A



[Watch Video Solution](#)

142. IF $ad \neq bc$ then the roots of

$$(a^2 + b^2)x^2 + 2x(ac + bd) + (c^2 + d^2) = 0 \text{ are}$$

A. real

B. real and equal

C. real and not equal

D. imaginary

Answer: D



[Watch Video Solution](#)

143. For $p, q \in R$, the roots of $(p^2 + q^2)x^2 + 2(p + q)x + 2 = 0$ are

- A. real and equal
- B. real and unequal
- C. equal complex numbers
- D. unequal complex numbers

Answer: D



[Watch Video Solution](#)

144. The roots of the equation $2(a^2 + b^2)x^2 + 2(a + b)x + 1 = 0$ are

- A. real
- B. real and equal
- C. real and not equal
- D. imaginary

Answer: D



[Watch Video Solution](#)

145. The roots of

$$(x - b)(x - c) + (x - a)(x - c) + (x - a)(x - b) = 0 \text{ are}$$

- A. real
- B. equal
- C. real and not equal
- D. imaginary

Answer: A



[Watch Video Solution](#)

146. The roots of

$$(x - a)(x - a - 1) + (x - a - 1) + (x - a - 2) + (x - a)(x - a - 2) =$$

$a \in \mathbb{R}$ are always

- A. equal
- B. imaginary
- C. real and distinct
- D. rational and equal

Answer: C



[View Text Solution](#)

147. IF l, m, n are rational the roots of $(m + n)x^2 - (l + m + n)x + l = 0$ are

- A. rational
- B. rational and equal
- C. rational and not equal
- D. irrational

Answer: A



[Watch Video Solution](#)

148. If the roots of $a^2x^2 + 2bx + c^2 = 0$ are imaginary then the roots of $b(x^2 + 1) + 2acx = 0$ are

- A. real and equal
- B. real and unequal
- C. equal complex numbers
- D. unequal complex numbers

Answer: B



[Watch Video Solution](#)

149. If the roots of $(a^2 + b^2)x^2 + 2(bc + ad)x + (c^2 + d^2) = 0$ are real and equal then

A. $a/c = b/d$

B. $ac + bd = 0$

C. $ad+bc=0$

D. $ac - bd = 0$

Answer: D



Watch Video Solution

150. IF m, n, k are rational and $m = k + \frac{n}{k}$ then the roots of $x^2 + mx + n = 0$ are

A. $k, n/k$

B. $k, n/k$

C. $-k, -n/k$

D. $-k, n/k$

Answer: C

 [Watch Video Solution](#)

151. IF α, β are the roots of $x^2 + px + q = 0$ and α^4, β^4 are the roots of $x^2 - rx + s = 0$ then the equation $x^2 - 4qx + 2q^2 - r = 0$ has always

- A. two real roots
- B. two negative roots
- C. two positive roots
- D. one positive root and one negative root

Answer: A

 [Watch Video Solution](#)

152. Let a, b, c be real numbers $a \neq 0$ if α is a roots of $a^2x^2 + bx + c = 0$, β is a roots of $a^2x^2 - bx - c = 0$ and $0 < \alpha < \beta$, then the equation $a^2x^2 + 2bx + 2c = 0$ has a root γ that always satisfies

A. $\gamma = (\alpha + \beta) / 2$

B. $\gamma = \left(\alpha + \frac{\beta}{2} \right)$

C. $\gamma = \alpha$

D. $\alpha < \gamma < \beta$

Answer: D

 [View Text Solution](#)

153. If $p(x) = ax^2 + bx + c$ and $Q(x) = -ax^2 + dx + c$

where $ac \neq 0$ then the equation $P(x) \cdot Q(x) = 0$ has atleast

A. two real roots

B. two negative roots

C. two positive roots

D. one positive root and oine negative

Answer: A

 [Watch Video Solution](#)

154. Let $f(x) = x^2 + ax + b$, where $a, b \in \mathbb{R}$. If $f(x) = 0$ has all its roots imaginary, then the roots of $f(x) + f'(x) + f(x) = 0$ are :

- A. real and distinct
- B. imaginary
- C. equal
- D. rational and equal

Answer: B

 [Watch Video Solution](#)

155. $\alpha, \beta, \gamma \in \mathbb{R}$ is a root of $a^2x^2 + bx + c = 0$ β is a root of $a^2x^2 - bx - c = 0$ and γ is a root of $a^2x^2 + 2bx + 2c = 0$ then

A.
$$\begin{vmatrix} -\beta^2 & \beta & 1 \\ \alpha^2 & \alpha & 1 \\ \gamma^2 & 2\gamma & 2 \end{vmatrix} = 0$$

$$\text{B. } \begin{vmatrix} -\beta^2 & \beta & 1 \\ \alpha^2 & \alpha & 1 \\ \gamma^2 & \gamma & 2 \end{vmatrix} = 0$$

$$\text{C. } \begin{vmatrix} -\beta^2 & \beta & 1 \\ \alpha^2 & \alpha & 1 \\ \gamma^2 & 2\gamma & 2 \end{vmatrix} = 0$$

D. none

Answer: A



[Watch Video Solution](#)

156. IF the roots of the equation $x^2 + a^2 = 8x + 6a$ are real , then a lies between

A. 1 and 2

B. -1 and 8

C. 2 and 8

D. -2 and 8

Answer: D

[Watch Video Solution](#)

157. If the equation $(\cos p - 1)x^2 + \cos px + \sin p = 0$ in the variable x has real roots then p can taken any value in the interval

- A. $(0, 2\pi)$
- B. $(-\pi, 0)$
- C. $(-\pi/2, \pi/2)$
- D. $(0, \pi)$

Answer: D

[Watch Video Solution](#)

158. If $m, \in Z$ and the equation $mx^2 + (2m - 1)x + (m - 2) = 0$ has rational roots then m is of the form

- A. $n(n + 2), n \in z$

B. $n(n + 1), n \in \mathbb{Z}$

C. $n(n - 2), n \in \mathbb{Z}$

D. none

Answer: B



[Watch Video Solution](#)

159. If $a \in \mathbb{Z}$ and the equation $(x - a)(x - 10) + 1 = 0$ has integral roots then the values of a are

A. 10, 8

B. 12, 10

C. 12, 8

D. none

Answer: C



[Watch Video Solution](#)

160. If $a \in \mathbb{R}$ and the equation $-3(x - [x]) + a^2 = 0$ (where $[x]$ denotes the greatest integer $\leq x$) has no integral solution, then all possible values of a lie in the interval

- A. $(-2, 1)$
- B. $(-\infty, -2) \cup (2, \infty)$
- C. $(-1, 0) \cup (0, 1)$
- D. $(1, 2)$

Answer: C



[View Text Solution](#)

161. If the roots of the equation $x^2 + kx + 64 = 0$ and $x^2 - 8x + k = 0$ are real then $k > 0$ are real then $k =$

- A. 8

B. 12

C. 16

D. 24

Answer: C



[Watch Video Solution](#)

162. IF the roots of the equation $ax^2 + bx + c = 0$ are real and distinct ,
then

A. both roots are greater than $-b/2a$

B. both roots are less than $-b/2a$

C. one of the roots exceeds $-b/2a$

D. none

Answer: C



[Watch Video Solution](#)

163. If the roots of $ax^2 + bx + c = 0$ are both positive, then

- A. $a < 0, c < 0$
- B. $a < 0, c > 0$
- C. $a > 0, c < 0$
- D. $a > 0, c > 0$

Answer: D



Watch Video Solution

164. IF the roots of $ax^2 + bx + c = 0$ are both negative and $b < 0$ then

- A. $a < 0, c < 0$
- B. $a < 0, c > 0$
- C. $a > 0, c < 0$
- D. $a > 0, c > 0$

Answer: A



[Watch Video Solution](#)

165. IF the roots of $ax^2 + bx + c = 0$ are equal in magnitude but opposite in sign then

- A. $a < 0, c < 0$
- B. $a < 0, c > 0, b > 0$
- C. $a > 0, b = 0, c < 0$
- D. $a > 0, b = 0, c > 0$

Answer: C



[Watch Video Solution](#)

166. if P, q, r are positive and are in A.P the roots of the quadratic $px^2 + qx + r = 0$ real for

A. $\left| \frac{p}{r} - 7 \right| \geq 4\sqrt{3}$

B. $\left| \frac{p}{q} - 7 \right| < 4\sqrt{3}$

C. all p and r

D. no p and r

Answer: A



Watch Video Solution

167. IF $(1 + k)\tan^2 x - 4\tan x - 1 + k = 0$ has real roots $\tan x_1$ and $\tan x_2$ then

A. $k^2 \leq 5$

B. $k^2 \geq 6$

C. $k = 3$

D. none

Answer: A

 [Watch Video Solution](#)

168. IF $0 < a$

A. $|\alpha| = |\beta|$

B. $|\alpha| < 1$

C. $|\beta| < 1$

D. none

Answer: A

 [View Text Solution](#)

169. if $0 < a < b < c < 0$ nad $c > 0$ then both the roots of the equation

$$2ax^2 + 3bx + 5c = 0$$

A. are real and negative

B. have negative real parts

C. have positive real parts

D. none

Answer: B



[View Text Solution](#)

170. IF $p, q, \in \{ 1, 2, 3, 4 \}$ the number of equation of the form $px^2 + qx + 1 = 0$ having real roots is

A. 15

B. 9

C. 7

D. 8

Answer: C



[Watch Video Solution](#)

171. If the equation $x^2 - 2mx + 7m - 12 = 0$ has equal roots then $m =$

A. 2 or 3

B. 3 or 4

C. 4 or 5

D. 5 or 6

Answer: B



[Watch Video Solution](#)

172. IF the roots of $(3m + 1)x^2 + 2(m + 1)x + m = 0$ are equal then $m =$

A. $4/2, 1$

B. $-1/2, 1$

C. $2, 1/2$

D. $2, -1/2$

Answer: B



[Watch Video Solution](#)

173. IF the roots of $x^2 - 2(5 + 2k)x + 3(7 + 10k)$ are equal then $k=$

A. $4/2, 1$

B. $-1/2, 1$

C. $2, 1/2$

D. $2, -1/2$

Answer: C



[Watch Video Solution](#)

174. If the quadratic expression $x^2 - (a - 1)x + \left(a + \frac{1}{4}\right)$ will be a perfect square then $a=$

A. 0, 4

B. 2, 6

C. 2, 4

D. 0, 6

Answer: D



[Watch Video Solution](#)

175. IF $(x - a)(x-b) + (x-b)(x-c) + (x-a)(x-c) = 0$ has equal roots then the relation between a , b and c is

A. $a+b+c=0$

B. $a=b=c$

C. $b^2 = ac$

D. $a + c = 2b$

Answer: B

 [Watch Video Solution](#)

176. If $c^2 \neq ab$ and the roots of $(c^2 - ab)x^2 - 2(a^2 - bc)x + (b^2 - ac) = 0$ are equal, then show that $a^3 + b^3 + c^3 = 3abc$ or $a = 0$

A. $a^3 + b^3 + c^3 = 3abc$ or $a = 0$

B. $a^2 + b^2 + c^2 = ab + bc + ca$

C. $abc = a + b + c$

D. $a = b = c$

Answer: A

 [Watch Video Solution](#)

177. The roots of the equation $(a - b)x^2 + (b - c)x + (c - a) = 0$ are

A. $a, b,$

B. b, c

C. $1, \frac{c - a}{a - b}$

D. $1, \frac{b - c}{a - b}$

Answer: C



Watch Video Solution

178. The roots of the equation $a(b - c)x^2 - b(c - a)x + c(a - b) = 0$ are

A. $ab + bc + ca, 1$

B. $a + b + c, ab + bc + ca$

C. $1, \frac{c(a - b)}{a(b - c)}$

D. $1, \frac{b(c - a)}{a(b - c)}$

Answer: C



Watch Video Solution

179. If $p(q - r)x^2 + q(r - p)x + r(p - q) = 0$ has equal roots then $2/q =$

A. $\frac{1}{p} + \frac{1}{r}$

B. $\frac{1}{p} - \frac{1}{r}$

C. $p + r$

D. pr

Answer: A



Watch Video Solution

180. IF the roots of $(b - c)x^2 + (c - a)x + (a - b) = 0$ are equal then a,

b,c are in

A. AP

B. GP

C. HP

D. AGP

Answer: A



[Watch Video Solution](#)

181. IF the roots of $a(b - c)x^2 + b(c - a)x + c(a - b) = 0$ are equal then a, b, c are in

A. A.P

B. G.P

C. H.P

D. none

Answer: C



[Watch Video Solution](#)

182. If the roots of $(a^2 + b^2)x^2 - 2b(a + c)x + (b^2 + c^2) = 0$ are equal then a, b, c are in

A. A.P

B. G.P

C. H.P

D. none

Answer: B



[Watch Video Solution](#)

183. IF the sum of the roots of $ax^2 + bx + c = 0$ is equal to the sum of the squares of the roots, then.....

A. A.P

B. G.P

C. H.P

D. none

Answer: A



[Watch Video Solution](#)

184. IF α, β are the roots of the equation $ax^2 + bx + c = 0$ and $\alpha < -1, \beta > 1$ then $1 + \frac{c}{a} + \left| \frac{b}{a} \right|$ is

A. positive

B. negative

C. non negative

D. non positive

Answer: B



[View Text Solution](#)

185. IF $3 + 4i$ is a root of the equation $x^2 + px + q = 0$ then

A. $p=6, q=25$

B. $p=6, q=1$

C. $p=-6, q=-7$

D. $p=-6, q=25$

Answer: D



[Watch Video Solution](#)

186. IF $3 + i$ is a root of the equation $x^2 + ax + b = 0$ then $a =$

A. 3

B. -3

C. 6

D. -6

Answer: D



Watch Video Solution

187. IF one root of the quadratic equation $ax^2 + bx + c = 0$ is $3 - 4i$ then $a + b + c =$

A. $40a$

B. $36a$

C. $-20a$

D. $20a$

Answer: D



Watch Video Solution

188. IF $(1 - p)$ is a root of quadratic equation $X^2 + px + (1 - p) = 0$ then its roots are

A. 0, 1

B. -1, 2

C. 0, -1

D. -1, 1

Answer: C



Watch Video Solution

189. IF one root of the equation $x^2 + px + 12 = 0$ is 4, while the equation $x^2 + px + q = 0$ has equal roots then the the value of q is

A. $49/4$

B. 4

C. 3

D. 12

Answer: A



Watch Video Solution

190. If $x^2 - 6x + 5 = 0$ and $x^2 - 12x + p = 0$ have a common root, then find p.

A. 11 or 35

B. 22 or 45

C. 40

D. 10

Answer: A



Watch Video Solution

191. If $x^2 - hx - 21 = 0$, $x^2 - 3hx + 35 = 0$ have a common root then h=

A. ± 2

B. ± 4

C. ± 6

D. ± 8

Answer: B



[Watch Video Solution](#)

192. The value of a such that $x^2 - 11x + a = 0$, $x^2 - 14x + 2a = 0$ may have a common root is

A. 6

B. 12

C. 24

D. 32

Answer: C



[Watch Video Solution](#)

193. IF the equation $x^2 - x - p = 0$ and $x^2 + 2px - 12 = 0$ have a common root then that root is

A. 1

B. $p+2$

C. 2

D. can not be determined

Answer: C



[View Text Solution](#)

194. IF $x^2 + bx + c = 0$, $x^2 + cx + b = 0$ ($b \neq c$) have a common root then $b+c=$

A. 0

B. 1

C. -1

D. 2

Answer: C



[Watch Video Solution](#)

195. IF $ax^2 + 2cx + b = 0$ and $ax^2 + 2bx + c = 0 (b \neq 0)$ have a common root, then $b+c=$

A. $-a/4$

B. $a/3$

C. $a/2$

D. a

Answer: A



[Watch Video Solution](#)

196. IF $x^2 + ax + bc = 0$ and $x^2 + bx + ca = 0$ have a common root ,
then $a + b + c =$

A. 0

B. 1

C. $ab + bc + ca$

D. $3abc$

Answer: A



Watch Video Solution

197. If the quadratic equations
 $ax^2 + 2bx + c = 0$ and $ax^2 + 2cx + b = 0$, ($b \neq c$) have a common
root, then show that $a + 4b + 4c = 0$

A. -2

B. -1

C. 0

D. 1

Answer: C



[Watch Video Solution](#)

198. If $x^2 - cx + d = 0$, $x^2 - ax + b = 0$ have one common root and second has equal roots then $2(b + d) =$

A. 0

B. ac

C. $a+c$

D. $a-c$

Answer: B



[Watch Video Solution](#)

199. IF $x^2 + bx + a = 0$, $ax^2 + x + b = 0$ have a common root and the first equation has equal roots then $2a^2 + b =$

A. 0

B. 1

C. -1

D. 2

Answer: A



Watch Video Solution

200. IF the equations $x^2 + 2x + 3 = 0$ and $ax^2 + bx + c = 0$, $a, b, c \in R$ have a common root then a:b:c is :

A. 1:3:2

B. 3:1:2

C. 1:2:3

D. 3 : 2 : 1

Answer: C

 [Watch Video Solution](#)

201. IF $x^2 + P_1x + q_1 = 0$, $x^2 + p_2x + q_2 = 0$, $x^2 + p_3x + q_3 = 0$ has a common root then $p_1^2 + p_2^2 + p_3^2 + 4(q_1 + q_2 + q_3) =$

A. $2(p_1p_2 + p_2p_3 + p_3p_1$

B. $(p_2p_1 + q_2q_3 + q_3p_1)$

C. $2(q_1p_2 + q_2p_3 + p_3q_1)$

D. none

Answer: A

 [View Text Solution](#)

202. The quadratic equation $x^2 - 6x + a = 0$ and $x^2 - cx + 6 = 0$ have one root in common the other roots of the first and second equations are integers in the ratio 4 : 3 then the common root is

A. 4

B. 3

C. 2

D. 1

Answer: C



[Watch Video Solution](#)

203. If $x^2 + bx + ca = 0$, $x^2 + cx + ab = 0$ have a common root then their other are the roots of the equation

A. $x^2 + 2ax - bc = 0$

B. $x^2 + ax + bc = 0$

C. $x^2 + ax - bc = 0$

D. none

Answer: B

 [Watch Video Solution](#)

204. IF $x^2 + px + q = 0$ and $x^2 + qx + p = 0$ have a common root ,
then their other roots are the roots of

A. $x^2 + x + pq = 0$

B. $x^2 - x - pq = 0$

C. $x^2 - x - pq = 0$

D. none

Answer: A

 [Watch Video Solution](#)

205. IF $(x - 2)$ is a common factor of the expression $x^2 + ax + b$ and $x^2 + cx + d$ then $\frac{b - d}{c - a} =$

A. -2

B. -1

C. 1

D. 2

Answer: D



Watch Video Solution

206. IF α, β are the roots of $ax^2 + bx + c = 0$, $\alpha_1, -\beta$ the roots of $a_1x^2 + b_1x + c_1 = 0$ then α, α are the roots of the equation

A. $\left(\frac{b}{a} + \frac{b_1}{a_1}\right)^{-1} x^2 + x \left(\frac{b_1}{c_1} + \frac{b}{c}\right)^{-1} = 0$

B. $\left(\frac{b}{a} - \frac{b_1}{a_1}\right)^{-1} x^2 - x \left(\frac{b_1}{c_1} + \frac{b}{c}\right)^{-1} = 0$

C. $\left(\frac{b}{a} + \frac{b_1}{a_1}\right)^{-1} x^2 - x \left(\frac{b_1}{c_1} + \frac{b}{c}\right)^{-1} = 0$

D. none

Answer: A



[View Text Solution](#)

207. The values of the parameter a for which the quadratic equation $(1 - 2a)x^2 - 6ax - 1 = 0$ and $ax^2 - x + 1 = 0$ have at least one root in common are

A. $0, 1/2$

B. $1/2, 2/9$

C. $2/9$

D. $0, 1/2, 2/9$

Answer: C



[Watch Video Solution](#)

208. IF $x^2 + ax + b, x^2 + cx + d$ has the common factor $x-1$ then

$$a + b - c - d =$$

A. 0

B. 1

C. -1

D. none

Answer: A



[Watch Video Solution](#)

209. If $ax^2 + 2bx + c = 0, a_1x^2 + 2b_1x + c_1 = 0$ have a common root ,

then the roots of the equation

$$(b^2 - ac)x^2 + (2bb_1 - aa_1 - a_1c)x + (b_1^2 - a_1c_1) = 0$$
 are

A. different

B. equal

C. zero

D. none

Answer: B



[View Text Solution](#)

210. IF $ax^2 + 2bx + c = 0$, and $px^2 + 2qx + r = 0$

A. $b^2 - ac$ and $q^2 - pr$ are both perfect squares

B. $b^2 - ac$ is perfect square and $q^2 - pr$ is not a perfect square

C. $q^2 - pr$ is a perfect square and $b^2 - ac$ is not a perfect square

D. both $b^2 - ac$ and $q^2 - pr$ are not perfect squares

Answer: A



[Watch Video Solution](#)

211. if $P, q, r, s \in \mathbb{R}$ such that $pr = 2(q + s)$ then

- A. both the equation $x^2 + px + q = 0, x^2 + rx + s = 0$ have real roots
- B. one of the equation $x^2 + px + q = 0, x^2 + rx + s = 0$ must have real roots
- C. Both the equations $x^2 + px + q = 0, x^2 + rx + s = 0$ cannot have real roots
- D. none

Answer: B



[Watch Video Solution](#)

212. IF every pair from the equation $x^2 + px + qr = 0, x^2 + qx + rp = 0$ and $x^2 + rx + pq = 0$ has a common root, then the product of three common roots is

A. pqr

B. $2pqr$

C. $p^2q^2r^2$

D. none

Answer: A



Watch Video Solution

213. IF the equation $k(6x^2 + 3) + rz + 2x^2 - 1 = 0$ and $6k(2x^2 + 1) + px + 4x^2 - 2 = 0$ have both the root common, then the value of $2r - p$ is

A. 0

B. 1

C. 2

D. 3

Answer: A



[View Text Solution](#)

214. The equation $ax^2 + bx + a = 0$ ($a, b, \in R$) and $x^3 - 2x^2 + 2x - 1 = 0$ have two roots common. Then $a + b$ must be equal to

A. 1

B. -1

C. 0

D. one of these

Answer: C



[Watch Video Solution](#)

215. The value of 'a' for which the equation $x^3 + ax + 1 = 0$ and $x^4 + ax^2 + 1 = 0$ have a common root is

- A. -2
- B. -1
- C. 1
- D. 2

Answer: A



[Watch Video Solution](#)

216. IF α, β are the roots of $x^2 + px + q = 0$ and also of $x^{2n} + P^n x^n + q^n = 0$ and if $\frac{\alpha}{\beta}, \frac{\beta}{\alpha}$ are the roots of $x^n + 1 + (x + 1)^n = 0$ then n is

- A. an odd integer
- B. an even integer

C. any integer

D. none of these

Answer: B



[View Text Solution](#)

217. IF a, b, c are in A.P. and if

$$(b - c)^2 + (c - a)x + (c - a)x + (a - b) = 0 \quad \text{and}$$

$2(c + a)x^2 + (b + c)x = 0$ have a common root then

A. a^2, b^2, c^2 are in A.P

B. a^2, c^2, b^2 are in A.P

C. a^2, c^2, b^2 are in G.P

D. none of these

Answer: B



[View Text Solution](#)

218. If $12^{4+2x^2} = (24\sqrt{3})^{3x^2-2}$, then $x =$

A. $\pm \sqrt{\frac{13}{12}}$

B. $\pm \sqrt{\frac{14}{5}}$

C. $\pm \sqrt{\frac{12}{13}}$

D. $\pm \sqrt{\frac{5}{14}}$

Answer: B



Watch Video Solution

219. for the equation $|x|^2 + |x| - 6 = 0$ the roots are

A. One and only one real number

B. real with sum one

C. real with sum zero

D. real with product zero

Answer: C

 [Watch Video Solution](#)

220. The number of real solution $x^2 - 7|x| + 12 = 0$ is

A. 4

B. 3

C. 2

D. 10

Answer: A

 [Watch Video Solution](#)

221. The number of real roots of $|x^2| - 5|x| + 6 = 0$ is

A. 2

B. 3

C. 4

D. 1

Answer: C



Watch Video Solution

222. If $|x^2| + |x| - 2 = 0$ then $x =$

A. ± 1

B. ± 3

C. ± 11

D. ± 15

Answer: A



Watch Video Solution

223. The product and sum of the roots of the equation $|x^2| - 5|x| - 24 = 0$ are respectively

- A. $-64, 0$
- B. $-24, 5$
- C. $5, -24$
- D. $0, 72$

Answer: A



[Watch Video Solution](#)

224. The real roots of the equation $|x^2 + 4x + 3| + 2x + 5 = 0$ are

- A. $4, -1 + \sqrt{3}$
- B. $-4, -1 - \sqrt{3}$
- C. $\pm 41 \pm \sqrt{3}$
- D. $-2, -4 - 1\sqrt{3}$

Answer: B



[Watch Video Solution](#)

225. The number of real solutions of the equation

$$|x|^2 - 5|x| + 6 = 0 \text{ is}$$

A. 4

B. 1

C. 3

D. 2

Answer: A



[Watch Video Solution](#)

226. The solutions of $|x^2 - 2x + 2| = 3x - 2$ are

A. 4, - 1

B. - 4, - 1

C. - 4, 1

D. 4, 1

Answer: D

 [Watch Video Solution](#)

227. IF $|x - 2| + 2|x - 9| = 7$ then $x =$

A. any real number between 0.7

B. any real number between 2.9

C. any real number between 2.7

D. any real number between 0.9

Answer: B

 [Watch Video Solution](#)

228. IF $|x - 2| + 2|x - 3| = 7$ then $x =$

A. 2 or $1/3$

B. 3 or $1/7$

C. 2 or $1/5$

D. none

Answer: D



Watch Video Solution

229. If $|x + 1| - |x| + 3|x - 1| - 2|x - 2| = x + 2$, then $x =$

A. $x = 2$ or $x \leq 2$

B. $x = -5$ or $x > 5$

C. $x = -2$ or $x \geq 2$

D. none of these

Answer: C



[View Text Solution](#)

230. The number of real roots of $(7 + 4\sqrt{3})^{|x| - 8} = 14$ is

A. 0

B. 2

C. 4

D. none of these

Answer: D



[View Text Solution](#)

231. The sum of all real values of x satisfying the equation

$$(x^2 - 5x + 5)^{x^2 + 4x - 60} = 1 \text{ is}$$

A. 3

B. -4

C. 6

D. 5

Answer: A



[Watch Video Solution](#)

232. The product of real roots of the equation

$$|x|^{6/5} - 26|x|^{3/5} - 27 = 0 \text{ is}$$

A. -3^{10}

B. -3^{12}

C. $-3^{12/5}$

D. $-3^{21/5}$

Answer: A



Watch Video Solution

233. IF α is a root of the equation $4x^2 + 2x - 1 = 0$ then $4\alpha^3 - 3\alpha$ is

- A. a root
- B. not a root
- C. may be a root
- D. none

Answer: A



Watch Video Solution

234. The value of the continued fraction $1 + \frac{1}{1 + \frac{1}{1 + \dots}}$ is

A. $\frac{\sqrt{5} - 1}{2}$

B. $\frac{\sqrt{5} + 1}{2}$

C. $\frac{\sqrt{5} - 1}{2}$

D. $\frac{\sqrt{5} + 1}{3}$

Answer: B



Watch Video Solution

235. IF $20^{3-2x^2} = (40\sqrt{5})^{3x^2-2}$, then $x =$

A. $\pm \sqrt{\frac{13}{12}}$

B. $\pm \sqrt{12/13}$

C. $\pm \sqrt{4/5}$

D. $\pm \sqrt{5/4}$

Answer: B



Watch Video Solution

236. If $x + \sqrt{x} = 6/25$ then $x =$

A. $1/5$

B. $1/25$

C. $1/625$

D. $1/125$

Answer: B



[Watch Video Solution](#)

237. If $\sqrt{x+1} - \text{sqrt}(x-1) = 1$ then $x =$

A. $7/11$

B. $2/3$

C. $5/4$

D. $-(3/2)$

Answer: C



Watch Video Solution

238. The solution set of $\sqrt{x+1} + \sqrt{2x-5} = 3$ is

A. {2}

B. {3}

C. {4}

D. {5}

Answer: B



Watch Video Solution

239. If $\sqrt{x+2} = \sqrt{3x-10}$, then $x=$

A. -6

B. 6

C. 3

D. -3

Answer: B



[Watch Video Solution](#)

240. The solution set of $\sqrt{x + 20} + \sqrt{x + 4} = 4\sqrt{x - 1}$ is

A. $\{2\}$

B. $\{3\}$

C. $\{4\}$

D. $\{5\}$

Answer: D



[View Text Solution](#)

241. IF $\sqrt{3x - 5} = 0$ then $x =$

A. 2

B. -2

C. $\frac{5}{3}$

D. none

Answer: C



[Watch Video Solution](#)

242. IF $x^2 - 4x - 12\sqrt{x^2 - 4x + 19} + 51 = 0$ then $x =$

A. 1 or 3 or -5 or 9

B. 2 or -3 or 5 or 7

C. -3 or 1 or -7

D. none

Answer: A



[View Text Solution](#)

243. IF $\sqrt{x^2 + 4a + 5} + \sqrt{x^2 + 4b + 5} = 2(a - b)$ then $x =$

A. $\frac{(a - b)^2 - 5}{2(a + b)}$

B. $\frac{a^2 - b^2}{2(a + b)}$

C. $\frac{(a + b)^2}{(a + b) - 5}$

D. none

Answer: A



[View Text Solution](#)

244. the solution set of $x^2 + x - 2 = 0$ is



Watch Video Solution

245. IF $2x^{1/3} + 2x^{-1/3} = 5$ then $x =$

A. 3 or $1/9$

B. 5 or $1/5$

C. 2 or $1/5$

D. 8 or $1/8$

Answer: D



Watch Video Solution

246. IF $(a + x)^{2/3} + (a - x)^{2/3} = 4(a^2 - x^2)^{1/3}$ then $x =$

A. $\pm \frac{3a}{4\sqrt{4}}$

B. $\pm \frac{4a}{3\sqrt{3}}$

C. $\pm \frac{5a}{3\sqrt{3}}$

D. $\pm \frac{5a}{6\sqrt{6}}$

Answer: C



View Text Solution

247. The number of rational roots of

$$(2x + 3)(2x + 5)(x - 1)(X - 2) = 30 \text{ is}$$

A. 4

B. 3

C. 2

D. 5

Answer: C



Watch Video Solution

248. The solution set of $(x - 1)(x - 3)(x - 5)(x - 7) = 9$ is

- A. $\{-4, 4, 4 \pm \sqrt{10}\}$
- B. $\{4, 4, 4 \pm \sqrt{10}\}$
- C. $\{-4, -4, 4 \pm \sqrt{10}\}$
- D. none

Answer: B



[View Text Solution](#)

249. The roots of $\frac{1}{(x - 1)^2} / (x - 2) = \frac{1}{x - 3}$ are

- A. $3 \pm \sqrt{2}$
- B. $2 \pm \sqrt{3}$
- C. $6 \pm \sqrt{8}$
- D. $8 \pm \sqrt{6}$

Answer: A



[View Text Solution](#)

250. The equation $x - \frac{2}{x-1} = 1 - \frac{2}{x-1}$ has

- A. no root
- B. one root
- C. two root
- D. infinitely many roots

Answer: A



[Watch Video Solution](#)

251. If $\frac{x}{b} + \frac{b}{x} = \frac{a}{b} + \frac{b}{a}$ then $x =$

- A. a^2 or b^2/a^3

B. a or b^2/a

C. a^2 or b/a

D. a or b^2/a^2

Answer: B



[View Text Solution](#)

252. A root of the equation $\frac{a+c}{x+a} + \frac{b+c}{x+b} = \frac{2(a+b+c)}{x+a+b}$ is

A. a

B. b

C. c

D. $a+b+c$

Answer: C



[Watch Video Solution](#)

253. The solution set of $\left(x + \frac{1}{x}\right)^2 - \frac{3}{2}\left(x - \frac{1}{x}\right) = 4$ when $x \neq 0$ is

- A. $\{1/2, 1, 1, 2\}$
- B. $\{-1/2, -1/2, -1, 1, 2\}$
- C. $\{1/2, -1, 1, 2\}$
- D. $\{-1/2, 1, 1, 2\}$

Answer: B



Watch Video Solution

254. Solve $2\left(x + \frac{1}{x}\right)^2 - 7\left(x + \frac{1}{x}\right) + 5 = 0$, when $x \neq 0$

- A. $\left\{2 \pm \frac{1}{2}, \frac{1 \pm i\sqrt{3}}{2}\right\}$
- B. $\left\{2, -\frac{1}{2}, \frac{1 + I\sqrt{3}}{2}\right\}$
- C. $\left\{-2, \frac{1}{2}, \frac{1 \pm i\sqrt{3}}{2}\right\}$

D. none

Answer: A

 Watch Video Solution

255. Solve $\left(x^2 + \frac{1}{x^2}\right) - 5\left(x + \frac{1}{x}\right) + 6 = 0$, when $x \neq 0$

A. $\left\{2 \pm \frac{\sqrt{3}}{2}\right\}$

B. $\left\{2, -\frac{1}{2}, \frac{1 + I\sqrt{3}}{2}\right\}$

C. $\left\{-2 \pm \sqrt{3}, \frac{1 + I\sqrt{3}}{2}\right\}$

D. none

Answer: A

 Watch Video Solution

256. If $\frac{\sqrt{1+x^2} + \sqrt{1-x^2}}{\sqrt{1+x^2} - \sqrt{1-x^2}} = 3$ then $x =$

A. $\sqrt{2/3}$

B. $\sqrt{1/3}$

C. $\sqrt{\frac{2}{5}}$

D. $\sqrt{3/5}$

Answer: D



Watch Video Solution

257. IF $\frac{x + \sqrt{12a - x}}{x - \sqrt{12a - x}} = \frac{\sqrt{a} + 1}{(\sqrt{a} - 1)}$, then x=

A. $2a^2$

B. $4a$

C. $3a$

D. $3a^2$

Answer: C



Watch Video Solution

258. IF $\sqrt{\frac{x}{1-x}} + \sqrt{\frac{1-x}{x}} = 2\frac{1}{6}$ then x=

A. $4/11$ or $7/11$

B. $3/11$ or $5/11$

C. $4/13$ or $9/13$

D. none

Answer: C

 [Watch Video Solution](#)

259. $\sqrt{\frac{x}{x-3}} + \sqrt{\frac{x-3}{x}} = \frac{5}{2}$, when $x \neq 0$ and $x \neq 3$

A. $\{1, 2\}$

B. $\{1, -1\}$

C. $\{1, 5\}$

D. $\{4, -1\}$

Answer: D

 [Watch Video Solution](#)

260. $\sqrt{\frac{3x}{x+1}} + \sqrt{\frac{x+1}{3x}} = 2$, when $x \neq 0$ and $x \neq -1$

A. $\{1/2\}$

B. $\{1\}$

C. $\{2\}$

D. $\{3\}$

Answer: A

 [Watch Video Solution](#)

261. If $2^x + 27(2^{-x}) = 12$ then $x =$

A. $\log_2 3$ or $2 \log_2 3$

B. $\log_2 5$ or $2\log_2 3$

C. $\log_2 7$ or $\log_2 5$

D. none

Answer: A

 [Watch Video Solution](#)

262. If $3^{2x} - 3^{x+1} - 3^{x-1} + 1 = 0$ then $x =$

A. 0

B. ± 7

C. ± 1

D. ± 3

Answer: C

 [Watch Video Solution](#)

263. Solve $7^{1+x} + 7^{1-x} = 50$ for real x .

- A. $\{1, 1\}$
- B. $\{1, -1\}$
- C. $\{-1, 1\}$
- D. none

Answer: B

 [Watch Video Solution](#)

264. IF $e^{(\cos^2 x + \cos^4 + \cos^6 x + \dots) \log 3}$ satisfies $y^2 - 10y + 9 = 0$ and

$(0 < x < \pi/2)$ then $\cot^2 x =$

- A. 0
- B. 1
- C. $1/2$
- D. 9

Answer: A



[Watch Video Solution](#)

265. The equation $x^{(3/4)} (\log_2 x)^2 + \log_2 x - 5/4 = \sqrt{2}$ has

- A. exactly two real roots
- B. no real root
- C. one irrational root
- D. one of these

Answer: C



[Watch Video Solution](#)

266. The number of real of the equation $\frac{2x - 3}{x - 1} + 1 = \frac{6x^2 - x - 6}{x - 1}$ is

- A. 0

B. 1

C. 2

D. none

Answer: B



[Watch Video Solution](#)

267. The equation $e^{\sin x} - e^{-\sin x} - 4 = 0$ has

A. 0

B. 1

C. 4

D. ∞

Answer: A



[Watch Video Solution](#)

268. The number of solutions of the system of equation given below is :

A. ∞

B. 2

C. 4

D. 8

Answer: D



[View Text Solution](#)

269. The number of solution of the equation $9x^2 - 18|x| + 5 = 0$

belonging to the domin of $\log_e \{(x + 1)(x + 2)\}$ is

A. 1

B. 2

C. 3

D. 4

Answer: C



Watch Video Solution

270. The number of solutions of the equation

$$5^x + 5^{-x} = \log_{10} 25, x \in R \text{ is}$$

A. 0

B. 1

C. 2

D. infinitely many

Answer: A



Watch Video Solution

271. IF the equation $ax+by=1, cx^2 + dy^2 = 1$ have only one solution then

$$\frac{a^2}{c} + \frac{b^2}{d} =$$

A. 1

B. -1

C. 0

D. 2

Answer: A



Watch Video Solution

272. If $a+b+c=0$ then the quadratic equation $3ax^2 + 2bx + c = 0$ has at least one root in

A. $[1, 2]$

B. $[0, 1]$

C. $[-1, 1]$

D. $[1, 3]$

Answer: B

 [Watch Video Solution](#)

273. If $2a + 3b + 6c = 0$ then the equation $ax^2 + bx + c = 0$ has atleast one root in

- A. $(0, 1)$
- B. $(1, 1)$
- C. $[-1, 1]$
- D. $[1, 2]$

Answer: A

 [Watch Video Solution](#)

274. Let $P(x)$ be a polynomial with integral coefficients . If there exists two integers a and b such that $p(a) - p(b) = 1$ then

- A. both a and b must be even

B. both a and b must be odd

C. a and b are two consecutive integers

D. none of these

Answer: C



[View Text Solution](#)

275. IF $0 < a$

A. real and distinct roots out of which one lies between c and d

B. real and distinct roots out of which one lies between a and b

C. real and distinct roots out of which one lies between b and c

D. non - real roots

Answer: A



[View Text Solution](#)

276. α and β are the roots of the equation $x^2 + px + p^3 = 0$, ($p \neq 0$). If the point (α, β) lies on the curve $x = y^2$, then the roots of the given equation are

A. 4, -2

B. 4, 2

C. 1, -1

D. 1, 1

Answer: A



[Watch Video Solution](#)

277. Find all number which exceed their square root by 12

A. 8

B. 16

C. 24

D. 32

Answer: B



[Watch Video Solution](#)

278. The two consecutive positive odd integers such that the sum of their squares is 290 are

A. 5, 7

B. 9, 11

C. 11, 13

D. 15, 17

Answer: C



[Watch Video Solution](#)

279. Find two consecutive positive even integers, the sum of whose squares is 340.

A. 10, 20

B. 12, 14

C. 14, 18

D. 16, 20

Answer: B



[Watch Video Solution](#)

280. The number having two digits such that it is 4 times the sum and three times the product of its two digits are

A. 8

B. 16

C. 24

D. 32

Answer: C



[Watch Video Solution](#)

281. A number of two digit s whose product is 30 . If the digits are interchanged the resulting number will exceed the previous by 9. the number is

A. 56

B. 54

C. 38

D. 28

Answer: A



[Watch Video Solution](#)

282. The sum of the ages of a father and a son is 45 years . Five years ago the product of their ages was four times the father's age at that time .

Their present ages are

A. 32, 8

B. 28, 7

C. 36, 9

D. 40, 10

Answer: C



[Watch Video Solution](#)

283. The cost of a piece of cable wire is Rs. 35/-, If the length of the piece of wire is 4 meters more and each meter costs, Rs. 1/- less, the cost would remain un-changed. What is the length of the wire ?

A. 10 metres

B. 12 metes

C. 15 metres

D. 20 metres

Answer: A



Watch Video Solution

284. One fourth of a herd of goats was seen in the forest. Twice the square root of the number in the herd had gone up the hill and the remaining 15 goats were on the bank of the river. Find the total number of goats.

A. 26

B. 28

C. 34

D. 36

Answer: D



Watch Video Solution

285. In the interior of a forest there are some apes . Of their total number square of $\frac{1}{9}$ th are playing at one place . The remaining are on the hills .
The total number of apes is

- A. 27 or 54
- B. 16 or 32
- C. 28 or 56
- D. 185 or 36

Answer: A

[View Text Solution](#)

286. In a cricket match Anil took one wicket less than twice the number of wickets taken by Ravi. If the product of the number of wickets taken by them is 15, find the number of wickets taken by each of them.

A. 5, 3

B. 3, 5

C. 2, 6

D. 7, 9

Answer: A



[Watch Video Solution](#)

287. Some points on a plane are marked and they are connected pair wise by line segments . IF the total number of line segments formed is 10 then the number of marked points on the plane is

A. 2

B. 3

C. 4

D. 5

Answer: D



[Watch Video Solution](#)

288. The sides of a right angled triangle containing the right angle are $5x$ cm and $(3x-1)$ cm . If the area of the triangle is 60 sq . Cm the length of the sides of the triangles are

A. 8 cm , 15 cm , 17 cm

B. 6cm , 12 cm , 18 cm

C. 10 cm , 18 cm , 20 cm

D. 9 cm , 116 , 24 cm

Answer: A



[Watch Video Solution](#)

1. If $x < 3$ or $x > 4$ then the value of $x^2 - 7x + 12$ is

A. zero

B. positive

C. negative

D. not determined

Answer: B



Watch Video Solution

2. IF $x \in R$ then the value of $x^2 - 6x + 10$ is

A. zero

B. positive

C. negative

D. not determined

Answer: B



[Watch Video Solution](#)

3. IF $-1/2 < x < 3$ then the value of $2x^2 - 5x - 3$ is

A. zero

B. positive

C. negative

D. not determined

Answer: C



[Watch Video Solution](#)

4. if $x \neq 3/2$ then the value of $4x^2 - 12x + 9$ is

A. zero

B. positive

C. negative

D. not determined

Answer: B



[Watch Video Solution](#)

5. IF $x \in R$ then the value of $4x - x^2 - 6$ is

A. zero

B. positive

C. negative

D. not determined

Answer: C



[Watch Video Solution](#)

6. IF $4 < x < 8$ then the value of $-7x^2 + 8x - 9$ is

- A. zero
- B. positive
- C. negative
- D. not determined

Answer: C



[Watch Video Solution](#)

7. If $x \in R$ then the value of $-7x^2 + 9x - 9$ is

- A. zero
- B. positive
- C. negative
- D. not determined

Answer: C



Watch Video Solution

8. if $a > 0$ and $b^2 - 4ac = 0$, then the curve $y = ax^2 + bx + c$

- A. Cuts the x- axis
- B. touches the x- axis lies below it
- C. lies entirely above the x- axis
- D. touches the x- axis and lies above it

Answer: D



Watch Video Solution

9. $x^2 - 2x + 10$ has minimum at $x =$

- A. 2

B. -1

C. 1

D. -2

Answer: C

 [Watch Video Solution](#)

10. $3x - 5x^2 + 12$ has maximum at $x =$

A. $2/5$

B. $-1/5$

C. $3/10$

D. $-3/10$

Answer: C

 [Watch Video Solution](#)

11. if $2x - 7 - 5x^2$ has maximum value at $x = a$ then $a =$

A. $-1/5$

B. $1/5$

C. $34/5$

D. $-34/5$

Answer: B



[Watch Video Solution](#)

12. the minimum value of $x^2 - 8x + 17, \forall x \in R$ is

A. 17

B. -1

C. 1

D. 2

Answer: C



Watch Video Solution

13. The maximum value of $10x - 5x^2 - 1$ is

A. -1

B. $-1/5$

C. 2

D. 4

Answer: D



Watch Video Solution

14. the maximum value of $(x - a)(b - x)$ is

A. $(a^2 - b)^2 / 4$

B. $(a - b)^2$

C. a

D. b

Answer: A



Watch Video Solution

15. The maximum value of $a^2 - abx - b^2x^2$ is

A. $5a^2/4$

B. $a^2/2$

C. a

D. $-a$

Answer: A



Watch Video Solution

16. the minimum value of $\left(x - \frac{5}{3}\right)^2 + \frac{7}{2}$ is

A. 1

B. 7

C. $7/5$

D. $7/2$

Answer: D



[Watch Video Solution](#)

17. the maximum value of $\frac{7}{5} - \left(x - \frac{2}{3}\right)^2$ is

A. $11/3$

B. $7/5$

C. 7

D. 5

Answer: B



Watch Video Solution

18. the maximum value of $c + 2bx - x^2$ is

A. b^2c

B. $b^2 - c$

C. $c - b^2$

D. $b^2 + c$

Answer: D



Watch Video Solution

19. the minimum value of the quadratic expression $x^2 + 2bx + c$ is

A. cb^2

B. c^2b

C. $c + b^2$

D. $c - b^2$

Answer: D



Watch Video Solution

20. the expression $2x^2 + 4x + 7$ has minimum value m at $x = \alpha$. The ordered pair (α, m) is

A. $(1, 5)$

B. $(1, -5)$

C. $(-1, -5)$

D. $(-1, 5)$

Answer: D



Watch Video Solution

21. The extreme value of $x^2 - 5x + 6$ is

A. $1/4$

B. $-1/4$

C. $1/2$

D. $-1/2$

Answer: B



[Watch Video Solution](#)

22. Find the changes in the sign of the following expressions and find their extreme values.

$$15 + 4x - 3x^2$$

A. $49/3$

B. $-49/3$

C. $47/3$

D. $-47/3$

Answer: A



Watch Video Solution

23. If a_1, a_2, \dots, a_n are in H.P. then

$$a_1 \cdot a_2 + a_2 \cdot a_3 + a_3 \cdot a_4 + \dots + a_{n-1} \cdot a_n =$$

A. $a_1 + a_2 + \dots + a_n$

B. $(a_1 + a_2 + a_3 + \dots + a_n)$

C. $n(a_1 + a_2 + \dots + a_n)$

D. none

Answer: D



Watch Video Solution

24. If a, b, c are positive then the least value of $(a + b + c)(1/a + 1/b + 1/c)$ is

A. 4

B. 3

C. 7

D. 9

Answer: D



[Watch Video Solution](#)

25. If a, b, c are distinct positive numbers then the expression $(b + c - a)(c + a - b)(a + b - c) - abc$ is

A. positive

B. negative

C. nonpositive

D. nonnegative

Answer: C

 [View Text Solution](#)

26. IF $a^2 + b^2 + c^2 = 1$ then the range of $ab + bc + ca$ is

A. $[1/2, 2]$

B. $[-1/2]$

C. $[-1/2, 1]$

D. $[1, 3/2]$

Answer: C

 [Watch Video Solution](#)

27. In triangle ABC, range of $\frac{a^2 + b^2 + c^2}{ab + bc + ca}$ is (a,b,c are sides of triangle)

A. $[1,2)$

B. $(-\infty, 1] \cup [2, \infty)$

C. zero

D. none

Answer: A

 [Watch Video Solution](#)

28. The solution set of $-x^2 + 3x + 4 > 0$ is

A. $(-1, 4)$

B. $(-\infty, -3] \cup [7, \infty)$

C. $(-\infty, 3) \cup (5, \infty)$

D. $[-4, 1]$

Answer: A

 [Watch Video Solution](#)

29. the solution set of $x^2 - 4x - 21 \geq 0$ is

- A. $(-1, 4)$
- B. $(-\infty, -3] \cup [7, \infty)$
- C. $(-\infty, 3) \cup (5, \infty)$
- D. $[-4, 1]$

Answer: B



[Watch Video Solution](#)

30. the solution set of $1 \leq x^2 - 2x$ is

- A. $(-\infty, 1 - \sqrt{2}] \cup [1 + \sqrt{2}, \infty)$
- B. $(-\infty, -3) \cup (1 + \sqrt{2}, \infty)$
- C. $(-1, 1/2)$

D. $[-1, 1/2]$

Answer: A



Watch Video Solution

31. the solution set of $x^2 > 4x + 5$ is

A. $(-\infty, 1 - \sqrt{2}) \cup (1 + \sqrt{2}, \infty)$

B. \mathbb{R}

C. $(-1, 1/2)$

D. $[-1, 1/2]$

Answer: B



Watch Video Solution

32. The solution set of $2x^2 - 4x + 5 > 0$ is

A. $[-1/2, 3]$

B. \mathbb{R}

C. $(-2, 1/2)$

D. ϕ

Answer: D



[Watch Video Solution](#)

33. the solution set of $x^2 - 2x + 2 < 0$ is

A. $[-1/2, 3]$

B. \mathbb{R}

C. $(-2, 1/2)$

D. ϕ

Answer: D



[Watch Video Solution](#)

34. the set of all solutions of the inequation $x^2 - 2x + 5 \leq 0$ in \mathbb{R} is

A. $\mathbb{R} - (-\infty, -5)$

B. $\mathbb{R} - (5, \infty)$

C. ϕ

D. $\mathbb{R} - (-\infty, -4)$

Answer: C



Watch Video Solution

35. The set of solutions of $|x|^2 - 5|x| + 4 < 0$ is

A. $(-4, -1)$

B. $(1, 4)$

C. $(-4, -1) \cup (1, 4)$

D. $(-4, 4)$

Answer: C



[Watch Video Solution](#)

36. if the expression $4x - 5x^2 + 1$ is positive if x lies in

A. $(-1/5, 1)$

B. $(1, 1)$

C. ϕ

D. R

Answer: A



[Watch Video Solution](#)

37. the expression $-7x^2 + 8x - 9$ is positive if x lies in

A. $(-\infty, 1) \cup (2, \infty)$

B. $(-\infty, 2) \cup (3, \infty)$

C. ϕ

D. R

Answer: C

 [Watch Video Solution](#)

38. the expression $x^2 - 5x - 6$ is negative is x lies in

A. $(-1, 6)$

B. $(2, 5)$

C. $(-3, 1/2)$

D. R

Answer: A

 [Watch Video Solution](#)

39. The greatest positive integral value of x for which $200 - x(10 + x)$ is positive is

A. 11

B. 10

C. 9

D. none

Answer: C



[Watch Video Solution](#)

40. The least integral value of x for which $33 - x(2 + 3x) > 0$ is

A. -11

B. -3

C. -2

D. - 1

Answer: B



[Watch Video Solution](#)

41. The integer k for which the inequality $x^2 - 2(4k - 1)x + 15k^2 - 2k - 7 > 0$ is valid for any x , is

A. 2

B. 3

C. 4

D. 5

Answer: B



[Watch Video Solution](#)

42. IF the difference between the roots the roots of the equation $x^2 + ax + 1 = 0$ is less than $\sqrt{5}$ then the set of possible values of a is

- A. $(-3, 3)$
- B. $(-3, \infty)$
- C. $(3, \infty)$
- D. $(-\infty, -3)$

Answer: A



[Watch Video Solution](#)

43. IF $x^2 + 6x - 27 > 0 - x^2 + 3x + 4 > 0$ then x lies in the interval

- A. $(3, 4)$
- B. $[3, 4]$
- C. $(-\infty, 3] \cup [4, \infty)$
- D. $(-9, 4)$

Answer: A



Watch Video Solution

44. The set of values of x for which the inequalities $x^2 - 3x - 10 < 0$, $10x - x^2 - 16 > 0$ hold simultaneously, is

A. $(-2, 5)$

B. $(2, 8)$

C. $(-2, 8)$

D. $(2, 5)$

Answer: B



Watch Video Solution

45. The set of values of x for which the inequalities $x^2 - 2x + 3 > 0$, $2x^2 + 4x + 3 > 0$ hold simultaneously, is

A. $(-4, 1)$

B. $(-4, -3] \cup [-2, 1)$

C. $(-4, -3) \cup (-2, 1)$

D. \mathbb{R}

Answer: D



[Watch Video Solution](#)

46. IF $x^2 - 2x + 3 > 0$, $2x^2 + 4x + 3 > 0$ then x lies in the interval

A. $(1, 2)$

B. \mathbb{R}

C. $(2, 5)$

D. ϕ

Answer: B



[Watch Video Solution](#)

47. the greatest negative integer satisfying $x^2 - 4x - 77 < 0$ and $x^2 > 4$ is

A. 1

B. 2

C. 3

D. -3

Answer: D



[Watch Video Solution](#)

48. If $x > 0$ then the least value of $x + \frac{1}{x}$ is

A. 2

B. -2

C. 1

D. 0

Answer: A



Watch Video Solution

49. the least value of $\cos^2 x + \sec^2 x$ is

A. 0

B. -2

C. 2

D. -1

Answer: C



Watch Video Solution

50. The range of $\frac{x^2 + 2x + 1}{x^2 + 2x - 1}$ is

A. $(-\infty, 0] \cup (1, \infty)$

B. $[1/2, 2]$

C. $(-\infty, -2/9] \cup (1, \infty)$

D. $(-\infty, -6) \cup (-2, \infty)$

Answer: A

 [Watch Video Solution](#)

51. The range of $\frac{x^2 - 2x + 9}{x^2 + 2x + 9}$ is

A. $(-\infty, 0] \cup (1, \infty)$

B. $[1/2, 2]$

C. $(-\infty, -2/9] \cup (1, \infty)$

D. $(-\infty, -6) \cup (-2, \infty)$

Answer: B

 [Watch Video Solution](#)

52. The range of $\frac{x^2 - 2x + 3}{x^2 - 2x - 8}$ is

A. $[-9/2, 1/2]$

B. $(-\infty, 1] \cup [2, \infty)$

C. $(-\infty, -9] \cup [-1, \infty)$

D. R

Answer: C



Watch Video Solution

53. The range of $\frac{x^2 - 2x + 3}{x^2 - 2x - 8}$ is

A. $[-9/2, 1/2]$

B. $(-\infty, 1] \cup [2, \infty)$

C. $(-\infty, -9] \cup [-1, \infty)$

D. R

Answer: D



Watch Video Solution

54. IF x is real then the value of $\frac{x^2 - 3x + 4}{x^2 + 3x + 4}$ lies in the interval

A. $\left[\frac{1}{3}, 3 \right]$

B. $\left[\frac{1}{5}, 5 \right]$

C. $\left[\frac{1}{6}, 6 \right]$

D. $\left[\frac{1}{7}, 7 \right]$

Answer: D



Watch Video Solution

55. Show that none of the values of the function $\frac{x^2 + 34x - 71}{x^2 + 2x - 7}$ over \mathbb{R} lies between 5 and 9.

A. does not lie

B. lies

C. none

D. cannot be determined

Answer: A



[Watch Video Solution](#)

56. IF x is real , then $\frac{x^2 - bc}{2x - b - c}$ has _____ values between b and c

A. no real

B. real

C. none

D. cannot be determined

Answer: A

 [Watch Video Solution](#)

57. If x is real then $y = \frac{2x^2 + 6x + 5}{x^2 + 3x + 2}$ ---- between -2 and +2

A. lies

B. does not lie

C. none

D. cannot be determined

Answer: B

 [Watch Video Solution](#)

58. If x is real, then the minimum value of $y = \frac{x^2 - x + 1}{x^2 + x + 1}$ is

A. $1/3$

B. 3

C. $1/2$

D. 1

Answer: A

 [Watch Video Solution](#)

59. the (relative) minimum value of $\frac{x^2 - 3x + 2}{x^2 + 3x + 2}$ is

A. $-1/11$

B. $-17 + 12\sqrt{2}$

C. $-17 - 12\sqrt{2}$

D. 0

Answer: B

 [View Text Solution](#)

60. For $x \in R$, the least value of $\frac{x^2 - 6x + 5}{x^2 + 2x + 1}$ is

A. -1

B. $-\frac{1}{2}$

C. $-\frac{1}{4}$

D. $-\frac{1}{3}$

Answer: D



Watch Video Solution

61. If $x \in R$, then the range of $\frac{x}{x^2 - 5x + 9}$ is

A. $-1/11$

B. -1

C. $1/11$

D. 1

Answer: D



[Watch Video Solution](#)

62. If x is real, then the maximum value of $\frac{x^2 + 14x + 9}{x^2 + 2x + 3}$ is

A. 2

B. 4

C. 6

D. 8

Answer: B



[Watch Video Solution](#)

63. the limits of $\frac{6x^2 - 18x + 21}{6x^2 - 18x + 17}$ are

A. 1, 15/7

B. 1, 15

C. 2, 3/5

D. 1, 7/15

Answer: A



[View Text Solution](#)

64. If x is real, then the range of $\frac{x^2 + 2x + 1}{x^2 + 2x + 7}$ is

A. 0, 1

B. 1, 2

C. 0, 2

D. 2, 3

Answer: A



[Watch Video Solution](#)

65. IF $\frac{x - a}{x^2 - 3x + 2}$ takes all real values for $x \in R$, then

A. $a = 2$

B. $a < 2$

C. $1 < a < 2$

D. $1 \leq a \leq 2$

Answer: D



[View Text Solution](#)

66. IF $x \in R$ then $\frac{x^2 + 2x + a}{x^2 + 4x + 3a}$ can take all real values if

A. $a \in (0, 2)$

B. $a \in [0, 1]$

C. $a \in [-1, 1]$

D. none

Answer: B



[View Text Solution](#)

67. If $Y = \tan x \cot 3x$, $x \in R$, then

A. $\frac{1}{3} < y < 1$

B. $\frac{1}{3} \leq y \leq 1$

C. $\frac{1}{3} \leq y \leq 3$

D. none

Answer: D



[Watch Video Solution](#)

68. If $x \in R$ then $\frac{2a(x-1)\sin^2 \alpha}{x^2 - \sin \alpha}$ cannot lie between

A. $a \sin^2 \alpha, a \cos^2 \alpha$

B. $a \sin^2(\alpha/2), a \cos^2(\alpha/2)$

C. $2a \sin^2 \alpha, 2a \cos^2 \alpha$

D. $2a \sin^2(\alpha/2), 2a \cos^2(\alpha/2)$

Answer: D



[View Text Solution](#)

69. If $a \neq b$ then the expression $x^2 - (a + b)x + (a^2 - ab + b^2)$ ___
negative values for any real value of x

A. does not take

B. take

C. none

D. cannot be determined

Answer: A



[View Text Solution](#)

70. If α, β are the roots of $x^2 - (a - 2)x - (a + 1) = 0$ where a is a variable then the least value of $\alpha^2 + \beta^2$ is

A. 2

B. 3

C. 5

D. 7

Answer: C



[Watch Video Solution](#)

71. IF the sum of the squares of the roots of the equation $x^2 - (\sin \alpha - 2)x - (1 + \sin \alpha) = 0$ is least, then $\alpha =$

A. $\pi/4$

B. $\pi / 3$

C. $\pi / 2$

D. $\pi / 6$

Answer: C



Watch Video Solution

72. The value of a for which the sum of the squares of the roots of the equation $x^2 - (a - 2)x - a - 1 = 0$ assume the least value is

A. 0

B. 1

C. 2

D. 3

Answer: B



Watch Video Solution

73. IF the roots of the equation $bx^2 + cx + a = 0$ be imaginary , then for all real values of x , the experssion $3b^2x^2 + 6bcx + 2c^2$ is

- A. less than $4ab$
- B. greater than $-4ab$
- C. less than $-4ab$
- D. greater than $4ab$

Answer: B

 [Watch Video Solution](#)

74. The smallest value of of the constant $m > 0$ for which $f(x) = 9mx - 1 + \frac{1}{x} \geq 0$ for all $x > 0$, is

- A. $\frac{1}{9}$
- B. $\frac{1}{16}$

C. $\frac{1}{36}$

D. $\frac{1}{81}$

Answer: C



Watch Video Solution

75. The real values of x for which $y = \sqrt{\frac{(x+1)(x-3)}{x-2}}$ takes real values are

A. $-1 \leq x < 2$ or $x \geq 3$

B. $1 < x < 2$ or $x > 2$

C. $x < 2$ or $x > 3$

D. $x > 2$

Answer: A



Watch Video Solution

76. IF $x > -c$ then the minimum value of $\frac{(a+x)(b+x)}{c+x}$ is

A. $\sqrt{a-c} + \sqrt{b-c}$

B. $\sqrt{a-c} - \sqrt{b-c}$

C. $(\sqrt{a-c} + \sqrt{b-c})^2$

D. $(\sqrt{a-c} - \sqrt{b-c})^2$

Answer: C



Watch Video Solution

77. IF $x > -c$ then the minimum value of $\frac{(a+x)(b+x)}{c+x}$ is

A. $\sqrt{a-c} + \sqrt{b-c}$

B. $\sqrt{a-c} - \sqrt{b-c}$

C. $(\sqrt{a-c} + \sqrt{b-c})^2$

D. $(\sqrt{a-c} - \sqrt{b-c})^2$

Answer: D



Watch Video Solution

78. For real x , the function $\frac{(x - a)(x - b)}{x - c}$ will assume all real values provided

A. $a < b < c$

B. $b < c < a$

C. $c < a < b$

D. none of these

Answer: B



Watch Video Solution

79. The range of values of x which satisfy $5x + 2 < 3x + 8$ and $\frac{x + 2}{x - 1} < 4$ are

A. $(-\infty, 1) \cup (2, 3)$

B. $(0, \infty)$

C. $(-\infty, 2)$

D. $(1, 3)$

Answer: A



Watch Video Solution

80. The range of values of x for which the inequality $\frac{x - 1}{4x + 5} < \frac{x - 3}{4x - 3}$

holds is

A. $(-4/3, 5/8)$

B. $(-4/3, 1/2)$

C. $(-5/4, 3/4)$

D. none

Answer: C



Watch Video Solution

81. The values of x for which $\frac{x-1}{3x+4} < \frac{x-3}{3x-2}$ holds, lie in

- A. $(-\infty, -5/4)$
- B. $(-4/3, 2/3)$
- C. $(3/4, \infty)$
- D. $(-\infty, -5/4 \cup (3/4, \infty)$

Answer: B



Watch Video Solution

82. $\left\{ x \in \mathbb{R} : \frac{14x}{x+1} - \frac{9x-30}{x-4} < 0 \right\}$ is equal to

- A. $(-1, 4)$
- B. $(1, 4) \cup (5, 7)$
- C. $(1, 7)$

D. $(-1, 1) \cup (4, 6)$

Answer: D



Watch Video Solution

83. IF the inequation $\sqrt{3x - 8} < -2$ then

A. ϕ

B. $[1, 2]$

C. $[12, \infty)$

D. $(1, 12]$

Answer: A



Watch Video Solution

84. if the inequation $\sqrt{x + 5} < -x$ then

A. $5 < x < 1$

B. $-5 < x < 1$

C. $-5 < x < -1$

D. none

Answer: C

 [View Text Solution](#)

85. If the inequation $\sqrt{x^2 - 18x + 72} < x - 1$ then

A. ϕ

B. $[1, 2)$

C. $[12, \infty)$

D. $(1, 2]$

Answer: C

 [Watch Video Solution](#)

86. IF $\sqrt{9x^2 + 6x + 1} < (2 - x)$ then

A. $x \in \left(+\frac{3}{2}, \frac{1}{4} \right)$

B. $x \in \left(-\frac{3}{2}, \frac{1}{4} \right)$

C. $x \in \left[-\frac{3}{2}, \frac{1}{4} \right)$

D. $x < \frac{1}{4}$

Answer: B



Watch Video Solution

87. If the inequation $\sqrt{(x + 2)(x - 5)} > 8 - x$ then x lies in

A. $\left(\frac{74}{36}, \infty \right)$

B. $\left[\frac{74}{13}, \infty \right)$

C. $\left(-\frac{74}{13}, \infty \right)$

D. $\left[-\frac{74}{13}, \infty\right)$

Answer: A



[View Text Solution](#)

88. If the inequation $\frac{\sqrt{6+x-x^2}}{x+10} \leq \frac{\sqrt{(8-2x-x^2)}}{2x+9}$ then

A. $2 \leq x \leq -1$

B. $-2 \leq x \leq -1$

C. $2 \leq x \leq , x = 3$

D. none

Answer: B



[View Text Solution](#)

89. if the inequation $\frac{\sqrt{8-2x-x^2}}{x+10} \leq \frac{\sqrt{(8-2x-x^2)}}{2x+9}$, then x lies in

A. $[-4, 1] \cup \{2\}$

B. $(-4, 1] \cup \{2\}$

C. $[-4, 1) \cup \{2\}$

D. none

Answer: A

 [View Text Solution](#)

90. if $x^2 + ky^2 + x - y$ is resolvable into two linear factors then $k =$

A. -1

B. 1

C. 2

D. 0

Answer: A

 [Watch Video Solution](#)

91. If $xy + 2x - 3y - k$ is resolvable into two linear factors then $k =$

A. -1

B. 4

C. 6

D. -5

Answer: C



[Watch Video Solution](#)

92. IF $x^2 - y^2 + 4x - 6y + k$ is resolvable into two linear factors then $k =$

A. -1

B. 4

C. 6

D. -5

Answer: D



[Watch Video Solution](#)

93. IF $x^2 + 4xy + 4y^2 + 4x + cy + 3$ can be written as a product of two linear factors , then $c=$

A. 2

B. 4

C. 6

D. 8

Answer: D



[Watch Video Solution](#)

94. IF $12x^2 - mxy + 3y^2 - 5y^2 - 2$ can be resolvable into two linear factors then $m=$

A. ± 6

B. ± 3

C. ± 5

D. ± 7

Answer: D



[View Text Solution](#)

95. IF $12^2 - 10xy + 2y^2 + 11x - 5y + k$ is resolvable into two linear factors factors then $k=$

A. 2

B. 1

C. 0

D. 4

Answer: A



[View Text Solution](#)

96. IF $3x^2 + 8xy - ky^2 + 29x - 3y + 18$ is resolvable into two linear factors then k=

A. 2

B. 1

C. 3

D. 4

Answer: C



[View Text Solution](#)

97. IF $mx^2 + 7xy - 3y^2 + 4x + 7y + 2$ is resolvable into two linear factors then m=of twp linear factors then the factors are

A. 7

B. 4

C. 2

D. -5

Answer: C



[View Text Solution](#)

98. IF $4x^2 + 4xy - ky^2 - 12 - 2y + 8$ can be written as the produ of two linear factors then the factors are

A. $(2x + 3y + 4)(3x + 5y + 2)$

B. $(3y + x + 9)(y - 3x - 2)$

C. $(2x + 3y - 4)(2x - y - 2)$

D. $(x - y + 4), (x - 2y + 5)$

Answer: C



[View Text Solution](#)

99. IF $3x^2 + 8xy + 5y^2 + 14x + 22y + 8$ is resolvable into two linear factors then the factors are

A. $(2x + 3y + 4)(3x + 5y + 2)$

B. $(3y + x + 9)(y - 3x - 2)$

C. $(2x + 3y - 4)(2x - y - 2)$

D. $(x - y + 4), (x - 2y + 5)$

Answer: A



[View Text Solution](#)

100. Examine $3y^2 - 8xy - 3x^2 - 29x + 3y - 18$ is re-solvable into two linear factors.

A. $(2x + 3y + 4)(3x + 5y + 2)$

B. $(3y + x + 9)(y - 3x - 2)$

C. $(2x + 3y - 4)(2x - y - 2)$

D. $(x - y + 4), (x - 2y + 5)$

Answer: B



[Watch Video Solution](#)

101. The condition for $ax^2 + 2cxy + by^2 + 2bx + 2ay + c$ is resolvable into two linear factors is

A. $a^3 + b^3 + c^3 = 3abc$

B. $a^3 + b^3 + c^3 = abc$

C. $a^2 + b^2 + c^2ab + bc + ca$

$$D. a^3 + b^3 + c^3 = 27abc$$

Answer: A



[View Text Solution](#)

102. The condition for $3x^2 + 2pxy + 2y^2 + 2ax - 4y + 1$ can be resolved into two linear factors is

A. $p^2 + 4ap + a^2 + 6 = 0$

B. $p^2 + 4ap + a^2 = 6$

C. $p^2 + 4ap + 2a^2 + 6 = 0$

D. $p^2 + 4ap + 2a^2 = 6$

Answer: C



[Watch Video Solution](#)

103. If $x^2 + 4y^2 - 8x + 12 = 0$ is satisfied by real values of x and y then y must lie between

A. 2, 6

B. 2, 5

C. -1, 1

D. -2, 1

Answer: A



[Watch Video Solution](#)

104. If $x^2 + 4y^2 - 8x + 12 = 0$ is satisfied by real values of x and y then y must lie between

A. 2, 6

B. 2, 5

C. -1, 1

D. $-2, 1$

Answer: C



Watch Video Solution

105. Let $f(x)$ be a polynomial for which the remainders when divided by $x - 1, x - 2, x - 3$ respectively are 3, 7, 13 then the remainder of $f(x)$ when divided by $(x - 1)(x - 2)(x - 3)$ is

A. $f(x)$

B. $x^2 + X + 1$

C. $x^2 + 1$

D. none

Answer: B



View Text Solution

106. Let two numbers have arithmetic mean 9 geometric mean 4. then these numbers are the roots of the quadratic equation

A. $x^2 + 18x + 16 = 0$

B. $x^2 - 18x - 16 = 0$

C. $x^2 + 18x - 16 = 0$

D. $x^2 - 18x + 16 = 0$

Answer: D



[Watch Video Solution](#)

EXERCISE 2 (SPECIAL TYPE QUESTIONOS) (SET -1)

1. I : If $(a\alpha + b)^2 + (a\beta + b)^{-2} = 1$ where α, β are the roots of $ax^2 + bx + c = 0$ then $ac(ac + 2) = b^2$

II : the value of 'a' for which one root of the quadratic equation

$(a^2 - 5a + 3)x^2 + (3a - 1)x + 2 = 0$ is twice as large as the quadratic equation

- A. Only I is true
- B. Only II is true
- C. both I and II are true
- D. niether I nor II true

Answer: A



[View Text Solution](#)

2. I: If $p(q - r)x^2 + q(r - p)x + r(p - q) = 0$ has equal roots then p,q,r in A.P

II : if the sum of the roots of $ax^2 + bx + c = 0$ has equal to the sum of the squares of their reciprocals then $bc^2, ca^2, + bx + c = 0$ is equal to the sum of the squares of their reciprocals then bc^2, ca^2, ab^2 are in A.P

- A. Only I is true

B. Only II is true

C. both I and II are true

D. neither I nor II true

Answer: B



[View Text Solution](#)

3. I) The maximum value of $c + 2bx - x^2$ is $c + b^2$

II) The minimum value of $x^2 + 2bx + c$ is $c - b^2$

Which of the above statements is true ?

A. Only I is true

B. Only II is true

C. both I and II are true

D. neither I nor II true

Answer: A



Watch Video Solution

4. E_1 , $a + b + c = 0$ if 1 is a root of $ax^2 + bx + c = 0$.

E_2 : $b^2 - a^2 = 2ac$ if $\sin \theta \cos \theta$ are the of $ax^2 + bx + c = 0$

which of the following is true?

- A. E_1 is true E_2 is true
- B. E_1 is true E_2 is false
- C. E_1 is false, E_2 is ture
- D. E_1 is false E_2 is false

Answer: A



View Text Solution

EXERCISE 2 (SPECIAL TYPE QUESTIONOS) (SET -2)

1. $\alpha, \beta, \gamma \in \mathbb{R}$ is a root of $a^2x^2 + bx + c = 0$ β is a root of $a^2x^2 - bx - c = 0$ and γ is a root of $a^2x^2 + 2bx + 2c = 0$ then

A. A,B,C,D

B. B,D,C,A

C. A,C,B,D

D. D,B,A,C

Answer: B



[Watch Video Solution](#)

2. if A, B, C, D are the sum of the roots of the roots of $2x^2 + x + 3 = 0, x^2 - x + 2 = 0, 3x^2 - 2x + 1 = 0, x^2 - x - x + 1 = 0$

then the ascending order of A,B,C,D is

A. A,B,C,D

B. B,D,C,A

C. A,C,B,D

D. C,D,A,B

Answer: C



[Watch Video Solution](#)

3. IF A, B, C , are the minimum value of $x^2 - 8x + 17$, $2x^2 + 4x - 5$, $3x^2 - 7x + 1$ then the ascending order of A, B, C is

A. A,B,C

B. B,C,A

C. C,A,B

D. A,C,B

Answer: B



[Watch Video Solution](#)

4. IF A,B,C are the maximum value of $2x + 5 - x^2$, $x - 1 - 2x^2$, $5x + 2 - 3x^2$ then the descending order of A,B,C is

A. A,B,C

B. B,C,A

C. C,A,B

D. A,C,B

Answer: D



[Watch Video Solution](#)

EXERCISE 2 (SPECIAL TYPE QUESTIONOS) (SET -3)

1. If α, β are the roots of $ax^2 + bx + c = 0$ then match the following

I. $\alpha\beta^2 + \alpha^2\beta + \alpha\beta =$

a) $\frac{b^2 - 2ac}{c^2}$

II. $\frac{1}{\alpha^2} + \frac{1}{\beta^2} =$

b) $\frac{3abc - b^3}{a^2 c}$

III. $\frac{\alpha^2}{\beta} + \frac{\beta^2}{\alpha} =$

c) $\frac{c^3}{a^3}$

IV. $\frac{\alpha^3 + \beta^3}{\alpha^{-3} + \beta^{-3}} =$

d) $\frac{ac - bc}{a^2}$

A. c,d,a,b

B. d,c,b,a

C. d,a,b,c

D. c,b,a,c

Answer: C



Watch Video Solution

2. Match the following

- I. If α, β are the roots of $x^2 + 5x - 4 = 0$ then the equation whose roots are $\frac{\alpha+2}{3}, \frac{\beta+2}{3}$ is a) $4x^2 + 3x - 2 = 0$
- II. If α, β are the roots of $2x^2 + x + 3 = 0$ then the equation whose roots are $\frac{1-\alpha}{1+\alpha}, \frac{1-\beta}{1+\beta}$ is b) $3x^2 - 2x + 1 = 0$
- III. If α, β are the roots of $x^2 - 2x + 3 = 0$ then the equation whose roots are $\frac{\alpha-1}{\alpha+1}, \frac{\beta-1}{\beta+1}$ is c) $9x^2 + 3x - 10 = 0$
- IV. If α, β are the roots of $2x^2 + 3x - 4 = 0$ then the equation whose roots are $2\alpha + \frac{3}{\beta}, 2\beta + \frac{3}{\alpha}$ is d) $2x^2 + x + 3 = 0$

A. a,c,d,b

B. d,c,b,a

C. b,c,d,a

D. c,d,a,c

Answer: D



View Text Solution

3. match the following .

Inequation

I. $x^2 - 7x + 12 \geq 0$

II. $x^2 + 3x - 4 \leq 0$

III. $2x^2 + 3x < 2$

IV. $x^2 - 2x + 2 < 0$

Solution set

a) \emptyset

b) $(-2, 1/2)$

c) $[-4, 4]$

d) $(-\infty, 3] \cup [4, \infty)$

A. a,c,d,b

B. d,c,b,a

C. b,c,d,a

D. c,d,a,c

Answer: B



Watch Video Solution

4. Let α and β be the roots of the quadratic equation $ax^2 + bx + c = 0$

. Observe the lists given below:

List -I

i) $\alpha = \beta \Rightarrow$

ii) $\alpha = 2\beta \Rightarrow$

iii) $\alpha = 3\beta \Rightarrow$

iv) $\alpha = \beta^2 \Rightarrow$

List-II

A) $(ac^2)^{1/3} + (a^2c)^{1/3} + b = 0$

B) $2b^2 = 9ac$

C) $b^2 = 6ac$

D) $3b^2 = 16ac$

E) $b^2 = 4ac$

F) $(ac^2)^{1/3} + (a^2c)^{1/3} = b$

Let correct match of List -I from List -II is

A. $I \quad ii \quad iii \quad iv$
 $E \quad B \quad D \quad F$

B. $I \quad ii \quad iii \quad iv$
 $E \quad B \quad A \quad D$

C. $I \quad ii \quad iii \quad iv$
 $E \quad D \quad B \quad F$

D. $I \quad ii \quad iii \quad iv$
 $E \quad B \quad D \quad A$

Answer: D



[View Text Solution](#)

EXERCISE 2 (SPECIAL TYPE QUESTIONOS) (SET -4)

1. A: The quadratic equation having roots $3 \pm 2I$ is $x^2 - 6x + 13 = 0$

R : the quadratic equation having roots α, β is $x^2 - (\alpha + \beta)x + \alpha\beta = 0$

A. Both A and R are true and R is the correct explanation of A

B. Both A and R are true but R is not the correct explanation of A

C. A is true but R is false

D. A is false but R is true

Answer: A

 [Watch Video Solution](#)

2. A: the quadratic equation whose roots are the reciprocals of the roots of $3x^2 - 7x + 2 = 0$ is $3x^2 + 7x + 2 = 0$

R: the quadratic equation whose roots are the reciprocals of the roots of the quadratic equation $f(x)$ is $f(1/x) = 0$

A. Both A and R are true and R is the correct explanation of A

B. Both A and R are true but R is not the correct explanation of A

C. A is true but R is false

D. A is false but R is true

Answer: D



Watch Video Solution

3. A : $x^2 + x + 1 > 0$ for all $x \in R$

R : if the roots of $ax^2 + bx + c = 0$ are imaginary then for $x \in R$, $ax^2 + bx + c$ and a have the same sign .

- A. Both A and R are true R is the correct explanation of A
- B. Both A and R are true but R is not correct explanation of A
- C. A is true but R is false
- D. A is false but R is true

Answer: A



View Text Solution

4. A : $x^2 + x - 12 \leq 0 \Rightarrow x \in [-4, 3]$

R : if α, β are the roots of $ax^2 + bx + c = 0$, $\alpha < \beta$ then $\alpha < x < \beta \Leftrightarrow ax^2 + bx + c$ and a have opposite signs .

- A. Both A and R are true R is the correct explanation of A
- B. Both A and R are true but R is not correct explanation of A
- C. A is true but R is false
- D. A is false but R is true

Answer: A



[View Text Solution](#)