Class 11 Physics

Motion In A Plane

Ncert Video Solutions - Only On Doubtnut

Download Doubtnut Now

Exercise 4.11

State, for each of the following physical quantities, if it is a scalar or a vector. Volume, mass speed acceleration, density, number of moles, velocity, angular frequency, displacement, angular velocity.

Watch Video Solution On Doubtnut App

Exercise 4.12

Pick out the two scalar quantities in the following list: force, angular momentum, work, current, linear momentum, electric field, average force, angular momentum, work, current , linear momentum, electric field, average velocity, magnetic moment,

relative velocity.

Watch Video Solution On Doubtnut App

Exercise 4.13

Pick out the only vector quantity in the following list:

Temperature, pressure, impulse, time, power , total path length, energy, gravitational tempeature, pressure, impulse, time, power, total path length, energy, gravitational potential, coefficent of friction, charge.

Watch Video Solution On Doubtnut App

Exercise 4.14

State with reasons, whether the following algebraic operations with scalar and vector physical quantities are meaningful:

(i) Adding any two scalars (ii) Adding a scalar to a vector of the

same dimensions

(iii) Multiplying any vector by any scalar (iv) Multiplying any two scalars

(v) Adding any two vectors

Watch Video Solution On Doubtnut App

Exercise 4.15

Read each statement below carefully and state with reasons, with it is true or false :

(a) The magnitude of vector is always a scalar.

(b) Each component of a vector is always a scalar.

(c) The total path length is always equal to the magnitude of the displacement vector of a particle.

(d) The average speed of a particle (defined as total path length divided by the time taken to cover the path) is greater or equal to the magnitude of average velocity of the particle over the same interval

of time.

(e) three vectors not lying in a plane can never add up to give a null vector.

Watch Video Solution On Doubtnut App

Exercise 4.16

Establish the following inequalities geometrically or otherwise ,

(a) $\left|\vec{A}+\vec{B}\right| \leq \left|\vec{A}\right| + \left|\vec{B}\right|$,

(b) $\left|\vec{A}+\vec{B}\right| \geq \left|\left|\vec{A}\right| - \left|\vec{B}\right|\right|$

(C) $\left|\vec{A}-\vec{B}\right| \leq \left|\vec{A}\right| + \left|\vec{B}\right|$

(d) $\left|\vec{A}-\vec{B}\right| \geq \left|\left|\vec{A}\right| - \left|\vec{B}\right|\right|$

When does the equality sign above apply ?

Watch Video Solution On Doubtnut App

Exercise 4.17

Given $\vec{A} + \vec{B} + \vec{C} + \vec{D} = 0$, which of the following statements are correct ?

(a) $\vec{A}, \vec{B}, \vec{C}$ abd $\vec{D}$ each must be a null vector.

(b) The magnitude of $\left(\vec{A} + \vec{C}\right)$ equals the magnitude of $\left(\vec{B} + \vec{D}\right)$.

(c) The magnitude of $\vec{A}$ can never be greater than the sum of the magnitude of $\vec{B}, \vec{C}$ and $\vec{D}$.


(d) $\vec{B} + \vec{C}$ must lie in the plane of $\vec{A} + \vec{D}$. if $\vec{A}$ and $\vec{D}$ are not collinear and in the line of $\vec{A}$ and $\vec{D}$, if they are collinear.

Watch Video Solution On Doubtnut App

Exercise 4.18

Three girls skating on a circular ice ground of radius $200m$ start from a point (P) on the edge of the ground and reach a point Q diametrically opposite to (P) following different paths as shown in Fig. (NCT) . 17. What is the magnitude of the displacement vector

for each ? which girl's displacement is equal to the actual length of path skated ?




Watch Video Solution On Doubtnut App

Exercise 4.19

A cyclist starts from the centre O of a circular park of radius 1km,


reaches the edge P of the park, then cycles along the PQ cicumference and returns to the centre along OQ as shown in fig. If the round trip taken ten minute, the net displacement and average speed of the cylists (in kilometer and kinetic per hour) is



Watch Video Solution On Doubtnut App

Exercise 4.20

On an open ground, a motorist follows a track that turns to his left by an angle of 60 after every $500m$. Starting from a given turn, The path followed by the motorist is a regular hexagon with side $500m$, as shown in the given figure specify the displacement of the motorist



at the end of sixth turn.

Watch Video Solution On Doubtnut App

Exercise 4.21

A passenger arriving in a new town wishes to go from the station to a hotel located 10km away on a straight road from the station. A dishonest cabman takes him along a circuitous path 23km long and reaches the hotel in 28 minutes . What is (a) the average speed of the taxi, (b) the magnitude of average velocity ? Are the two equal ?

Watch Video Solution On Doubtnut App

Exercise 4.22

Rain is falling vertically with a speed of $30ms^{-1}$. A woman rides a bicycle with a speed of $10ms^{-1}$ in the North to South direction.

What is the direction in which she should hold her umbrella ?

Watch Video Solution On Doubtnut App

Exercise 4.23

A man can swim with a speed of $4kmh^{-1}$ in still water. He crosses a river 1km wise that flows steadly at $3kmh^{-1}$. If he makes his strokes normal to the river current, how far down the river does he go when he reaches the other bank?

Watch Video Solution On Doubtnut App

Exercise 4.24

In a harbour, wind is blowing at the speed of $72km/h$ and the flag on the mast of a boat anchored in the harbour flutters along the N-E *direction*. If the boat starts moving at a speed of $51km/h$ to the North, what is the direction of flag on the mast of the bat ?

Watch Video Solution On Doubtnut App

Exercise 4.25

The ceiling of a long hall is $25m$ high. What is the maximum horizontal distance if a ball thrown with a speed of $40ms^{-1}$ can go without hitting the ceiling of the hall ?

Watch Video Solution On Doubtnut App

Exercise 4.26

A cricketer can throw a ball to a maximum horizontal distance of 100m. With the same speed how much high above the ground can the cricketer throw the same ball?

Watch Video Solution On Doubtnut App

Exercise 4.27

A stone tied to end of a string $80cm$ long is whirled in a horizontal

plane with a constant speed. If the stone makes 14 revolutions in 25 seconds, what is the magnitude and direction of acceleration of the stone ?

Watch Video Solution On Doubtnut App

Exercise 4.28

An aircraft executes a horizontal loop of radius 1 km with a steady speed of $900 kmh^{-1}$. Compare its centripetal acceleration with the acceleration due to gravity.

Watch Video Solution On Doubtnut App

Exercise 4.29

Read each statement below carefully and state, with reasons, if it is true or false : (a) The net acceleration of a particle in circular motion is always along the radius of the circle towards the centre.

(b) The velocity vector of a particle at a point is always aling the tagent to the path of the partile at that point. ltbRgt (c) The acceleration vector of a particle in uniform circular motion acraged over one cycle is a null vector.

Watch Video Solution On Doubtnut App

Exercise 4.30

The position of a particle is given by $\vec{r} = 3.0t\hat{i}$ - $2.0t^2\hat{j}$ + $4.0\hat{k}m$

$\overrightarrow{r} = 3.0t\hat{i} - 2.0t^2\hat{j}$

$+ 4.0\hat{k}m$

, wher (t) in seconds and the coefficients have the proper units for $\vec{r}$ to be in metres. (a) Fing the $\vec{v}$ and $\vec{a}$ of the particle ? (b) What is the magnitude and direction fo velocity of the particle at $t = 2s$?

Watch Video Solution On Doubtnut App

Exercise 4.31

A particle starts from the origin at $t = 0$ with a velocity of $10.0\hat{j}m/s$ and moves in the X-y plane with a constant accleration of $\left(8.0\hat{I} + 2.0\hat{j}\right)ms^{-2}$. (a) At wht time is the x-coordinate of the particle $16m$? What is the y-coordinate of the particle at that time ? (b) What is the speed of the particle at that time ?

Watch Video Solution On Doubtnut App

Exercise 4.32

$\hat{i}$ and $\hat{j}$ are unit vectors along x-and y-axes respectively. What is the magnitude and the direction of the vectors $\hat{i} + \hat{j}$ and $\hat{i} - \hat{j}$? What are the components of a vector $\vec{A} = 2\hat{i} + 3\hat{j}$ along the direction $\hat{i} + \hat{j}$and $\hat{i} - \hat{j}$?

Watch Video Solution On Doubtnut App

Exercise 4.33

For any arbitary motion in space, which of the following relations are true?

a) $v_{\text{average}} = (1/2)\left(v\left(t_1 + v\left(t_2\right)\right)\right.$

b) $v_{\text{average}} = \left[r\left(t_2\right) - r\frac{t_1}{t_2 - t_1}\right.$

$v(t) = v(0) + at$

d) $a_{\text{average}} = \left[v\left(t_2\right) - v\frac{t_1}{t_2 - t_1}\right.$

The average stands for average of the quantity over time interval t_1 to t_(2)`

Watch Video Solution On Doubtnut App

Exercise 4.34

Read each statement below carefully and state with reason and examples, if it is true or false. A scalar quantity is one that (a) is conserved in a process (b) can never take negative values (c) must

be dimensionless (d) does not vary from one point to another in space (e) has the same value for observers with different orientations of axes.

Watch Video Solution On Doubtnut App

Exercise 4.35

An aircraft is flying at a height of 3400 m above the ground. If the angle subtended at a ground observation point by the aircraft positions 10 s apart is 30 , what is the speed of the aircraft ?

Watch Video Solution On Doubtnut App

Exercise 4.36

A vector has magnitude and direction (i) Does it have a location in the space ? (ii) Can it vary with time ? (iii) Will two equal vectors $\vec{a}$ and $\vec{b}$ different locations in space necessarily have identical

physical effects ? Give examples in support of your answer.

Watch Video Solution On Doubtnut App

Exercise 4.37

A vector has both magnitude and direction. Does that mean anything that has magnitude and direction is necessarily a vector ? The rotation of a body can specified by the direction of the axis of rotation and the angle of rotation about the axis. Does the make any rotation a vector ?

Watch Video Solution On Doubtnut App

Exercise 4.38

Can you associate vectors with (a) the length of a wire bent into a loop (b) a plane area (c) a sphere ? Explain.

Watch Video Solution On Doubtnut App

Exercise 4.39

A bullet fired at an angle of 30 with the horizontal hits the ground 3.0 km away. By adjusting its angle of projection, can one hope to hit a target 5.0 km away? Assume the muzzle speed to be fixed and neglect air resistance.

Watch Video Solution On Doubtnut App

Exercise 4.40

A fighter plane flying horizontally at an altitude of $1.5km$ with speed $720kmh^{-1}$ passes directly over head an anticraft gun. At what angle from the vertical should the gun be fired from the shell with muzzle speed $600ms^{-1}$ to hit plane.

At what minimum altitude should the pilot fly the plane to avoid

being hit ? (Take g= 10 ms^{-2}).

Watch Video Solution On Doubtnut App

Exercise 4.41

A cyclist is riding with a speed of $27kmh^{-1}$. As he approaches a circular turn on the road of radius 80 m, he applies brakes and reduces his speed at the constant rate $0.5ms^{-1}$. What is the magnitude and direction of the net acceleration of the cyclist on the circular turn ?

Watch Video Solution On Doubtnut App

Exercise 4.42

Show that for a projectile the angle between the velocity and the x-axis as function of time is given

by $\theta_t = \tan^{-1}\left(\frac{4h_m}{R}\right)$

where the sybols have thir usual meanings.

Watch Video Solution On Doubtnut App


Solved Example 1

Rain is falling vertically with a speed fo $35ms^{-1}$. Winds starts blowing after sometime with the speeds of $12ms^{-1}$ in east to west direction. At what angles with the vertical should a boy waiting at a bus stop hold his umbrella to protect himself from rain?

Watch Video Solution On Doubtnut App

Solved Example 2

Find the magnitude and direction of the resultant of two vectors A and B in the terms of their magnitudes and angle θ between them.



Watch Video Solution On Doubtnut App

Solved Example 3

A motor boat is racing towards North at $25km/h$ and the water current in that region is $10km/h$ in the direction of 60^{00} East of South. Find the resultant velocity of the boat.

Watch Video Solution On Doubtnut App

Solved Example 4

The position of a particle is given by

$r = 3t\hat{i} + 2t^2\hat{j} + 8\hat{k}$

where, t is in seconds and the coefficients have the proper units for r

to be in metres.

(i) Find v (t) and a(t) of the particles.

(ii) Find the magnitude and direction of v(t) and a(t) at $1 = 1s$.

Watch Video Solution On Doubtnut App

Solved Example 5

A paarticle starts from origin at $t = 0$ with a velocity $5.0\hat{i}m/s$ and

moves in x-y plane under action of a force which produces a

constant acceleration of $\left(3.0\hat{i} + 2.0j\right)m/s^2$.

(a) What is the y-cordinate of the particle at the instant its x-

coordinate is `84 m ? (b) What is the speed of the particle at this

time?

Watch Video Solution On Doubtnut App

Solved Example 6

Rain is falling vertically with a speed of $35ms^{-1}$. A woman rides a bicyle with a speed of $12ms^{-1}$ in east to west direction. What is the direction in which she should hold her umbrealla?

Watch Video Solution On Doubtnut App

Solved Example 7

Galileo, in his book Two new sciences, stated that "for elevations which exceed or fall short of 45 by equal amounts, the ranges are equal. Prove this statement.

Watch Video Solution On Doubtnut App

Solved Example 8

A hiker stands on the edge of a cliff $490m$ above the ground and throwns a stone horiozontally with an initial speed of $15ms^{-1}$ neglecting air resistance.The time taken by the stone to reach the ground in seconds is $\left(g = 9.8ms^{2}\right)$

Watch Video Solution On Doubtnut App

Solved Example 9

A cricket ball is thrown at a speed of $28ms^{-1}$ in a direction 30 above the horizontal. Calculate (a)the maximum height (b) the time taken by ball to return ti the same level, and (c)the distance from the thrower to the point wher the ball restance from the throwrt to the point where the ball returns to the same level.

Watch Video Solution On Doubtnut App

Solved Example 10

An insect trapped in a circular groove of radius 12 cm moves along the groove steadily and completes 7 revolutions in 100 s. (i) What is the angular speed and the linear speed of the motion ? (ii) Is the acceleration vector a constant vector ? What is its magnitude ?


Watch Video Solution On Doubtnut App

Apne doubts ka Instant video solution paayein

Abhi Doubtnut try karein!

GET IT ON Google Play

Whatsapp your doubts on 8400400400



doubtnut